

OMICS GROUP



OMICS Group International through its Open Access Initiative is committed to make genuine and reliable contributions to the scientific community. OMICS Group hosts over **400** leading-edge peer reviewed Open Access Journals and organizes over **300** International Conferences annually all over the world. OMICS Publishing Group journals have over **3 million** readers and the fame and success of the same can be attributed to the strong editorial board which contains over **30000** eminent personalities that ensure a rapid, quality and quick review process. OMICS Group signed an agreement with more than **1000** International Societies to make healthcare information Open Access.

OMICS Journals are welcoming Submissions

OMICS Group welcomes submissions that are original and technically so as to serve both the developing world and developed countries in the best possible way.

OMICS Journals are poised in excellence by publishing high quality research. OMICS Group follows an Editorial Manager® System peer review process and boasts of a strong and active editorial board.

Editors and reviewers are experts in their field and provide anonymous, unbiased and detailed reviews of all submissions.

The journal gives the options of multiple language translations for all the articles and all archived articles are available in HTML, XML, PDF and audio formats. Also, all the published articles are archived in repositories and indexing services like DOAJ, CAS, Google Scholar, Scientific Commons, Index Copernicus, EBSCO, HINARI and GALE.

For more details please visit our website:

<http://omicsonline.org/Submitmanuscript.php>



Donetsk National Medical University named after M. Gorky


DISODERS OF PURINE AND PYRIMIDINE NUCLEOTIDES METABOLISM IN HUMAN GASTROINTESTINAL TRACT CANCER

Author: Doctor of Biology Science in the field of Oncology,

Professor of the Chemistry Department of DonNMU **BORZENKO BERTA GEORGIYEVNA**

Co-authors: associate prof. Elena Bakurova, assistant Kseniia Mironova

Stomach cancer is the fourth most common cancer in the world



Stomach cancer


Stomach cancer is the fourth most common cancer in the world, with 989,000 new cases diagnosed in 2008 [WCRF International].

Key enzymes of thymidine metabolism

Thymidine kinase (TK; EC 2.1.2.1.)


Thymidine phosphorylase (TP; EC 2.4.2.4.)

TP is similar to the platelet-derived endothelial cell growth factor (PD-ECGF) and hence plays dual role in cell biology




DE NOVO

The activity of the pathway
in norm 90-95%
In tumor ~80% ↓



Simple precursors



Thymidine
synthase

FU

SALVAGE

The activity of the pathway
in norm 5-10%
In tumor ~20% ↑

TP



TK


Total pool

dTDP

dTTP

Cancer
feed back
mechanism is
absent






The thymidine salvage pathway enzymes TK and TP compete for thymidine as a substrate and catalyze opposing synthetic and catabolic reactions that have been implicated in the control of proliferation and angiogenesis, respectively.

Key enzymes of adenosine (Ado) metabolism

Ecto-5'-nucleotidase (5'-NT AMP) or CD73 (5'-NT: 3.1.3.5)
Adenosine deaminase (ADA: 3.5.4.4)



The aim of this study

is to investigate the DNA precursors metabolism in healthy and patients with gastric cancer $T_{3-4}N_{0-x}M_0$ and to determine the possible role of its regulatory enzymes as biomarkers.

PATIENTS AND METHODS

- ✗ Seventy-four adult patients, 35–80 year old, with gastric cancer were included in this study.
- ✗ Patients were staged in the accordance with TNM classification (the 6th edition) as $T_{3-4}N_{0-x}M_0$.
- ✗ Eighty-two non-cancerous patients of the same age without gastroduodenal zone pathology formed control group and also were examined.
- ✗ The study protocol was approved by Ethical Committee permission of M. Gorky National Medical University (Donetsk, Ukraine) for studies with human materials.
- ✗ The features of enzymes activity were studied in blood serum and tissues. Activity of tissue enzymes was determined in the surgically removed material in thirty tumors (the mucosa margin of resection distant from the carcinoma had being as a control).


PATIENTS AND METHODS

- × TK activity was determined by the radioisotope method. Radioactivity of diethyl-amino-ethyl-cellulose paper disks (DEAE-cellulose disks paper "Limbro" (UK) was counted in a liquid scintillation SL-8 on a biological scintillation counter (CBS-2, Russia). TK activity was given in nanomoles of thymidylate formed within an hour per 1 mg of protein.
- × TP, ADA [Fritch], 5`NT [Edwards. Clin. Path. Immunology,1980] activity was determined spectrophotometrically.
- × Protein concentration in biological material was determined using the method described by Lowry et al.
- × Statistical analysis of results was performed using Medstat software package. On significance of differences was analyzed by parametric and nonparametric methods.

Miszczak-Zaborska E., Smolarek M., Bartkowiak J. Influence of the thymidine phosphorylase (platelet-derived endothelial cell growth factor) on tumor angiogenesis. Catalytic activity of enzyme inhibitors// Postery Biochem. – 2010. - 56(1). – P. 61-66.

Tertil M., Skrzypek K., Flirczyk U. et al. Regulation and Novel Action of Thymidine Phosphorylase in Non-Small Cell Lung Cancer: Crosstalk with Nrf2 and HO-1 //PLoS One. - 2014. - 9(5): e97070.

TP and TK activity in the serum of control group, patients with gastric cancer and in tissues, nmol/mg[•]min and nmol/mg[•]h, respectively




B.G. Borzenko, E.M. Bakurova, Yu.A. Popovich, et al. Activity of thymidilate “salvage pathway” enzymes in human gastric cancer and blood serum: correlation with treatment modalities// Exp.Oncol. – 2013. – Vol. 35, №1. – P. 37 – 40


Sho Tabada, Ryuji Ikeda, Masatatsu, Sunii Shimaoka et al. Thymidine phosphorylase activates NFkB and stimulates the expression of angiogenic and metastatic factors in human cancer cells// Oncotarget. - 2014. - 5(21). – P. 10473 – 85

TK and TP activity in blood serum of control group in dependence of age

The thymidine kinase activity



The thymidine phosphorylase activity




The activity of enzymes, which regulates metabolism of adenosine in blood serum of control group and gastric cancer patients

*Regateiro FS, Cobbold SP, Waldmann H. CD73 and adenosine generation in the creation of regulatory microenvironments.

Clin Exp Immunol. 2013. Sir William Dunn School of Pathology, University of Oxford, Oxford, UK.



*Zimmermann H, Zebisch M, Sträter N. Cellular function and molecular structure of ectonucleotidases. Purinergic Signal. 2012. Institute of Cell Biology and Neuroscience, Molecular and Cellular Neurobiology, Biologicum, Goethe-University Frankfurt, Max-von-Laue-Str. 13, 60438, Frankfurt am Main, Germany.

ADA and 5'-NT activity in blood serum of control group in dependence of age




•Borzenko B., Bacurova H., Verhova O., Zhebelenko Y. Can we used age-dependent changes of enzymes activity in benign disease as poor prognostic factors? //Joint ECCO 15-34 ESMO Multidisciplinary Congress Abstract book. - Vol.7, N 2, September 2009. – P. 313


TP and TK activity of gastric cancer patients in different age groups



The activity of enzymes, which regulates metabolism of adenosine in blood serum of control group and gastric cancer patients




ADA activity in blood cells of gastic cancer patients



Sauer AV, Brigida I, Carriglio N, Hernandez RJ, Scaramuzza S, et.al. Alterations in the adenosine metabolism and CD39/CD73 adenosinergic machinery cause loss of Treg cell function and autoimmunity in ADA-deficient SCID. Blood. 2012. San Raffaele Telethon Institute for Gene Therapy (HSR-TIGET), Via Olgettina 58, Milan, Italy


Analysis of lymphocytes and erythrocytes functional power

The blast transformation reactions of lymphocytes



The decline of cellular immune function


Sorption capacity of erythrocyte membrane



Disorder of transport function of erythrocytes

Regulatory enzymes of adenosine metabolism in blood serum and lymphocytes of gastric cancer patients

Age, years	ADA, nmol/mg*min				5`-NT, nmol/mg*min			
	Healthy		Oncology patients		Healthy		Oncology patients	
	serum	lymphocytes	serum	lymphocytes	serum	lymphocytes	serum	lymphocytes
men								
36-45	1,83 ±0,14	182,4± 9	6,92 ±0,27	94,2 ±1,1	5,48 ±0,52	50,3 ±3,1	5,2± 0,33	25,4± 0,2
46-60	1,74 ±0,22	178,4± 2,1	9,46 ±2,6	89,6 ±3,6	5,76 ±0,26	48 ±5,7	10±0,5	36,8± 0,9
61-70	2,18 ±0,45	100,7± 7,2	8,98 ±1,2	73,4 ±3,9	5,63 ±0,46	45,1 ±2,1	10,3±0 ,6	29±1,9




The association between serum enzymes activity and volume of surgical intervention

Enzymes activity	The type of surgery					
	radical		diagnostic		palliative	
	before	after 9 days	before	after 9 days	before	after 9 days
TK, nmol/mg·h	6,29 ± 1,47	3,40 ± 0,58*	6,43 ± 1,76	4,02 ± 0,45	5,50 ± 0,67	1,97 ± 0,20*
TP, nmol/mg·min	15,28 ± 2,1	38,33 ± 1,9**	10,03 ± 2,4	11,43 ± 2,2	13,0 ± 1,8	14,52 ± 1,7*
ADA, nmol/mg·min	7,19 ± 2,9	5,95 ± 0,6	7,45 ± 2,2	6,57 ± 1,3	10,31 ± 2,5	10,96 ± 1,5
5`-NT, nmol/h·mg	63,02 ± 8,1	40,58 ± 6,3*	51,48 ± 2,1	31,52 ± 3,9*	48,62 ± 6,5	38,86 ± 5,2


Note: authentic differences versus previous value: *- p < 0.05, ** - p < 0.01

Comparative dynamic of thymidine kinase and thymidine phosphorylase activity during treatment

Radical type of surgery



Palliative type of surgery



References

B.G. Borzenko, E.M. Bakurova, Yu.A. Popovich, et all. Activity of thymidilate “salvage pathway” enzymes in human gastric cancer and blood serum: correlation with treatment modalities// *Exp.Oncol.* – 2013. – Vol. 35, №1. – P. 37 – 40

Regateiro FS, Cobbold SP, Waldmann H. CD73 and adenosine generation in the creation of regulatory microenvironments. *Clin Exp Immunol.* 2013. Sir William Dunn School of Pathology, University of Oxford, Oxford, **UK.**

Sauer AV, Brigida I, Carriglio N, Hernandez RJ, Scaramuzza S, et.all. Alterations in the adenosine metabolism and CD39/CD73 adenosinergic machinery cause loss of Treg cell function and autoimmunity in ADA-deficient SCID. *Blood.* 2012. San Raffaele Telethon Institute for Gene Therapy (HSR-TIGET), Milan, **Italy**

Zimmermann H, Zebisch M, Sträter N. Cellular function and molecular structure of ecto-nucleotidases. *Purinergic Signal.* 2012. Institute of Cell Biology and Neuroscience, Molecular and Cellular Neurobiology, Biologicum, Goethe-University Frankfurt, Frankfurt am Main, **Germany.**

Ghiringhelli F, Bruchard M, Chalmin F, Rébé C. Production of adenosine by ectonucleotidases: a key factor in tumor immunoescape. *J Biomed Biotechnol.* 2012. Faculté de Médecine, Université de Bourgogne, Dijon, **France.**

Jin D, Fan J, Wang L, Thompson LF, Liu A, Daniel BJ, Shin T, Curiel TJ, Zhang B. CD73 on tumor cells impairs antitumor T-cell responses: a novel mechanism of tumor-induced immune suppression. *Cancer Res.* 2010. Cancer Therapy and Research Center, Department of Medicine, University of Texas Health Science Center, **USA.**

E. Mischak-Zaborska, R. Kubiak, A. Bienkiewicz et al. The cytosol activity of thymidine phosphorylase in endometrial cancer // *J. of Experimental and Cancer Research*. – 2008. - Vol. 27 – P. 64 – 69.

A. Kawahara, Y. Akagi, S.Hattori et al. Higher expression of deoxyuridine triphosphatase may predict the metastasis potential of colorectal cancer/ // *J. Clin. Phathology*. – 2009. - Vol. 62, № 4. – P. 364 - 369.

Peter P. Grimminger, Paul M. Schneider, Ralf Metzger et al. Low Thymidylate Synthase, Thymidine Phosphorylase, and Dihydropyrimidine Dehydrogenase mRNA Expression Correlate With Prolonged Survival in Resected Non–Small-Cell Lung Cancer// *Clinical Lung Cancer*. – 2010.- Vol. 11, № 5. – P. 328-334.

Zhang J.M., Mizoi T., Shiiba K. et al. Expression of thymidine phosphorylase by macrophages in colorectal cancer tissues // *World J. Gastroenterol*. – 2004. - Vol.10, №4. – P. 545– 549.

Daozhen Chen, Qiusha Tang. An experimental study on cervix cancer with combination of HSV-TK/GCV suicide gene therapy system and 60Co radiotherapy // *BMC Cancer*. – 2010. - №10. – P. 609- 614.

Sam J. Lubner, Noelle K. LoConte, Kyle D. Holen et al. A Phase II Study of Oxaliplatin, 5-Fluorouracil, Leucovorin, and High-Dose Capecitabine in Patients With Metastatic Colorectal Cancer // *Clinical Colorectal Cancer*. – 2010.- Vol. 9, № 3. – P. 157-161.


T. Hotta, Y. Kobayashi, K. Taniguchi et al. Thymidine phosphorylase and dihydropyrimidine dehydrogenase expression in hepatocellular carcinoma and metastatic liver cancer // *Oncol. Rep*. – 2004. – Vol.12, №2. – P. 347 –351.

Sho Tabada, Ryuji Ikeda, Masatatsu , Sunii Shimaoka et al. Thymidine phosphorylase activates NFkB and stimulates the expsression of angiogenic and metastatic factors in human cancer cells// *Oncotarget*. - 2014. - 5(21). – P. 10473 – 85

Conclusions

An activity of TK, TP, ADA and 5`-NT has an effect on the tumor disorders.

The control of individual dynamics of this enzymes activity in blood serum of gastric cancer patients may be used as informative tool for monitoring of patients and treatment optimization.





**Thank
for your attention!**

Prof. Borzenko Berta
Georgiyevna

E-mail: borzenko.bg@rambler.ru

Metabolomics: Open Access Related Journals

- Drug Metabolism & Toxicology
- Diabetes & Metabolism
- Cancer Science & Therapy
- Carcinogenesis & Mutagenesis
- Metabonomics & Metabolites
- Metabolomics: Open Access



Metabolomics: Open Access Related Conferences

- 4th International Conference and Exhibition on Metabolomics & Systems Biology 2015



OMICS Group Open Access Membership

OMICS publishing Group Open Access Membership enables academic and research institutions, funders and corporations to actively encourage open access in scholarly communication and the dissemination of research published by their authors.

For more details and benefits, click on the link below:

<http://omicsonline.org/membership.php>

