

# Bio-Suppression of Strawberry Fruit Rot Disease Caused by *Botrytis cinerea*

Abeer A El-ghanam<sup>1</sup>, Safinaz A Farfour<sup>2\*</sup> and Seham S Ragab<sup>1</sup><sup>1</sup>Plant Pathology Research Institute, Agricultural Research Center, Giza, Egypt<sup>2</sup>Environmental Biotechnology Department, Genetic Engineering and Biotechnology Research Institute (GEBRI), University of Sadat City, Egypt

## Abstract

In grade to obtain high quality and safety food along with the environment protection from pesticides hazards, four bio agents were used to control fruit rot disease caused by *Botrytis cinerea* in strawberry fruits. *Chlorella vulgaris*, *Spirulina platensis*, *Azotobacter chroococcum*, *Trichoderma harzianum* and their combinations (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>4</sub>, T<sub>12</sub>) were used as spray treatment in the open field and on strawberry after picking. The individual bio agents treatments decreased liner growth and spore production of *B. cinerea* in comparing with mixed bio agents. Also, all the treatments decreased Disease Severity% (DS%). The treatments of T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> caused 13.9, 22.3, 22.3 and 24.1 DS% during 2014; 12.26, 15.5, 17.0 and 21.86 DS% during 2015 in comparing with the mixed bio agents (T<sub>12</sub>) which caused 29.3 and 29.03 DS% during 2014 and 2015, respect., after three sprayings in the field. Also, the treatments of T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> caused 0.00 DS% after 2<sup>nd</sup> spray through the storage at 5°C. The treatment of T<sub>12</sub> caused the highest increase in the total sugar content in strawberry fruits which infected naturally with *B. cinerea* and stored at 5°C for three weeks increased with T<sub>12</sub>. In addition, the treatment of T<sub>12</sub> caused the highest increase in polyphenol oxidase (PPO) activity and peroxidase (PO) which were 0.459 and 0.360 U/mg fresh weight respect., in comparing with T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> which caused 0.278, 0.287, 0.298, 0.313 and 0.139, 0.202, 0.276, 0.302 U/mg fresh weight respect., after storage for 6 weeks. The treatment of T<sub>1</sub> gave the best result in K% in leaves (1.82). T<sub>5</sub> gave the best value of N% in soil (72.2%), T<sub>3</sub> gave the best one in P% in soil (36.0%) and T<sub>4</sub> caused the best amount of K% (16.0) in the soil.

**Keywords:** Bio agents; Strawberry; *Clorella sp*; *Spirulina sp*; *Azotobacter sp*; *Trichoderma sp*

## Introduction

Strawberry (*Fragaria ananassa*) is one of the most widely grown small fruit crops in the world. Fungal diseases of strawberry, mainly caused by *Colletotrichum acutatum* and *Botrytis cinerea*, are responsible for severe economic losses [1,2]. Strawberry fruit have a very short postharvest life, gray mold caused by *B. cinerea* Infection may occur in the lower, remain quiescent until fruits mature, and then develop abundantly, causing fruit decay accompanied by profuse sporulation of the pathogen *B. cinerea* also causes significant losses during shipping and marketing making it one of the most economically important pathogens of strawberry [1].

*Botrytis cinerea* is an airborne and soil borne plant pathogen with anecrotrophic life style attacking over 200 crop hosts worldwide over 200 mainly dicotyledonous plant species, including important protein, oil, fiber and horticultural crops, in temperate and subtropical regions. It can cause soft rotting of all aerial plant parts, and produce prolific grey conidiophores and (macro) conidia typical of disease [2] *Botrytis* fruit rot, also known as gray mold, caused by the fungus *B. cinerea* and is one of the most important diseases of strawberry worldwide. The disease affects fruit in the field resulting in severe pre-harvest losses. It also affects fruit after harvest, since infections that begin in the field continue to develop during storage and transit at refrigeration temperatures. Strawberry flowers are highly susceptible to *B. cinerea*, and may be blighted directly. However, symptoms usually are observed later on green and ripening fruit. Lesions typically develop on the stem end of the fruit and are often associated with infected stamens or dead petals adhering to the fruit or trapped beneath the calyx *B. cinerea* is a common colonizer of strawberry foliage in the nursery, and is also present on dying vegetation around strawberry fields. During the last 50 years, management of *B. cinerea* relied heavily upon the use of synthetic chemicals [3].

Red and brown algae are mainly used as human healthy food sources, due to their high concentration in polysaccharides, natural richness in minerals, polyunsaturated fatty acids and vitamins. Macroalgae (seaweeds) are rich sources of structurally new and biologically active metabolites. In recent years, there have been many reports of macro algae derived compounds that have a broad range of biological activities, such as antibacterial, antiviral, antioxidant, anti-inflammatory, cytotoxic and antimitotic activities Special attention has been reported for antiviral, antibacterial and/or antifungal activities against human pathogens and biostimulant properties of seaweeds are explored for use in agriculture [4-10]. Biochemical resistance through the accumulation of various phenolic compounds and phytoalexins and the activation of peroxidases, polyphenol oxidases and key enzymes in the phenylpropanoid and isoflavonoid pathways may play a crucial role in the biological control of and resistance to pathogenic attack in plants. The defense strategy of plants consists of two stages. The first stage is assumed to involve the rapid accumulation of phenols at the infection site, which function to slow the growth rate of the pathogen and to allow for the activation of "secondary" strategies that will more thoroughly restrict the pathogen. The secondary responses involve the activation of specific defense mechanisms, such as the synthesis of molecules related to pathogen stress [11]. However, it is still not easy and costly in application. It can serve as the best control

**\*Corresponding authors:** Safinaz A. Farfour, Environmental Biotechnology Department, Genetic Engineering and Biotechnology Research Institute (GEBRI), University of Sadat City, Egypt, Tel: +20235720944; E-mail: farfour.safinaz@yahoo.com

Received June 11, 2015; Accepted June 24, 2015; Published June 28, 2015

**Citation:** El-ghanam AA, Farfour SA, Ragab SS (2015) Bio-Suppression of Strawberry Fruit Rot Disease Caused by *Botrytis cinerea*. J Plant Pathol Microbiol S3: 005. doi:10.4172/2157-7471.S3-005

**Copyright:** © 2015 El-ghanam AA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

measure under green-house conditions. The concern of pesticide use with respect to human health and environment has brought increasing interest in alternatives by avoiding negative effects on the environment. Today, there are strict regulations on chemical pesticide use, and there is political pressure to remove the most hazardous chemicals from the market.

*Trichoderma harzianum* isolates were also reported to control strawberry grey mold [12]. Various isolates of *Trichoderma spp.*, that was effective in controlling anthracnose and grey mold in strawberry, under laboratory and greenhouse conditions [13,14]. It has been reported on various occasions that the combination of a number of biological control agents (BCAs) can further improve grey mold control [13,14]. Also, it was reported that co-application of T-39 of *T. harzianum* and AQ10 of *Ampelomyces quisqualis* resulted in improved control of grey mould in cucumber than with each organism alone [15].

Calcium is one of the main components in the plant cell wall and, it causes the cell wall solidity [16]. Calcium addition showed positive effects at fruit density and fruit nutrition value growth and crop yields [17,18]. Nitrogen assisted in amino acids and enzymes of the plant and this element participate at contain of the fruit taste and lignin. Also it effects on plant vegetative growth, leave and stems. Increasing of N causes yield enchasing, fruit size and TSS addition. Of course extra Nitrogen causes to decline the fruit density in ripening proses by function of cell wall dissolution enzymes [19].

Khafagi, [20] found that sugar contents increased in extract of strawberry fruit inoculated with *phytophthora cactorum* as compared with uninoculated fruits However, the least susceptible cultivar (Aliso) showed the lowest amount of reducing and total sugars as compared with the most susceptible one (Baugar). El-neshway [21] reported that sugar contents of strawberry extracts increased as a total and reducing sugars in the least susceptible cultivars. However, reverse trend was noticed with the non- reducing sugars. Generally, the amount of reducing and total sugars varied from one cultivar to another regardless of the infection percentage or level of susceptibility.

The present study is directed for testing some bio agents against strawberry fruit rot disease and its effect on N, P and K contents of strawberry leaves, the activities of polyphenol oxidase and peroxidase and fruits sugar content.

## Materials and Methods

### Source of fungal pathogen

In these experiments; four bio agents were used alone or as mixtures besides the fungicide (switch) and untreated one as controls. Diseased fruits of strawberry showing various types of rots collected from field (fresh planting system) at EI-Tahreer city regions. Fruit samples were rinsed several times in sterilized water and surface sterilized by using 70% ethyl alcohol for two minutes and dried between sterilized filter papers. They were then cut into small pieces and placed in petri- dishes containing potato dextrose agar medium (PDA). The dishes were incubated for 7 days at 25°C. Developed mycelium was transferred and kept on PDA slants.

### Preparation of bio agents

**Trichoderma harzianum:** In order to isolate from healthy plants, it collected from different cultivated areas at Sadat City. Roots and stem bases of the obtained samples were washed tap water, surface sterilized by 70% ethanol and then left to dry on sterilized filter papers. The samples were cut into small pieces. Plated on PDA medium and

incubated at 25°C. Petri dishes were examined daily. Isolates were purified and identified according to Barnett and Hunter [22]. The spores of 7 days old culture was collected and suspended in water and adjusted at  $2 \times 10^7$  spore/ml water using haemocytometer, then applied as foliar spray at the rate of 5 ml/plant.

**Azotobacter chroococcum:** *Azotobacter chroococcum* isolate was grown on Jensen medium for 48 hrs in a rotary shaker at 25°C. The bacterial inoculum was applied as a foliar treatment at the rat of 5 ml bacterial suspension per plant ( $1 \times 10^8$  cfu/ml).

**Algae extract:** *Spirulina platensis* and *Chlorella vulgaris* extract were obtained from International Research Center, Giza, Egypt as digested solution; the extract was suspended by the rate of 50 g/L water. The algal extract was applied as foliar spray at the rate of 5 ml/plant [23].

### Determining liner growth of *Botrytis cinerea*

Effect of the tested antagonists on the mycelia growth of *B. cinerea* was carried out by inoculating PDA culture with 5 mm. discs of the tested pathogens against discs of the *T. harzianum* and against streak of the *Azotobacter* and algae, 6 cm apart. The tested antagonistic bio agents were inoculated at the same time with the pathogen. Four replicates were used for each treatment. Petri-dishes were kept at  $20 \pm 2^\circ\text{C}$  until the mycelia growth covered the entire surface on the control plates, and the percentage of the mycelia growth inhibition was calculated as the following formula:

$$\text{Percentage of mycelia growth inhibition} = \frac{dc-dt}{dc} \times 100$$

Where:

dc=average diameter of fungal colony in the check.

dt=average diameter of fungal colony in the treatment.

### Treatments

Four bio agents *i.e.*, *Chlorella vulgaris*, *Spirulina platensis*, *Azotobacter chroococcum*, *Trichoderma harzianum* and their combinations were applied in 12 treatments (from  $T_1$  to  $T_{12}$ ) in comparing with chemical control ( $T_{13}$ ) and untreated control ( $T_{14}$ ) as shown in Table 1.

**Effect of tested bio agents on *Botrytis cinerea* spore production:** After determining linear growth of the pathogen, spores were collected and transferred to sterilized water (50 ml/treatment). Spores were then counted by using haemocytometer and spore concentration/ml was determined according to the following equation:

$$\text{Spore conc.} = \text{total spores count four squares} \times 2500 \times \text{dilution factor.}$$

**Filed experiment:** Experiment was carried out at El-Tahreer city regions. Healthy strawberry transplants cv. Festival were used for fresh planting system. Treatments were mentioned in Table 1: ( $T_1$ ,  $T_2$ ,  $T_3$ ,  $T_4$ ,  $T_5$ ,  $T_6$ ,  $T_7$ ,  $T_8$ ,  $T_9$ ,  $T_{10}$ ,  $T_{11}$ ,  $T_{12}$ ,  $T_{13}$ ,  $T_{14}$ ) and sprayed in January, 2014 and 2015 three times a week interval with four replicates; each replicate contained 50 plants. The percentage of fruit rot severity caused by *B. cinera* was estimated by the equation adopted by Hanounik.

$$\text{Disease severity (DS)\%} = \frac{\sum(\text{NPC} \times \text{CR})}{\text{NIP} \times \text{MSC}} \times 100$$

Where:

CR= Class rate

NPC=No of plants in each class rate

NIP=No of infected fruits

MSC=Maximum severity class rate

Treatment	Code
<i>Chlorella vulgaris</i>	T <sub>1</sub>
<i>Spirulina platensis</i>	T <sub>2</sub>
<i>Azotobacter chroococcum</i>	T <sub>3</sub>
<i>Trichoderma harizianum</i>	T <sub>4</sub>
<i>Trichoderma harizianum</i> + <i>Chlorella vulgaris</i>	T <sub>5</sub>
<i>Trichoderma harizianum</i> + <i>Spirulina platensis</i>	T <sub>6</sub>
<i>Trichoderma harizianum</i> + <i>Azotobacter chroococcum</i>	T <sub>7</sub>
<i>Azotobacter chroococcum</i> + <i>Chlorella vulgaris</i>	T <sub>8</sub>
<i>Azotobacter chroococcum</i> + <i>Spirulina platensis</i>	T <sub>9</sub>
<i>Azotobacter chroococcum</i> + <i>Chlorella vulgaris</i> + <i>Trichoderma harizianum</i>	T <sub>10</sub>
<i>Azotobacter chroococcum</i> + <i>Spirulina platensis</i> + <i>Trichoderma harizianum</i>	T <sub>11</sub>
<i>Azotobacter chroococcum</i> + <i>Chlorella vulgaris</i> + <i>Spirulina platensis</i> + <i>Trichoderma harizianum</i>	T <sub>12</sub>
Chemical control (switch)	T <sub>13</sub>
Untreated control	T <sub>14</sub>

**Table 1:** The experimental treatments codes.

**Effect of bio agents on the incidence of fruit rot in laboratory:** Healthy mature fruits of cv. Festival were sprayed with different bio agents as follow:

- 1- Sprayed on laboratory.
- 2- Sprayed once only in the field.
- 3- Sprayed two times in the field, one week interval.
- 4- Sprayed three times in the field, one week interval.

All the previous treatments were collected after a week from the last spray and stored six weeks at 5°C.

**Determination of sugar contents:** Quantitative analysis of total, reducing and non-reducing sugars were determined, according to Tomas and Ducher [23] by using picric acid method.

The determinations were calculated as milligrams glucose per gram fresh weight.

**Determination of nitrogen, phosphorous and potassium:** Contents in strawberry leaves, soil, *Azotobacter chroococcum* and algae were determined in the central laboratory of soil fertility, Agriculture ministry, Giza, Egypt using the methods described by Awad [24] (Table 2).

**Determination of enzymes activity:** The enzymes activity of polyphenol oxidase and peroxidase were determined. Samples of 100 gm of strawberry of each treatment were blended in 200 ml distilled water for 2 min. The mixture was squeezed through several layers of cheese cloth and centrifuged at 3000 rpm for 20 minutes. Supernatants were kept at 5°C until assaying. The measurement of enzyme activity was carried out in the prepared filtrates by measuring according to the method of Aneja [25] using spectrophotometric procedure; as described by Matta and Dimond [26].

## Results and Discussion

Data in Table 3 show that reduction in mycelia growth was noticed with all bio agents and the highest inhibition of *Botrytis cinerea* mycelia growth was exhibited in case of T<sub>3</sub> and T<sub>1</sub> followed by T<sub>4</sub> and T<sub>2</sub> with averages of 61.11, 58.30, 55.50 and 50.00 mm., respectively. Moreover, inhibition was moderately with the other mixed bio agents; T<sub>5</sub>, T<sub>9</sub>, T<sub>6</sub>, T<sub>11</sub>, T<sub>7</sub> and T<sub>8</sub>. Sporulation of *B. cinerea* showed significant reduction

with all the tested bio agents (*Chlorella vulgaris*, *Spirulina platensis*, *Azotobacter chroococcum* and *Trichoderma harizianum*) alone or as mixture. Regardless of the sequence of application rates differed according to the potency of the tested bio agents and application sequence.

The highest reduction in sporulation of *B. cinerea* were obtained with T<sub>1</sub> treatment (2.2%), compared with those of the other tested bio agents, followed by T<sub>3</sub> and T<sub>4</sub> (3.3%).

In this respect, cyanobacteria (blue-green algae) and eukaryotic algae produce biologically active compounds that have antifungal activity antibiotic and toxic activity (Bonjouklian and Kiviranta [27,28]). Also Hussien [29] evaluated the effect of culture filtrates of nine algal strains at concentrations of 10, 20, 30, and 40% on mycelium growth and spore production of *Cercospora beticola* causing leaf spot disease in sugar beet. Generally, they found that all the algal culture filtrates reduced the fungal mycelium growth, but the best results were obtained by *Spirulina platensis*, *Oscillatoria* sp. and *Nostoc muscorum*; the highest fungal mycelium growth inhibition percentage was achieved by the concentrations of 30% (100, 100, and 82%, respectively.) and at 40% (100, 100, and 100%, respectively). Fungal spore production (number of spores) was completely inhibited by the previous three algal culture filtrates particularly at the concentration of 40%. They conclude that the antifungal activity of the algal culture filtrates has been attributed to the presence of bioactive compounds, that is, total phenolic compounds, total saponins, and alkaloids in the algal culture filtrates. Also, several workers reported that the extracts of *Nostoc muscorum* significantly inhibited the growth of *Candida albicans* and *Sclerotinia sclerotiorum* and [30] aBis (2, 3-dibromo-4, 5-dihydroxybenzyl) ether (BDDE) is a bromophenol isolated from marine algae which possesses cytotoxic and antibacterial activity. BDDE inhibits the growth of *B. cinerea* cultured on a solid medium

Bio agents	N%	P%	K%
<i>Chlorella vulgaris</i>	1.62	0.39	3.96
<i>Spirulina platensis</i>	1.17	0.19	2.09
<i>Azotobacter chroococcum</i>	3.2	0.28	2.25

**Table 2:** Nitrogen, phosphorous and potassium content of some bio agents contents.

Treatment	Linear growth (mm)		Spores production x10 <sup>3</sup>
	Reduction	Mycelia growth (mm)	
T <sub>1</sub>	58.3	37.5	2.2
T <sub>2</sub>	50.1	45.0	4.5
T <sub>3</sub>	61.1	35.0	3.3
T <sub>4</sub>	55.5	40.0	3.3
T <sub>5</sub>	25.0	67.5	5.5
T <sub>6</sub>	23.3	72.5	5.5
T <sub>7</sub>	22.2	70.0	6.1
T <sub>8</sub>	20.5	71.5	6.2
T <sub>9</sub>	25.0	67.5	5.5
T <sub>10</sub>	30.5	62.5	13.6
T <sub>11</sub>	23.3	72.5	14.0
T <sub>12</sub>	19.4	74.0	14.9
Switch (2200 ppm)	100.0	0.0	0.0
Untreated Control t	0.0	90.0	100
L.S.D at 5%	0.23		0.60

**Table 3:** The effect of different bio agents on the liner growth and spores production of *B. cinerea*.

of potato dextrose agar (PDA) as well as on the potato dextrose broth (PDB) medium. Further studies have revealed that BDDE decreases the germination rate and inhibits the mycelia growth of *B. cinerea*. The inhibition mechanisms are related to the disruption of the cell membrane integrity in *B. cinerea* spores and newly formed germ tubes. *Spirulina platensis* – green algae (*Chlorophyta*); *Fucus vesiculosus*, inhibited the mycelia growth *Botrytis cinerea* Bc 2107 with at 50% 0.5% and 90% at 2.02% algae the presence of bioactive molecules, as phenolic compounds (phlorotannins, terpenes, alkaloids), polysaccharides or fatty acids, many of these structures being identified as antimicrobials [31]. *Azotobacter* is able to inhibit the growth of phytopathogenic fungi species such as *Alternaria*, *Venturia*, *Sclerotinia*, *Rhizoctonia*, and *Pythium* [32]. The isolates of *Trichoderma* which chose possessed variety of potential modes of action *in vitro* [33]. Ap-PCR indicated that isolate T-105 survived better than the other isolates in the mix [34].

Data presented in Table 4 indicate that percentage of disease severity of fruit rot decreased by spraying strawberry fruits with all the different bio agents in field and laboratory after cold storage at 5°C for six weeks. Results showed that the lowest disease severity were (0.0) with (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, and T<sub>4</sub>) with two, three and four spray treatments compared with untreated control (100%), where at the first treatment, disease severity were 0.3, 0.6, 0.3 and 1.6 with T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, and T<sub>4</sub>, respectively. Therefore, the highest reduction rate (40.3%) was obtained on spraying strawberry fruits, growing on field with mixed bio agents (T<sub>12</sub>) and stored at 5°C for six weeks.

These compounds could be environmentally friend means of plant disease control and could be utilized in organic farming and for vegetable cropping systems where application of synthetic fungicides or chemicals needs to be avoided [24]. Antifungal compounds have been detected in different algal species. For example, extracts of the brown alga *Cystoseira tamariscifolia* showed *In Vitro* fungal activity against the plant pathogens *B. cinerea*, *F oxysporum* and *Verticillium album-atrum* [35].

Application of biological control using antagonistic microorganisms has proved to be successful for controlling various plant diseases [36]. Recently algae are one of the chief biological agents that have been studied for the control of plant pathogenic fungi [37].

Data in Table 5 show the effect of sprayed strawberry by some bio agents on the disease severity% (DS%) of fruit rot under field conductions. It is clear that at the first season (2014) T<sub>1</sub>, T<sub>7</sub>, T<sub>9</sub>, T<sub>2</sub> and T<sub>3</sub> had the best effect in reducing DS% on the other hand the same treatment was cleared in the second season (2015) with light differences in arrangement where T<sub>1</sub> had the first grade followed by T<sub>10</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>7</sub>, respect., and generally the fungicide Switch decreased DS% more than any other treatment. Generally disease severity decreased when sprayed strawberry fruit under field conditions by *Clorella vulgaris* (T<sub>1</sub>) compared with (T<sub>2</sub>, T<sub>5</sub>, T<sub>6</sub>, T<sub>8</sub>, and T<sub>9</sub>) during (2014&2015). The results indicated that the application of bio agents (T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>7</sub>) decreased disease severity during season 2014 and 2015 control.

This result is in agreement with Edra et al., [38]. who found that the brown-alga *Lessonia trabeculata* inhibited bacterial growth and reduced both the number and size of the necrotic lesions in tomato leaves following infection with *Botrytis cinerea*. Marine algae represent a great source of a vast variety of complex natural products and could be a promising source of a novel bioactive compound that can help plant survival by offering protection against stress imposed by pathogens. Aqueous and ethanolic extracts from the red-alga *Gracillaria chilensis* prevent the growth of *Phytophthora cinnamomi*. Similarly, aqueous

and ethanolic extracts from the brown-alga *Durvillaea antarctica* were able to diminish the damage caused by tobacco mosaic virus (TMV) in tobacco leaves. These results suggest that macro-algae contain compounds with different chemical properties which could be considered for controlling specific plant pathogens. On the other hand, *Trichoderma* isolates, as well as T-39 and TRICHODEX were used effectively, for the control of *Colletotrichum* and *Botrytis* on strawberry petioles and leaves [14]. Isolates T-115, T-161, T-166, T-39 and TRICHODEX were effective in reducing *Botrytis* leaf grey mould incidence whereas T-118, T-165, T-166, T-39 and TRICHODEX were effective in reducing anthracnose in petioles [14].

Data in Table 6 summarize the amounts of reducing, non-reducing and total sugars in naturally infected fruits with *Botrytis cinerea* and treated by different bio agents, after 1, 2, 3 weeks of storage at 25°C. Reducing sugars increased with increasing incubation periods from one to three weeks. The highest value of reducing sugar treated with mixed

Treatment	Disease severity%			
	1 <sup>st</sup> spray*	2 <sup>nd</sup> spray**	3 <sup>rd</sup> spray**	4 <sup>th</sup> spray**
T <sub>1</sub>	0.3	0.0	0.0	0.0
T <sub>2</sub>	0.6	0.0	0.0	0.0
T <sub>3</sub>	0.3	0.0	0.0	0.0
T <sub>4</sub>	1.6	0.0	0.0	0.0
T <sub>5</sub>	24.0	20.0	9.0	3.0
T <sub>6</sub>	22.3	14.3	8.6	2.6
T <sub>7</sub>	27.3	13.0	7.6	3.3
T <sub>8</sub>	19.9	12.6	9.0	6.6
T <sub>9</sub>	19.3	13.6	6.7	2.6
T <sub>10</sub>	36.6	20.8	13.0	8.2
T <sub>11</sub>	33.6	19.7	13.5	7.0
T <sub>12</sub>	40.3	25.0	20	9.0
Switch (T <sub>13</sub> )	0.0	0.0	0.0	0.0
Untreated control (T <sub>14</sub> )	100	100	100	100
L.S.D at 5%	0.48	0.47	0.47	0.46

\* In the Lab.  
\*\*In the field.

**Table 4:** Effect of some bio agents on disease severity percentage of strawberry fruits after storage for six weeks at 5°C.

Treatment	Disease severity							
	2014				2015			
	1 <sup>st</sup> spray	2 <sup>nd</sup> spray	3 <sup>rd</sup> spray	Mean	1 <sup>st</sup> spray	2 <sup>nd</sup> spray	3 <sup>rd</sup> spray	Mean
T <sub>1</sub>	19.6	13.3	9.0	13.9	17.5	12.3	7.0	12.2
T <sub>2</sub>	21.0	18.0	11.6	22.3	20.0	17.0	9.6	15.5
T <sub>3</sub>	32.0	19.0	16.0	22.3	19.0	18.0	14.0	17.0
T <sub>4</sub>	35.3	20.3	17.0	24.1	30.3	20.3	15.0	21.8
T <sub>5</sub>	38.3	23.6	20.0	27.3	38.3	22.6	19.0	26.6
T <sub>6</sub>	28.3	32.6	16.3	25.7	26.3	30.6	15.3	24.6
T <sub>7</sub>	23.8	23.0	15.3	20.0	22.6	20.0	14.3	18.9
T <sub>8</sub>	29.0	27.0	20.3	25.4	28.0	26.6	19.3	24.0
T <sub>9</sub>	23.3	22.0	17.3	20.8	22.6	21.0	16.3	19.9
T <sub>10</sub>	35.3	23.6	20.6	26.4	18.3	15.6	12.6	15.4
T <sub>11</sub>	40.3	29.0	20.6	29.9	39.6	28.0	18.3	28.6
T <sub>12</sub>	41.6	27.0	19.3	29.3	40.0	26.6	20.5	29.0
Switch (T <sub>13</sub> )	13.6	8.6	5.3	9.16	13.6	7.6	4.3	8.5
Untreated control (T <sub>14</sub> )	55.7	53.3	54.0	54.3	56.0	53.3	50.0	53.1
L.S.D at 5%	0.58	0.92	1.29		0.60	0.95	1.30	

**Table 5:** Effect of some bio agents on disease severity percentage (DS%) of Strawberry fruits in field experiment during two successive seasons 2014 and 2015.

Treatment	Sugar content (mg/g fresh weight of fruit)								
	1 week			2 weeks			3 weeks		
	R	NR	T	R	NR	T	R	NR	T
T1	73.1	11.6	84.7	83.5	11.7	95.2	93.2	11.7	210.2
T2	88.3	11.7	100	93.5	21.4	114.9	93.6	26.6	120.2
T3	86.3	12.1	98.4	92.5	20.4	112.9	92.6	25.5	118.1
T4	86.3	12.1	98.4	92.3	27.3	119.6	104.2	22.5	126.4
T5	93.2	22.2	115.4	98.4	27.3	125.7	100.1	27.3	127.4
T6	93.1	21.3	114.4	98.1	31.2	129.3	104	27.2	131.2
T7	94.4	22.5	116.9	99.3	31.2	130.5	108	32.4	140.4
T8	98.2	27.0	125.2	102.1	46.5	148.6	109	46.4	155.4
T9	99.3	27.6	126.9	104.0	51.5	155.5	118.5	47.2	165.7
T10	102.1	31.2	133.3	116.4	67.4	183.4	128.1	67.4	195.5
T11	109.8	50.4	160.2	114.9	77.3	192.2	134.8	82.6	217.4
T12	118.7	62.3	181	120.2	83.2	203.4	138.7	95.6	234.3
Switch(T <sub>13</sub> )	73.1	11.6	84.7	78.7	21.2	99.9	93.1	11.7	104.8
Untreated control(T <sub>14</sub> )	128.1	64.3	192.4	130.3	85.2	215.5	140.3	97.5	237.8
L.S.D at 5%	0.49	0.54	0.60	0.49	0.59	0.44	0.57	0.4	0.59

R: Reducing sugars; NR: Non-reducing sugars T: Total sugars.

**Table 6:** Sugar content mg/g fresh weight of strawberry fruits and naturally infected with *B. cinerae* treated with different bioagents after storage one, two and three weeks.

Treatment	Polyphenoloxidase (U/mg fresh weigh)			Peroxidase(U/mg fresh weigh)		
	1 <sup>st</sup> spray	2 <sup>nd</sup> spray	3 <sup>rd</sup> spray	1 <sup>st</sup> spray	2 <sup>nd</sup> spray	3 <sup>rd</sup> spray
T1	0.278	0.309	0.320	0.139	0.231	0.267
T2	0.287	0.310	0.325	0.202	0.305	0.325
T3	0.298	0.314	0.330	0.276	0.312	0.368
T4	0.313	0.330	0.350	0.302	0.313	0.370
T5	0.325	0.452	0.520	0.304	0.345	0.375
T6	0.367	0.509	0.624	0.314	0.391	0.400
T7	0.375	0.596	0.600	0.326	0.401	0.435
T8	0.329	0.620	0.630	0.335	0.423	0.450
T9	0.363	0.724	0.730	0.345	0.444	0.455
T10	0.379	0.858	0.860	0.350	0.445	0.456
T11	0.380	0.870	0.880	0.358	0.488	0.470
T12	0.459	0.897	0.910	0.360	0.490	0.475
Switch(T <sub>13</sub> )	0.220	0.310	0.310	0.150	0.228	0.270
Untreated control(T <sub>14</sub> )	0.410	0.911	0.920	0.360	0.495	0.500
L.S.D at 5%	0.012	0.003	0.82	0.45	0.04	0.29

**Table 7:** Effect of different bio agents on fruits Polyphenol oxidase and Peroxidase activities in treated strawberry fruits after storage six weeks at 5°C.

bio agents treatment (T<sub>12</sub>) after storage periods 1, 2 and 3 weeks with averages 118.7, 123.0 and 138.7 mg/g fresh weight of fruit, respectively. Generally, reducing, non-reducing and total sugars increased sharply in case of treatments T<sub>10</sub>, T<sub>11</sub> and T<sub>12</sub> compared with the other treatments but the treatment (T<sub>14</sub>) had the highest values.

Early studies have reported that germination in most *B. cinerea* isolates is dependent on the presence of nutrients, mainly sugars, and that this process can be stimulated by exudates from mature fruits and [39-41]. Fouri and Holzm showed that in *B. cinerea* fructose at micro molar concentration was a more potent germination inducer than glucose. Glucose, fructose and sucrose were the better inducers of germination.

Data presented in Table 7 indicate that the enzyme activities increased with treatments of mixed bio agents. The least activities of Polyphenol oxidase and Peroxidase cleared with T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> after spraying three times with different bio agents and stored for six weeks under 5°C. At the same time treatment with the fungicide (switch) led to decrease in the enzymes activity, in contrast control treatment had the highest values. Generally, mixing the bio agents resulted excess in the activity for both enzymes compared to the treatment separately.

Disease reduction was accompanied with a gradual increase in peroxidase activity during experiment period (Table 7).

The compared works showed that applying chemical inducers resulted in a gradual increase in peroxidase activity in *faba bean* plants pre-sprayed and inoculated with *Botrytis fabae* or *Botrytis cinerea* during examination periods. Increases in peroxidase activities were higher compared with untreated inoculated plants (control). Maximum increase in peroxidase activity was detected after 24 h. in *B. fabae* and 48 h. in *B. cinerea*, resulted in remarkable increase in disease reduction resulted from the inoculation of both pathogens [42]. Also Vance et al. [43] and Fry, [44]. Fry stated that peroxidase is known to be involved in the oxidation of polymerization of hydroxycinnamyl alcohols to yield lignin and crosslinking isodityrosine bridges in cell wall. Also, Ride [45] and Tarrad [46] reported that increase in peroxidase activity enhance lignification in response to chocolate spot infection which may restrict the fungal penetration. These findings indicate a positive relationship between resistance and peroxidase activity. Peroxidase also produces free radicals and hydrogen peroxide which are toxic to many microorganisms [47] Another supportive suggestion was brought by Nawar and Kuti [48] who stated that an increase in peroxidase activity is considered as a preliminary indicator for. Laccase (PPO), a fungal

polyphenol oxidase from *Botrytis cinerea*, has been widely studied in relation to its effect on the composition and quality of must and wine [49-51]. In particular, PPO has been shown to be the main enzyme involved in oxidation of red and white wines from rotten grapes. This involves oxidation at all stages the winemaking process, ageing, and while stored in bottles. This is largely because PPO is ubiquitous enzyme in the plant world, requiring a polyphenol substrate, and is well adapted to the physicochemical conditions of wine (pH, ethanol, SO<sub>2</sub>) [52-55]. Peroxidase also functions in the metabolism of reactive oxygen species (ROS) and reactive nitrogen species (RNS), thus activating the hypersensitive response (HR), a form of programmed cell death at the infection site that is associated with limiting pathogen development [56]. Peroxidase is thought to be involved in the polymerization of phenolic monomers to generate the aromatic matrix of suberin. In addition, polyphenol oxidase and peroxidase are activated by pathogens, resulting in the oxidation of phenolics to form quinones, which are effective inhibitors of phytopathogens. In addition, a large number of toxic phytoalexins can be derived from phenolic compounds. Peroxidases, polyphenol oxidases and key enzymes in the phenylpropanoid and isoflavonoid pathways may play a crucial role in the biological control of and resistance to pathogenic attack in plants. The defense strategy of plants consists of two stages. The first stage is assumed to involve the rapid accumulation of phenols at the infection site, which function to slow the growth rate of the pathogen and to allow for the activation of “secondary” strategies that will more thoroughly restrict the pathogen. The secondary responses involve the activation of specific defense mechanisms, such as the synthesis of molecules related to pathogen stress [12].

Nitrogen, phosphorous and potassium contents were determined in soil, strawberry leaves and algae extracts (*Chlorella vulgaris* and *Spirulina platensis*) and *Azotobacter chroococcum* after spraying strawberry plants with different bio agents. Data presented in Table 8 cleared that, N content in strawberry leaves increased in the treatments T<sub>3</sub>, T<sub>4</sub>, T<sub>12</sub> and T<sub>11</sub> compared to the other treatments except T<sub>13</sub> (Fungicide switch) which had the highest value (1.41%). From the same table, P content showed an increase with the treatments especially in case of T<sub>12</sub>, T<sub>10</sub>, T<sub>1</sub>, T<sub>5</sub> and T<sub>6</sub> with the average of 0.35, 0.35, 0.34, 0.32 and 0.32% respectively. In this respect K content increased with the treatments T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, T<sub>7</sub> and T<sub>8</sub> with averages of 1.82, 1.79, 1.77, 1.69 and 1.62%, respectively. Generally, leaves, algae, *Azotobacter* and Soil

contented of calcium is the highest than nitrogen and phosphorous. Data in Table 8 indicate that soil content of nitrogen, phosphorous and calcium is the highest compared with leaves, algae and *Azotobacter* contents of nitrogen, phosphorous and potassium caused by the highest fertilization the Soil. When compared leaves, algae, *Azotobacter* and soil contents of potassium noted that leaves contents of potassium is agree with algae and *Azotobacter* contents.

Therefor algae and *Azotobacter* contents of potassium decreased strawberry fruit rot, when nitrogen is deficit in plants it inhibits the synthesis of 6-phosphogluconate and glucose-6-phosphate dehydrogenase and enhances the synthesis and builds up of phenolic compounds which gives rise browning of curds. In the other hand Eman and Abd-Allahwhen, found that using algal extract at concentrations above 50% caused a progressive increase in N, P and K% in the leaves which was observed as a result of increasing concentration of algal extract till 50% while with concentrations above 50% the increase was slightly appeared.

## References

- Mertely JC, Peres NA (2009) Botrytis fruit rot on gray mold of strawberry University of Florida. IFAS Extension p. 230.
- Williamson BB, Tudzynski P, Tudzynski JAL, Vankan (2007) Botrytis cinerea the cause of grey mold disease. Molecular Plant Pathology 8: 561-580.
- Rosslensbroich HD, Stuebler (2000) Botrytis cinerea - history of chemical control and novel fungicides for its management. Crop Protection 19: 557-61.
- Antimicrobial Demirel ZF, Yilmaz-Koz GU, Karabay-Yavasoglu A, Ozdemir Sukatar (2009) antioxidant activity of brown algae from the Aegean Sea. J.Serb.ChemSoc 74: 619-628.
- Lima-Filho JA, Carvalho S, Freitas V, Melo (2002) Antibacterial activity of extracts of six macroalgae from the northeastern Brazilian coast. Braz.J. Microbiol 33: 311-313
- El-Baz FKAM, AboulEnein GS, El-Baroty AM, Youssef H, Abd El-Baky H (2002) Anticarcinogenic activity of algal extracts. J. Med. Sci 2: 243-251.
- Kandhasamy MKD, Arunachalam (2008) Evaluation of in vitro antibacterial property of seaweeds of southeast coast of India. African Journal of Biotechnology 7: 1958-1961.
- MaasJ L (1998) P. 28 in Compendium of strawberry diseases. (2ndedn) APS Press St.
- Li AK, Cheng C, WongF, KingWai, Feng C, et al. (2007) Evaluation of antioxidant capacity and total phenolic content of different fractions of selected microalgae. Food Chem 102: 771-776.
- Rajasulochana P, Dhamotharan R, Krishnamoorthy P, Murugesan S (2009) Antibacterial activity of the extracts of marine red and brown algae. Journal of American Science 5: 20-25.
- Vallinayagam R, Arumugam RR, Kannan, Thirumaran G, Anantharaman P (2009) Antibacterial activity of some selected seaweeds from Pudumadam Coastal Regions. Global Journal of Pharmacology 3: 50-52.
- Mohamed C, Arbia A, Azza R (2007) Phenolic Compounds and their Role in Biocontrol and Resistance of Chickpea to Fungal Pathogenic Attacks. Tunis.J. Plant Protect. 2: 7-21.
- Trosno A, Dennis C (1977) The use of Trichoderma species to control strawberry fruit rots. Netherlands Journal of Plant pathology 83: 449-455.
- Freeman S, Barbul O, Rav David D, Nitzani Y, Zveibil A, et al. (2001) Trichoderma spp. for biocontrol of Colletotrichum acutatum and Botrytis cinerea in strawberry. Biocontrol of Fungal and Bacterial Plant Pathogens.IOBC/WPRS Bulletin 24: 147-150.
- Elad Y, Barbul O, Nitzani Y, Rav David D, Zveibil A, et al. (2001) Inter- and intra-species variation in biocontrol activity. In Proceedings of the 5<sup>th</sup> Congress of the European Foundation of Plant Pathology 474-478.
- EladY, Kirshner N Yehuda, Szejnberg A (1998) Management of powdery mildew and gray mold of cucumberby Trichoderma Harzianum T39 and Ampelomyces quisqualis. AQ10. Biocontrol 43: 241-251.
- Khayyat M, Tehranifar A, Akbarian A, Shayesteh Nia SH, Khabatri S (2009)

Treatment	In Leaves of strawberry			In soil		
	N%	P%	K%	N%	P%	K%
T <sub>1</sub>	1.21	0.34	1.82	27.7	15.0	112.0
T <sub>2</sub>	1.31	0.32	1.79	37.8	20.0	120.0
T <sub>3</sub>	1.41	0.26	1.77	36.1	36.0	120.0
T <sub>4</sub>	1.33	0.29	1.54	34.4	10.0	160.0
T <sub>5</sub>	1.32	0.32	1.51	72.2	8.0	128.0
T <sub>6</sub>	1.24	0.32	1.44	47.0	8.0	72.0
T <sub>7</sub>	1.17	0.30	1.69	42.0	10.0	88.0
T <sub>8</sub>	1.23	0.31	1.62	61.3	9.0	96.0
T <sub>9</sub>	1.24	0.28	1.45	56.3	8.0	80.0
T <sub>10</sub>	1.30	0.35	1.46	52.1	8.0	80.0
T <sub>11</sub>	1.32	0.29	1.47	60.5	9.0	88.0
T <sub>12</sub>	1.33	0.35	1.43	50.0	10.0	80.0
Switch(T <sub>13</sub> )	1.41	0.27	1.62	27.7	15.0	112.0
Untreated control(T <sub>14</sub> )	1.28	0.38	1.54	37.8	20.0	120.0
L.S.D at 5%	0.49	0.01	0.54	0.63	0.57	0.54

**Table 8:** Effect of some bio-agents on NPK% contents of Strawberry after harvesting.

- Effects of calcium form on electrolyte leakage, total nitrogen, and yield and biomass production by strawberry plant under NaCl salinity. Journal Central European Agriculture 10: 297-302.
18. Pilar Hernandez M, Maria J, Rafael G (2006) Effect of calcium dips and Chitosan coating on postharvest life of strawberry (*Fragaria ananasa*). Postharvest Biology and Technology 39: 247-253.
19. Levent Tuna A, Cengiz K, Muhammad A, Hakan A, Ibrahim Y et al. (2007) The effect of calcium sulphate on growth, membrane stability and nutrient uptake of tomato plants grown under salt stress. Environmental and Experimental Botany 59: 173-178.
20. Tabatabaei SY, Yusefi M, Hajiloo Y (2008) Effects at shading and  $\text{NO}_3\text{NH}_4$  ratio on the yield, quality and metabolism in strawberry. Scintia Horticulture 116: 264-272.
21. Khafagi YS (1989) Studies on powdery mildew disease on strawberry in Egypt. Ph.D. Thesis, Fac. Agric., Suez Canal Univ 106p
22. Barnett HI, Hunter BB (1972) Illustrated genera of imperfect fungi. (3rd edn) Burgess Publ. Co., Minneapolis, Minnesota, USA.
23. Tomas W, Ducher RA (1924) The colorimetric determination of treatise on fungi as experimental Systems for basic and applied research. XI. Agricultural Applications. Springer, Heidelberg, Germany pp. 93-109.
24. Awad NE (2000) Biologically active steroid from the green alga *Ulva lactuca*. Phytother Res 14: 641-643. Veg Sci 32: 62-64
25. Aneja KK (2001) Experiments in microbiology plant pathology tissue culture and mushroom production technology. Third Edition p. 468.
26. Matta J, Dimond AE (1963) Symptoms of Fusarium wilt in relation to quantity of fungus and enzyme cavity in tomato stems. Phytopathology 53: 574-578.
27. Bonjouklian, Smitka RTA, Doolin LE (1991) Tjipanazoles, new antifungal agents from the blue-green alga *Lyngbyella tjeppanensis*. tetrahedron 47(37): 7739-7750.
28. Kiviranta J, Abdel-Hameed A, Sivonen K, Niemela SI, Arlberg G (1993) Toxicity of cyanobacteria to mosquito larvae-screening of active compounds. Environmental Toxicology and Water Quality 8: 63-71.
29. Hussien MY, Abd-El-Aal AAM, Mostafa SS (2009) Bioactivity of algal extracellular byproducts on cercospora leaf spot disease, growth performance and quality of sugar beet. In Proceeding of the 4th Conference on Recent Technologies in Agriculture, Cairo, Giza.
30. Borowitzka M A (1995) Microalgae as sources of pharmaceuticals and other biologically active compounds. Journal of Applied Phycology 7: 3-15.
31. Andree C, Axine O, Iacomi B (2010) Antifungal activity of macroalgae extracts Scientific Papers, UASVM Bucharest, Series A, Vol. LIII, ISSN 1222-5339.
32. Lila NB (2011) Response of cauliflower to nitrogen fixing biofertilizer and graded levels of nitrogen. The Journal of Agriculture and Environment I: 12, Technical Paper 41.
33. Freeman SO, Barbul, Rav David D, Nitzani Y, Zveibil Y, et al. (2001) *Trichoderma* spp. for biocontrol of *Colletotrichum acutatum* and *Botrytis cinerea* in strawberry. Biocontrol of Fungal and Bacterial Plant Pathogens. IOBC/WPRS Bulletin 24: 147-150
34. Elad Y (2000) Biological control of foliar pathogens by means of *Trichoderma harzianum* and potential modes of action. Crop Protection 19: 709-714
35. Abourriche, Harrouf AM, Berrada M, Bennamara A, Chaib N, Francisco C (1999) Fitoterapia Antimicrobial activities and cytotoxicity of the brown alga *Cystoseira tamariscifolia* 70: 611-614.
36. Sivan A (1987) Biological control of Fusarium crown rot of tomato by *Trichoderma harzianum* under field conditions. Plant Disease 71: 587-592.
37. Abo-Shady, Bahia AM, Alghaffar A, Rahhal MMH, Abd-El-Monen HA (2007) Biological control of faba bean pathogenic fungi by three cyanobacterial filtrates. Pok J Bio Sci 10: 3029-3038.
38. Edra J, Dorta F, Medina C, Ramirez A, Ramirez I, Peña-Cortés H (2011) Anti-Phytopathogenic Activities of Macro-Algae Extracts Mar Drugs 9: 739-756.
39. Brodie IDS, Blakeman JP (1976) Competition for exogenous substrates in vitro by leaf surface micro-organisms and germination of conidia of *Botrytis cinerea*. Physiol Pl Pathol 9: 227-239.
40. Fouri JF, Holzm G (1997) Effects of fruit and pollen exudates on growth of *Botrytis cinerea* and infection of plum and nectarines fruit. Pl. Dis. 82: 165-170.
41. Doehlemann G, Molitor F, Hahn M (2005) Molecular and functional characterization of a fructose specific transporter from the gray mold fungus *Botrytis cinerea*. Fungal Genet Biol 42: 601-610.
42. Maggie, Hassan EM, El Abbasi SSAIH, Mikhail MS (2007) Changes in Peroxidase Activity Due to Resistance Induced Against Faba Bean Chocolate Spot Disease Egypt. J. Phytopathol 35: 35-48.
43. Vance CP, Kirk TK, Sherwood RT (1980) Lignification as a mechanism of disease resistance. Annu Rev. Phytopathol 18: 259-288
44. Fry SC (1982) Isodityrosine a new amino acid from plant cell wall glycoprotein. Biochem. J, 204: 449-455
45. Ride JP (1983) Cell walls and other structural barriers in defence. In: Biochemical Plant Pathology. Calloz JA (ed.), John Wiley and Sons, New York, USA
46. Tarred AM, El-Hyatem YY, Omar SA (1993) Weyerone derivatives and activities of peroxidase and polyphenol oxidase in faba bean leaves as induced by chocolate spot disease. Plant Sci 89: 161-165.
47. Pena MJ, Kuc JA (1992) Peroxidase-generated hydrogen peroxidase as a source of antifungal activity In Vitro and on tobacco leaf disks. Phytopathology 82: 696-699.
48. Nawar HF, Kuti JD (2003) Weyerone acid phytoalexin synthesis and peroxidase activity as markers for resistance of broad beans to chocolate spot disease. J. Phytopathol 151: 564.
49. Ribereau-Gayon PD, Donneche BA, Lonvaud-Funel (1998) Handbook of enology. I: The microbiology of wine and vinification 410-419.
50. Minussi RC, Rossi M, Bologna L, Rotilio D, Pastore GM, et al. (2007) Phenols removal in musts: Strategy for wine stabilization by laccase. Mol Catal J B Enzymatic 45: 102-107.
51. Li H, Guo A, Wang A (2008) Mechanism of oxidative browning of wine. Food Chem 108: 1-13.
52. Almagro LL, Gomez RA, Belchi-Navarro S, Bru R, Ros BA, et al. (2009) Plants defend themselves against pathogen attack by activating a multicomponent defense response. J Exp Bot. 60: 377-390.
53. Alfred MM (2006) Polyphenol oxidases in plants and fungi: Going places? A review. Phytochemistry 67: 2318-2331.
54. Orcutt DM, Nilsen, ET (2000) Physiology of Plants Under Stress. John Wiley & Sons, INC, New York. p. 342-344.
55. Gupta AK, Samnotra RK (2004) Effect of biofertilizers and nitrogen on growth, quality and yield of cabbage cv. Golden Acre. Environment and Ecology. 22 551-553.
56. Eman A, Abd El Moniem Abd-Allah ASE (2008) Effect of Green Alga Cells Extract as Foliar Spray on Vegetative Growth, Yield and Berries Quality of Superior Grapevines American-Eurasian J Agric & Environ Sci 4: 427-433.

This article was originally published in a special issue, **Pathological Findings in Plants** handled by Editor(s). Dr. Chioma Okeoma, Department of Microbiology University of Iowa, USA