

## Editor's Note: Volume 7 Issue 9

Purushottam S Narute\*

Clinical Center, National Institutes of Health, USA

Since the first case of mysterious disease reported in 1981, HIV-1 pandemic caused death of more than 25 million people around the world. Due to intrinsic mutagenic ability and high variability among HIV genome it is very difficult to develop globally effective therapeutics as well as vaccines. The advancement of highly active anti-retroviral therapy (HAART) and several experimental vaccine approaches are less effective in controlling new infections as well as quality of life in developing countries. In essence prevention is better strategy to combat with this HIV/AIDS global epidemic. The current issue focuses on several different prevention as well as therapeutic approaches in developing countries. Navaneethapandian et al. compared the growth patterns of HIV-infected and HIV-exposed, uninfected infants from India and reported that mortality is high in the absence of early antiretroviral treatment in HIV-1 infected infants [1]. With similar lines Birlie et al. focused on prevention of mother-to-child transmission of HIV and performed retrospective study in Southwest Ethiopia. This study concluded that pregnant HIV positive mothers should be enrolled in PMTCT programs at earlier stage and exclusive breast feeding should be encouraged so as to decrease MTCT [2].

To develop new therapeutics, Tincho et al. used *in silico* approach to identify anti-HIV peptide-based drugs [3]. Indeed prevention is better than cure, Girma T, et al. studied factors affecting knowledge, attitude and voluntary counseling and testing among students from Ethiopia [4]. In similar lines, Mapara describes use of color pictures to raise awareness about AIDS [5]. Similarly, Applewhite S discusses use of pre-exposure prophylaxis and antiretroviral treatment to reduce infections [6]. Swartz et al. performed LGBTI case study to understand psychosocial well-being of LGBTI individuals and communities in South Africa [7]. Raheel studied the stigmatizing attitudes and factors determining stigma toward PLWA prevailing in the Saudi Arabia [8]. Moreover, Avanzi et al. studied significance of genetic variation in interleukin 28B (IL28B) and inosine triphosphatase (ITPA) genes in cohort of HIV/HCV co-infected patients in Southern Brazil [9]. Saleem et al. studied antibiotic resistance in Gonorrhea for prevention of STD [10]. In addition to this Ramdial et al. studied lymph node biopsy in patients with nodal donovanosis [11].

Overall this issue provides recent research on HIV prevention and development of new therapeutic which will be very interesting for readers.

### References

1. Navaneethapandian PGD, Rajendran M, Ramesh K, Elangovan T, Subramanyan S, et al. (2016) Growth of infants born to HIV-infected women in Madurai, South India. J AIDS Clin Res 7: 604.
2. Birlie B, Diriba TA, Sisay K, Gurmessa A, Seyoum D, et al. (2016) Mother to child HIV transmission and its predictors among HIV-exposed infants: A retrospective follow-up study in Southwest Ethiopia. J AIDS Clin Res 7: 605.
3. Tincho MB, Gabere MN, Pretorius A (2016) *In silico* identification and molecular validation of putative antimicrobial peptides for HIV therapy. J AIDS Clin Res 7: 606.
4. Girma T, Tolera D, Melaku F (2016) Factors affecting voluntary counseling and testing VCT among Ambo secondary school students, West Shoa, Ethiopia. J AIDS Clin Res 7: 607.
5. Mapara EM (2016) Picturing AIDS: Using colour pictures to raise global awareness. J AIDS Clin Res 7: 611.
6. Applewhite S (2016) An HIV prevention and treatment strategy for serodiscordant black men who have sex with men couples: Using pre-exposure prophylaxis (PrEP) and antiretroviral treatment (ART) to reduce infections in the present and the future. J AIDS Clin Res 7: 609.
7. Swartz L, Erasmus G (2016) LGBTI-psychosocial services and support adolescents and young people in Gauteng, Pretoria South Africa: Profiling the psychosocial support services of OUT (Servicing the lesbian, gay, bisexual and transgender community). J AIDS Clin Res 7: 608.
8. Raheel H (2016) Stigma towards people living with HIV/AIDS (PLWAs) among adolescents of Riyadh, Kingdom of Saudi Arabia. J AIDS Clin Res 7: 612.
9. Avanzi VM, Vicente BA, Gomes-da-Silva MM, Vidal LRR, Nogueira MB, et al. (2016) IL28B and ITPA single nucleotide polymorphisms in a cohort of patients with HCV and HIV, Southern Brazil. J AIDS Clin Res 7: 610.
10. Saleem K, Azim W (2016) Patterns of presentation in gonococcal urethritis and assessment of antibiotic sensitivity in gonorrhea. J AIDS Clin Res 7: 613.
11. Ramdial PK, Sing Y, Ramburan A, Bagratee JS, Naidu TK, et al. (2016) Nodal donovanosis as the sentinel clue to acquired immunodeficiency syndrome. J AIDS Clin Res 7:614.

\*Corresponding author: Purushottam Narute, PhD, Critical Care Medicine Department, National Institutes of Health Clinical Center, Building 10, 4D10, 10 Center Drive, Bethesda, MD, 20892, USA, Tel: 301-402-7745; E-mail: [Purushottam.narute@gmail.com](mailto:Purushottam.narute@gmail.com)

Received October 10, 2016; Accepted October 11, 2016; Published October 17, 2016

Citation: Narute PS (2016) Editor's Note: Volume 7 Issue 9. J AIDS Clin Res 7: e123. doi:10.4172/2155-6113.1000e123

Copyright: © 2016 Narute PS. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.