

Foliar Phosphorus and Zinc Application Improve Growth and Productivity of Maize (*Zea mays* L.) Under Moisture Stress conditions in Semi-Arid Climates

Amanullah^{1*}, Amir Saleem¹, Asif Iqbal¹ and Shah Fahad²

¹Department of Agronomy, Faculty of Crop Production Sciences, The University of Agriculture, Peshawar, Pakistan

²Crop Physiology and Production Center (CPCC), College of Plant Science and Technology Huazhong Agricultural University, Wuhan, Hubei, PR China

Abstract

Foliar fertilizers application is beneficial for improving growth, yield and yield components of field crops under moisture stress condition in semiarid climates. Field experiment was conducted to study the response of dryland maize (*Zea mays* L., cv. Azam) to foliar phosphorus (1, 2 and 3% P) and zinc levels (0.1, 0.2, and 0.3% Zn) and their application time (T_1 =at boot stage and T_2 =at silking stage) at the Agronomy Research Farm of The University of Agriculture Peshawar during summer 2014. The experiment was conducted under moisture stress condition (with three irrigations only: 1st at emergence, 2nd at knee height and third at seed development stage). The results revealed that foliar treated plots (rest) had significantly ($P<0.05$) better growth, higher yield components and grain yield than control (no foliar spray). Plots applied with 3% foliar P had improved growth and resulted in significantly ($P<0.05$) higher yield and yield components. Yield and yield components of maize increased significantly ($P<0.05$) with 0.3% foliar Zn application. Maize growth, yield and yield components also increased significantly ($P<0.05$) when foliar nutrients were applied at early stage (booting) than late (silking) application. It was concluded from this study that application of 3% foliar P+0.3% foliar Zn at boot stage improve growth and increase maize productivity and profitability under moisture stress condition in semiarid climates.

Keywords: Maize; Growth; Yield; Foliar; Phosphorus; Zinc; Phosphorus × zinc; Limited irrigation

Introduction

Maize (*Zea mays* L.) is the 3rd highest yielding cereal crop in the world after wheat and rice. In Pakistan 64% maize is grown in irrigated and 36% in rainfed areas. In northwest Pakistan (Khyber Pakhtunkhwa) having semiarid climate maize yield is very low 1868 kg ha⁻¹ [1]. Improper water and nutrients management are the two main factors that adversely affect the growth and crop productivity under moisture stress (drought) condition in the Khyber Pakhtunkhwa (province of Pakistan [2]. Balance fertilization is essential for improving crop growth [3], increasing yield and harvest index [4], grain quality [5,6], fertilizer use efficiency [7] and profitability [8-11].

The diffusion coefficient of soil applied phosphorus (P) is very low and plants cannot get P when needed [12] particularly in the calcareous soils under semiarid condition [4]. Therefore, foliar P application is very important under semiarid climate. The mechanistic processes by which foliar applied nutrients are taken up are through leaf stomata [13] and hydrophilic pores within the leaf cuticle [14]. Foliar applications of KH_2PO_4 is reported to delay leaf senescence and increase winter wheat grain yields during hot and dry summers [15,16]. Increased yields in barley were obtained using dilute solutions of foliar P [17]. Foliar fertilization with nitrogen, phosphorus, and potassium (N-P-K) can be supplemented with soil applied fertilizers but cannot replace soil fertilization in the case of maize [18,19] reported that when initial P deficiency symptoms appeared 25 days after sowing in wheat, higher doses of ammonium phosphate as a foliar spray gave the greatest reduction in P deficiency and highest yields. Foliar P can increase fertilizer use efficiency [20].

Zinc (Zn) is very important for various physiological functions in plants [21-24]. Zinc deficiency not only reduces the crop production but also cause Zn deficiency in our diet [25]. Application of Zn to zinc deficient soils increased maize grain yield as well as the Zn and N concentrations in maize grains. Application of Zn increases dry

matter by increasing leaf chlorophyll contents [26] and increase in N and P efficiencies [27]. In soils the P interferes Zn uptake by the plants [2,28]. About the interaction of zinc and phosphorus numerous studies have been done and all confirms this point that zinc and phosphorus imbalance in the plant, as a result excessive accumulation of phosphorus, causing zinc imposed deficiency [29]. Therefore, foliar Zn application is a simple way for making quick correction of plant nutritional status in wheat [30] and maize [31]. Foliar application of Zn enhances the uptake and accumulation of nitrogen and finally increased the maize grain. Foliar application of Zn significantly improved starch contents of forage maize [32]. In case of greater bioavailability of the grain zinc derived from foliar applications than from soil, agronomic bio-fortification would be a very attractive and useful strategy in solving zinc deficiency related health problems globally and effectively [24,33,34]. Therefore, foliar application of nutrients to crop plants is very effective for improving crop growth and yield [35]. However, there is lack of research on crops response to combined use of foliar P and Zn. The present study was therefore designed to investigate impact of various foliar P (1, 2 and 3%) and foliar Zn levels (0.1, 0.2 and 0.3%) applied at different growth stages (T_1 =boot stage and T_2 =silking stage) on the growth and yield of maize (*Zea mays* L., cv. Azam) under limited irrigation in semi-arid condition.

***Corresponding author:** Amanullah, Department of Agronomy, Faculty of Crop Production Sciences, The University of Agriculture, Peshawar, Pakistan, Tel: +92 91 9221144; E-mail: amanullah@aup.edu.pk

Received July 01, 2016; **Accepted** September 30, 2016; **Published** October 07, 2016

Citation: Amanullah, Saleem A, Iqbal A, Fahad S (2016) Foliar Phosphorus and Zinc Application Improve Growth and Productivity of Maize (*Zea mays* L.) Under Moisture Stress conditions in Semi-Arid Climates. J Microb Biochem Technol 8: 433-439. doi: 10.4172/1948-5948.1000321

Copyright: © 2016 Amanullah, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Materials and Methods

To study the effect of foliar phosphorus (P) and zinc (Zn) levels on growth and yield of maize (*Zea mays* L., cv. Azam), field experiment was conducted at the Agronomy Research Farm of The University of Agriculture Peshawar, during summer, 2014. The experiment consisted of three foliar P levels ($P_1=1$, $P_2=2$ and $P_3=3\%$), three foliar Zn levels ($Zn_1=0.1$, $Zn_2=0.2$ and $Zn_3=0.3\%$), two application times (T_1 =boot stage and T_2 =silking stage) and one control ($P_0 Zn_0$). The experiment was laid out in a randomized complete block design having three replications. Each replication consisted of 10 treatments per replication with a plot size of 3 m × 3 m (4 rows, 3 m long and 70 cm apart) at the rate of 66,000 plants ha⁻¹. Local variety of maize “Azam” was used as test crop and sown on 29th June, 2014. Data were recorded on plant height, leaf area index, number of grains ear⁻¹, thousand grains weight, biomass yield, grain yield and shelling percentage. Data on plant height (cm) at physiological maturity was recorded with the help of meter rod by selecting five plants randomly from each plot and then average plant height was calculated. Leaf area index at silking was calculated as leaf area plant⁻¹ (mean single leaf area × leaves plant⁻¹) divided by ground area per plant. Number of grains ear⁻¹ was calculated on ten randomly selected ears from each plot and then averaged. Thousand grains weight were calculated by taking of randomly 1000 grains from seed lot of each plot and was weighted with the help of electronic balance. Data on biomass yield was recorded by harvesting the four central rows in each plot, the material was sun dried for several days and weighed, and then converted into biomass yield in kg ha⁻¹. For grain yield, the four central rows of each treatment harvested for biomass yield, the ears were separated, dried, threshed, grains were cleaned and weighed and then converted into grain yield (kg ha⁻¹). Shelling percentage was calculated by using the formula [36,37]:

$$\text{Shelling percentage (\%)} = \frac{\text{Grains weight of 10 ears} \times 100}{\text{Total weight of 10 ears}}$$

Statistical Analysis

The data were statistically analyzed according to Steel et al. [38] for randomized complete block design and means among different treatment were composed using least significant differences (LSD) test ($P \leq 0.05$). Brief ANOVA and level of significance is given in Table 1.

Results

Growth

Plant height (cm) was significantly ($P < 0.05$) affected by control vs. rest (average of all foliar P and Zn treated plots), application time (T), foliar Zn, foliar P, Zn × P and T × P (Table 1). The rest had produced significantly taller plants (187 cm) over control (180 cm) (Table 2). Maize produced taller plants with early (boot stage) foliar application than late (silking stage) application. Among the foliar P levels, plant height was increased with 3% P (189.2 cm) and lower plant height (184.3 cm) was obtained with 1% P. Among the foliar Zn levels, the tallest plants were recorded with 0.3% Zn (188.1 cm) and minimum (185 cm) with 0.2% foliar Zn. Interaction between P × Zn indicated that at lowest foliar P level (1 % P); all the foliar Zn levels resulted in shorter plants (Figure 1). At the highest foliar P (3%), application of 0.1% and 0.3% foliar Zn increased plant height over 0.2% foliar Zn (Figure 1). However, 0.2% foliar Zn increased plant height over 0.1% and 0.3 foliar Zn when applied in combination with 2% P (Figure 1). Interaction between P × T indicated that the highest foliar P level (3%) resulted in the tallest plant heights when applied at boot stage (Figure 2). Application of 2% foliar P was better in terms of taller plants when




Figure 1: Interaction between foliar P and foliar Zn (P × Zn) influence plant height (cm) in maize (*Zea mays*).




Figure 2: Interaction between foliar P and application time (P × T) influence plant height (cm) in maize (*Zea mays*).

applied at silking stage (Figure 2). However, no difference in plant heights was observed when 0.1% foliar Zn was applied either at booting or silking (Figure 2).

Leaf area index (LAI) was significantly ($P < 0.05$) affected by control vs. rest, foliar P and Zn levels (Table 1). The rest had produced significantly higher LAI over control (Table 2). Maximum LAI was obtained with 3% foliar P (2.7) and minimum (2.5) was obtained with 1% foliar P. Among the foliar Zn levels, maximum LAI was obtained with 0.3% Zn (2.8) and lowest with 0.1% foliar Zn (2.5).

Yield components

Number of grains ear⁻¹ was significantly ($P < 0.05$) affected by control vs. rest, foliar P and Zn levels, and P × Zn interaction (Table 1). The rest had produced significantly more number of grains ear⁻¹ (456) as compared to control (321) (Table 2). Among the foliar P levels, the highest number of grains ear⁻¹ was obtained with 2% P (478.5) and lower with 3 % P (429.6). Among the foliar Zn levels, maximum number of grains ear⁻¹ were recorded with 0.2% Zn (498.7) and lowest number of grains ear⁻¹ were obtained with 0.3% Zn (411.8). Interaction between P × Zn indicated that at lowest foliar P level (1% P); application of 0.2% foliar Zn increased number of grains ear⁻¹ than other P and Zn

Source of Variance	Plant height (cm)	Leaf area index	Grains ear ⁻¹	1000 grains weight (g)	Biomass yield (kg ha ⁻¹)	Grain yield (kg ha ⁻¹)	Shelling (%)
Control vs. Rest	***	***	***	***	***	***	***
Application time (T)	*	*	*	*	***	***	***
Zinc (Zn)	**	***	**	*	***	***	***
Phosphorus (P)	***	***	**	*	*	***	***
P × Zn	***	ns	*	*	ns	**	ns
Zn × T	ns	ns	ns	ns	**	***	*
P × T	***	ns	ns	ns	ns	ns	ns
P × Zn × T	ns	ns	ns	ns	ns	ns	ns

*, **, *** indicates that data is significant at 5, 1 and 0.1% level of probability, respectively. The word ns stand for the non-significant data at 5% level of probability

Table 1: Analysis of variance for plant height (cm), leaf area index, number of grains ear⁻¹, thousand grains weight, biomass yield (kg ha⁻¹), grain yield (kg ha⁻¹) and shelling percentage (%) of maize as affected by foliar phosphorus (P) and zinc (Zn) levels and their application time (T).

Treatments	Plant height (cm)	Leaf area index	Number of Grains ear ⁻¹	Thousand Grains Weight (g)	Biomass yield (kg ha ⁻¹)	Grain yield (kg ha ⁻¹)	Shelling Percentage (%)
P 1%	184.3 c	2.5 c	460.5 b	213.50 b	9471 b	3560 c	75.8 c
P 2%	186.6 b	2.6 b	478.5 a	216.60 a	9544 ab	3624 b	77.1 b
P 3%	189.2 a	2.7 a	429.6 c	205.50 c	9567 a	3708 a	78.2 a
SD	± 2.5	± 0.1	± 24.7	± 5.7	± 50.1	± 74.2	± 1.2
Zn 0.1%	186.1 b	2.5 b	458.2 b	206.90 b	9321 c	3446 c	73.9 c
Zn 0.2%	185.9 b	2.5 b	498.7 a	217.70 a	9506 b	3603 b	76.4 b
Zn 0.3%	188.1a	2.8 a	411.8 c	211.00 b	9754 a	3842 a	80.8 a
SD	± 1.2	± 0.2	± 43.5	± 5.5	± 217.3	± 199.4	± 3.5
Boot stage	187.9 a	2.7 a	441.7 b	211.00 a	9779 a	3698 a	78.6 a
Silking stage	185.5 b	2.5 b	470.7 a	212.70 a	9276 b	3563 b	75.5 b
SD	± 1.7	± 0.1	± 20.5	± 1.2	± 355.7	± 95.5	± 2.2
Control plots	180.0 b	2.0 b	321.2 b	213.50 b	8841 b	3289 b	72.0 b
Treated plots	187.0 a	3.0 a	456.4 a	216.60 a	9527 a	3631 a	77.0 a
SD	± 4.9	± 0.7	± 95.6	± 2.2	± 485.1	± 241.8	± 3.5

Where SD stands for standard deviations

Means of the same category followed by different letters are significantly different from each other using LSD test (P ≤ 0.05)

Table 2: Plant height (cm), leaf area index, number of grains ear⁻¹, thousand grains weight, biomass yield (kg ha⁻¹), grain yield (kg ha⁻¹) and shelling percentage (%) of maize as affected by foliar phosphorus (P) and zinc (Zn) levels and their application time (T).




Figure 3: Interaction between foliar P and foliar Zn (P × Zn) influence number of grains ear⁻¹ in maize (*Zea mays*).

combinations (Figure 3). The combined application of 2% foliar P+0.1% foliar Zn stood second in the ranking in terms of higher number of grains ear⁻¹ (Figure 3).



Thousand grains weight (g) was significantly (P<0.05) affected by control vs. rest, foliar P and Zn levels, and P × Zn interaction (Table 1). The rest had significantly higher thousand grains weight (216.6 g) as compared to control (213.5 g) (Table 2). Among the foliar Zn levels,

application of 0.2% Zn gave highest thousand grains weight (217.7 g), while 0.1% Zn gave lowest thousand grains weight (206.9 g). Among the foliar P levels, the highest thousand grains weight was obtained with 2% P (216.6 g), while the lowest thousand grains weight was recorded with 3% P (205.5 g). Interaction between P × Zn indicated that combination of 2% foliar P+0.2% foliar Zn increased 1000 grains weight in maize than any other foliar P and Zn combinations (Figure 4). The combined application of 1% foliar P+0.2% foliar Zn stood second in the ranking in terms of heavier grains (Figure 4).

Yield and shelling percentage

Biomass yield (kg ha⁻¹) was significantly (P<0.05) affected by control vs. rest, foliar P and Zn levels, and interaction between foliar P and application time (P × T) (Table 1). The rest plots had significantly higher biomass yield (9527 kg ha⁻¹) as compared to control (8841 kg ha⁻¹) (Table 2). Among the foliar P levels, the highest biomass yield was obtained with 3% P (9566 kg ha⁻¹) and the lowest biomass yield was obtained with 1% P (9470 kg ha⁻¹). Among the foliar Zn levels, the highest biomass yield was obtained with 0.3% Zn (9754 kg ha⁻¹) and the lowest biomass yield (9321 kg ha⁻¹) with 0.1% foliar Zn. The interaction between P × T indicated that the biomass yield in maize increased significantly when any of the three P levels (1, 2 and 3% P) was applied at the early growth stage (booting) than applied late at silking stage of maize (Figure 5).




Grain yield (kg ha⁻¹) of maize was significantly (P<0.05) affected by control vs. rest, foliar P and Zn levels, and interactions between P



$\times Zn$ and $P \times T$ (Table 1). The rest had significantly higher grain yield (3631 kg ha⁻¹) as compared to control (3289 kg ha⁻¹) (Table 2). Among the foliar Zn levels, the highest grain yield (3842 kg ha⁻¹) was obtained with 0.3% Zn, while the lowest grain yield (3446 kg ha⁻¹) was obtained with 0.1% foliar Zn. Among the foliar P levels, the highest grain yield (3707 kg ha⁻¹) was obtained with 3% P, while the lowest grain yield (3560 kg ha⁻¹) was obtained with 1% P. The interaction between $P \times Zn$ indicated that the grain yield in maize increased significantly with the increase in both foliar P and Zn levels and vice versa (Figure 6). The interaction between $P \times T$ indicated that the grain yield in maize increased significantly when any of the three P level (1, 2 and 3% P) was applied at the early growth stage (booting) than applied late at silking stage of maize (Figure 7).

Shelling percentage (%) was significantly ($P < 0.05$) affected by control vs. rest, foliar P and Zn levels, and $Zn \times T$ interaction (Table 1). The rest had significantly higher shelling (77.0%) as compared to control (72.0%) (Table 2). Among the foliar P levels, maximum shelling percentage was obtained with 3% P (78.2%), while the lowest shelling percentage was obtained with 1% P (72.2%). Among the foliar Zn levels, maximum shelling percentage was obtained with 0.3% Zn (80.8%) and the lowest shelling percentage was obtained with 0.1% Zn (73.9%). The interaction between $P \times T$ indicated that the shelling percentage in

maize increased significantly when any of the three P levels (1, 2 and 3% P) was applied at the early growth stage (booting) than applied late at silking stage of maize (Figure 8) and showed positive relationship with increase in grain yield.



Discussion

Both foliar P and Zn application improved the growth parameters (plant height and LAI) over control (no foliar P and Zn application) under calcareous soils in semiarid condition. Amanullah et al. [2] reported significant variation ($P < 0.05$) in plant height and LAI of maize while spraying NPK at different growth stages. The increase in plant height with P and Zn application was earlier reported by many researchers [39-46]. Similarly, the increase in LAI in wheat with P and Zn application was earlier reported by Shukla and Warsi [47]. Other researchers [48]; Mohsin et al. [46] reported increase in the LAI of maize with foliar Zn application. The increase in LAI with Zn application probably may be due to increase in tryptophan amino acid and indole acetic acid hormone which are two main factors in leaf area expansion [49]. According to Firouzi [50], foliar spray of Zn increase LAI due to an increase in length and width of leaves with increase in auxin. Amanullah et al. [6] reported that foliar N application time had significant ($P < 0.05$) effects, while foliar N source had no significant ($P > 0.05$) effects on mean single leaf area of maize. Amanullah et al. [6] reported that foliar spray of N+P+K (2% each) resulted in maximum mean single leaf area, followed by foliar N application alone and the lowest mean single leaf area was observed with sole foliar P.

The yield components (number of grains ear⁻¹ and 1000 grains weight) also showed positive and significant ($P < 0.05$) response to foliar P and Zn application over control. Earlier, Amanullah et al. [6] found that number of grains ear⁻¹ in maize increased with foliar spray of NPK in combination over control. The increase in grains ear⁻¹ with foliar P application are in agreement with those of Alston [51], they reported that grains spike⁻¹ of wheat significantly increased with foliar P. Similarly, Soleimani [52] reported marked increase in number of grains spike⁻¹ of wheat for foliar application Zn. The tendency of increasing number of grains ear⁻¹ with Zn application was also in line with the findings of many workers [53,54]. The increase in 1000 grains weight in maize with foliar nutrition was in accordance with our early study [6]. According to Samad et al. [55], the combine application of soil and foliar applied P increased 1000-grains weight over control but did not improve it over the sole application of soil or foliar applied P. Tahir et al. [43] and Mohsin et al. [46], reported increase in the 1000-grains weight with foliar Z application over control. These results are in line with the results of Guenis et al. [56], Soyly et al. [57], Harris et al. [54] and Farajzadeh et al. [58]; they reported significant increase in thousand grains weight with foliar application of micronutrients.

Both biomass and grain yields as well as shelling increased significantly ($P < 0.05$) with foliar P and Zn application over control and the increase was more when foliar nutrients were applied at boot stage than applied at silking. The increase in biomass yield with foliar nutrition was attributed to the improvement in growth parameters, yield and yield components of maize. According to Amanullah et al. [6], combined foliar application of the three major nutrients (N+P+K) at the rate of 1% each in two equal splits at 30 and 60 days after emergence increased maize productivity under moisture stress condition under semiarid condition. Many researchers [20,59] reported increase in grain and biomass yields with foliar P. Other researchers [46,60] showed positive response of biomass and grain yields to foliar Zn application. The increase in biomass yield with foliar application of Zn probably might be due to their critical role of Zn in crop growth, involving in photosynthesis processes, respiration and other biochemical and physiological activates and thus their importance in achieving higher yields [61]. Higher yield due to Zn fertilization was also attributed to the enhanced synthesis of carbohydrates and their transport to the site of grain production [62-65].

Conclusion

The foliar P and Zn treated plots (rest) had better performance in terms of improved growth, higher yield and yield components than control (no foliar spray). Application of foliar P at the rate of 3% and Zn at the rate of 0.3% improved growth, increased yield and yield components of maize under moisture stress condition. Early application of foliar P and Zn at boot stage had positive impact on growth, yield and yield components of maize as compared to late spray at silking.

References

1. MINFAL (2012) Agricultural statistics of Pakistan. Government of Pakistan. Ministry of Food Agri. Livestock, Food Agricultural and Livestock, Division (Economic wing) Islamabad.
2. Amanullah, Inamullah (2014) Preceding rice genotypes, residual phosphorus and zinc influence crop productivity under rice-wheat cropping system. 4th International & 13th National Conference of Botany at SBB University, Sheringal, Upper Dir, Khyber Pakhtunkhwa.
3. Amanullah (2014a) Effects of variable nitrogen source and rate on leaf area index and total dry matter accumulation in maize (*Zea mays* L.) genotypes under calcareous soils. Turkish J Field Crops 19: 276-284.
4. Amanullah M, Asif SS, Malhi RA, Khattak (2009) Effects of P-fertilizer source and plant density on growth and yield of maize in Northwestern Pakistan. J Plant Nutr 32: 2080-2093.
5. Amanullah M, Shah P (2010) Timing and rate of nitrogen application influence grain quality and yield in maize planted at high and low densities. J Sci Food & Agric 15: 21-29.
6. Amanullah A, Khan Z, Khan F (2013) Foliar application of nitrogen at different growth stages influence the phenology, growth and yield of maize (*Zea mays* L.). Soil & Environment 32: 135-140.
7. Amanullah (2014b) Source and rate of nitrogen application influence agronomic N-use efficiency and harvest index in maize (*Zea mays* L.) genotypes. Maydica 59: 80-89.
8. Amanullah LK, Almas, Shah P (2010a) Timing and rate of nitrogen application influence profitability of maize planted at low and high densities in Northwest Pakistan. Agron J 102: 575-579.
9. Amanullah M, Zakirullah, Khalil SK (2010b) Timing and rate of phosphorus application influence maize phenology, yield and profitability in Northwest Pakistan. Intl J Plant Production 4: 281-292.
10. Amanullah M, Khan W (2011) Interactive effect of potassium and phosphorus on grain quality and profitability of sunflower in Northwest Pakistan. Pedosphere. 21: 532-538.
11. Amanullah M, Asif LK, Almas (2012) Agronomic efficiency and profitability of P-fertilizers applied at different planting densities of maize in Northwest Pakistan. J Plant Nutr 35: 331-341.
12. Clarkson DT (1981) Nutrient interception and transport by root systems. Butterworths 307-330.
13. Eichert TJ, Burkhardt J (1999) A novel model system for the assessment of foliar fertilizer efficiency. In Technology and Applications of foliar fertilizers. Proceedings of the Second International Workshop on Foliar Fertilization, Bangkok.
14. Tyree M, Scherbatskoy D, Tabor CA (1990) Leaf cuticles behave as asymmetric membranes: Evidence from measurement of diffusion potentials. Plant Physiology 92: 103-109.
15. Sherchand K, Paulsen GM (1985) Response of wheat to foliar phosphorus treatments under field and high temperature regimes. J Plant Nutri. 12: 1171-1181.
16. Batten GD, Wardlaw IF, Aston MJ (1986) Growth and the distribution of phosphorus in wheat developed under various phosphorus and temperature regimes. Aus J Agric Res 3: 459-469.
17. Qaseem SM, Afridi MM, Samiullah RK (1978) Effect of leaf applied phosphorus on the yield characteristics of ten barley varieties. Indian J Agric Sci 48: 215-217.
18. Ling F, Silberbush M (2002) Response of maize to foliar versus soil application of nitrogen-phosphorus-potassium fertilizers. J Plant Nutri 11: 2333-2342.

19. Haloi B (1980) Effect of foliar application of phosphorus salt on yellowing of wheat seedlings. *J Res Assam Agric Uni* 1: 108-109.
20. Dixon RC (2003) Foliar fertilization improves nutrient use efficiency. *Fluid Journal* 11: 22-23.
21. Outten CE, O'Halloran TV (2001) Femtomolar sensitivity of metalloregulatory protein controlling Zn homeostasis. *Science* 292: 2488-2492.
22. Pandey N, Pathak GC, Sharma CP (2006) Zinc is critically required for pollen function and fertilization in lentil. *J Trace Elements in Medicine and Biol* 20: 89-96.
23. Alloway B (2004) Zinc in soils and crop nutrition. *Areas of the World with Zinc Deficiency Problems*.
24. Cakmak I (2008) Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant and Soil* 302: 1-17.
25. Bagei SA, Ekiz H, Yilmaz A, Cakmak I (2007) Effects of zinc deficiency and drought on grain yield of field-grown wheat cultivars in Central Anatolia. *J Agron Crop* 193: 198-206.
26. Sharafi S, Tajbakhsh M, Majidi M, Pourmirza A (2002) Effect of iron and zinc fertilizer on yield and yield components of two forage corn cultivars in Urmia. *Soil and Water* 12: 85-94.
27. Mahaleh JK, Roshdi M (2008) Effect of low-used elements spraying on qualitative and quantitative properties of silo corn 704 in Khoy. *Seed and Plant* 24: 281-293.
28. Salimpour S, Khavazi K, Nadian H, Besharati H, Miransari M (2010) Enhancing phosphorus availability to canola (*Brassica napus* L.) using P solubilizing and sulfur oxidizing bacteria. *Aust J Crop Sci* 4: 330-334.
29. Khorgamy A, Farnis A (2009) Effect of phosphorus and zinc fertilisation on yield and yield components of chick pea cultivars. *African Crop Science Conference Proceedings* 9: 205-208.
30. Erenoglu B, Nikolic M, Römhold V, Cakmak I (2002) Uptake and transport of foliar applied zinc in bread and durum wheat cultivars differing in zinc efficiency. *Plant and Soil* 241: 251-257.
31. Grzebisz W, Wrońska M, Diatta JB, Dullin P (2008) Effect of zinc foliar application at early stages of maize growth on patterns of nutrients and dry matter accumulation by the canopy. Part I. Zinc uptake patterns and its redistribution among maize organs. *J Elementolgy* 13: 17-28.
32. Leach KA, Hameleers A (2001) The effects of a foliar spray containing phosphorus and zinc on the development, composition and yield of forage maize. *Grass Forage Sci* 56: 311-315.
33. Yosefi K, Galavi M, Ramrodi M, Mousavi SR (2011) Effect of bio-phosphate and chemical phosphorus fertilizer accompanied with micronutrient foliar application on growth, yield and yield components of maize (Single Cross 704). *Aust J Crop Sci* 5: 175-180.
34. Abd El-Baky, MMH, Ahmed AA, El-Nemr MA, Zaki MF (2010) Effect of potassium fertilizer and foliar zinc application on yield and quality of sweet potato. *Res J Agric Biol Sci* 6: 386-394.
35. Aghtape AA, Ghanbari A, Sirousmehr A, Siahsar B, Asgharipour M, et al. (2011) Effect of irrigation with wastewater and foliar fertilize application on some forage characteristics of foxtail millet. *Int J Plant Physiol Biochem* 3: 34-42.
36. Amanullah, Khalid S (2016) Integrated use of phosphorus, animal manures and biofertilizers improve maize productivity under semiarid condition. *Organic Fertilizers - From Basic Concepts to Applied Outcomes*. InTech, Rijeka, Croatia.
37. Amanullah, Khan A (2015) Phosphorus and compost management influence maize (*Zea mays*) productivity under semiarid condition with and without phosphate solubilizing bacteria. *Frontiers in Plant Science (Plant Biotic Interactions)* 6: 1083.
38. Steel RGD, Torrie JH, Dickey D (1996) *Principals and procedures of statistics*. McGraw Hill, USA.
39. Kausar K, Akbar M, Rasul E, Ahmad AN (1993) Physiological response of nitrogen, phosphorus and potassium on growth and yield of wheat. *Pakistan. J Agric Res* 14: 126-130.
40. Singaram P, Kothandaraman GV (1994) Studies on residual, direct and cumulative effect of phosphorus sources on the availability, content and uptake of phosphorus and yield of maize. *Madras Agric J* 81: 425- 429.
41. Allmarus RR, Pumphrey EV, Cochlar FE, Robert S (1998) Method and rate of supplying ZnSO₄ for corn on Zn deficient soil in west Nebraska. *Agron J* 53: 236-238.
42. Sahoo SC, Panda M (2001) Effect of phosphorus and detasseling on yield of babycorn. *Indian J Agri Sci* 71: 21-22.
43. Tahir M, Fiaz N, Nadeem MA, Khalid F, Ali M (2009) Effect of different chelated zinc sources on the growth and yield of maize (*Zea mays* L.). *Soil Environ*. 28: 179-183.
44. El-Badawy, El M, Mehasen SAS (2011) Multivariate analysis for yield and its components in maize under zinc and nitrogen fertilization levels. *Aust J Basic & Appl Sci* 5: 3008-3015.
45. Piri I (2012) Effect of phosphorus fertilizer and micronutrients foliar application on sorghum yield. *Annals. Biol Res* 3: 3998-4001.
46. Mohsin AU, Ahmad M, Farooq, Ullah S (2014) Influence of zinc application through seed treatment and foliar spray on growth, productivity and grain quality of hybrid maize. *J Animal & Plant Sci* 24: 1494-1503.
47. Shukla SK, Warsi AS (2000) Effect of sulphur and micronutrients on growth, nutrient content and yield of wheat (*Triticum aestivum*) *Indian J. Agri. Res.* 34: 203-205.
48. Safyan N, Naderidarbaghshahi MR, Bahari B (2012) The effect of microelements spraying on growth, qualitative and quantitative grain corn in Iran. *Intl R J Appl Basic Sci* 3: 2780-2784.
49. Seifi-Nadergholi M, Yarnia M, Rahimzade KF (2011) Effect of zinc and manganese and their application method on yield and yield components of common bean (*Phaseolus vulgaris* L. CV. Khomein). *Middle-East J. Sci. Res.* 8: 859-865.
50. Firouzi M (2005) The effect of foliar absorption of microelements on growth and yield of foliage corn. MS thesis in Agronomy, Agriculture College, Islamic Azad University, Khorasgan Branch.
51. Alston AM (1979) Effects of soil water content and foliar fertilization with nitrogen and phosphorus in late season on the yield and composition of wheat. *Aus J Exper Agric Anim Husb* 30: 577-585.
52. Soleimani R (2006) The effects of integrated application of micronutrient on wheat in low organic carbon conditions of alkaline soils of western Iran. *Soil Sci* 21: 382-385.
53. Bakyt K, Sade B (2002) Response of field grown barley cultivars grown on zinc deficient soil to zinc application. *Comm Soil Sci Plant Anal* 33: 533-544.
54. Harris D, Rashid A, Miraj G, Arif M, Shah H (2007) On farm' seed priming with zinc sulphate solution - a cost effective way to increase the maize yields of resource - poor farmers. *Field Crops Res* 102: 119-127.
55. Samad A, Muhammad D, Musarat M, Ullah W (2014) Enhancing wheat yield and phosphorus use efficiency through foliar application in calcareous soil. *J Natur Sci Res* 4: 70-74.
56. Guenis A, Alpaslan M, Unal A (2003) Effects of boron fertilization on the yield and some yield components of bread and durum wheat. *Turk J Agric* 27: 329-335.
57. Soylu SB, Sade A, Topal N, Akgun S, Gezgin (2005) Responses of irrigated durum and bread wheat cultivars to boron application in low boron calcareous soil *Turk. J Agric* 29: 275-286.
58. Farajzadeh MT, Khorshidi MB, Ahmadzadeh V (2009) Effect of micronutrients and their application method on yield, crop growth rate and net assimilation rate of corn. *J Food Agric Envir* 7: 611-615.
59. Poulsen KH, Nagy R, Gao LL, Smith SE, Bucher M, et al. (2005) Physiological and molecular evidence for P uptake via the symbiotic pathway in reduced mycorrhizal colonization with a compatible fungus. *New Phytologist*. 168: 445-453.
60. Thalooth M, Tawfik M, Mohamed MH (2006) A comparative study on the effect of foliar application of zinc, potassium and magnesium on growth, yield and some chemical constituents of mung bean plants growth under water stress conditions. *World J Agric* 2: 37-46.
61. Zeidan MS, Manal F, Hamouda HA (2010) Effect of foliar fertilization of Fe, Mn and Zn on wheat yield and quality in low sandy soils fertility. *World J Agric Sci* 6: 696-699.

62. Pedda-Babu P, Shanti M, Prasad BR, Minhas PS (2007) Effect of zinc on rice in rice – black gram cropping system in saline soils. Andhra Agric J 54: 47-50.
63. Amanullah K, Kakar M, Khan A, Khan I, Shah Z (2014) Growth and yield response of maize (*Zea mays* L.) to foliar NPK-fertilizers under moisture stress condition. Soil & Environ 33: 116-123.
64. Amanullah M, Yasir SK, Khalil MT (2010) Phenology, growth and grain yield of maize as influenced by foliar applied urea at different growth stages. J Plant Nutri 33: 71-79.
65. Girma K, Raun B, Zhang H, Mosali J (2006) What about foliar P on corn and winter wheat. Fluid J 14: 17-19.

Citation: Amanullah, Saleem A, Iqbal A, Fahad S (2016) Foliar Phosphorus and Zinc Application Improve Growth and Productivity of Maize (*Zea mays* L.) Under Moisture Stress conditions in Semi-Arid Climates. J Microb Biochem Technol 8: 433-439. doi: [10.4172/1948-5948.1000321](https://doi.org/10.4172/1948-5948.1000321)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid peer review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled for better prominence and citations
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Best discounts for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/jmbt>