

Molecular basis of anti-inflammatory strategies in cancer cachexia

Martins Thomas

Cardiothoracic Surgery Unit, College of Medicine of University of Lagos, Nigeria

Background: There are newer diagnostic and therapeutic armamentaria for primary lung cancer. Application of molecular genetics in lung cancer management is evolving rapidly. However, the traditional knowledge and practices that were applicable before the 1980s still hold sway in most developing countries.

Aims and Objectives: This research was conducted to highlight the staggering gap in the current aetiopathology and management profile of primary lung cancers and to assess the readiness of developing world for the challenges of lung cancer management in the new decade.

Methods: We studied the patients referred to Lagos University Teaching Hospital with suspicion of primary lung cancer. We noted their bio-data, predisposing factors and final diagnosis on completion of investigations. We also noted the therapeutic modalities that were applied - especially the type of operation that was done for each of the patients.

Results: The research lasted 99 months beginning in October 1999 and 267 patients were enlisted. There were 148 males (55.4%) and 119 females (44.5%). Stage IV patients were 183 (68.5%) while only 3 patients were found at stage I. Histology showed squamous cell carcinoma in 27.7% of cases while adenocarcinoma constituted 64.0%. Curative surgery was performed for 13.1% while non curative surgery was performed for 16.5%. Correlation between smoking and malignancy was stronger among the males than the female patients.

Discussion and Conclusion: There is increasing incidence of primary lung cancers among non-smoking females.