


# 3<sup>rd</sup> International Conference and Exhibition on **Obesity & Weight Management**

December 01-03, 2014 DoubleTree by Hilton Hotel San Francisco Airport, USA



## *Yoonjung Park*

University of Houston, USA

### **Role of physical activity in obesity-induced vascular dysfunction in heart**

The purpose of this study was to investigate whether physical activity initiated with the start of high-fat feeding would primordially prevent development of endothelial dysfunction, and if it does, then to determine some potential mechanisms. C57BL/6 female mice were randomly divided into three groups: 1) control low-fat diet (LF-SED; 15% of calories from fat), 2) high-fat diet (HF-SED; 45% of calories from fat), and 3) HF diet given access to a voluntary running wheel (HF-RUN). Our hypothesis was that HF-RUN would differ in multiple markers of endothelial dysfunction from HF-SED after 10 weeks of 45%-fat-diet, but would did not differ from LF-SED. HF-RUN differed from HF-SED in nine determinations in which HF-SED either had decreases in 1) Acetylcholine (ACh)-induced and flow-induced vasodilations in isolated, pressurized coronary arterioles, 2) heart phosphorylated endothelial nitric oxide synthase (p-eNOS/eNOS) protein, 3) coronary arteriole leptin (ob) receptor protein, 4) phosphorylated signal transducer and activator of transcription 3 (p-STAT3/STAT3) protein, and 5) coronary arteriole superoxide dismutase 1 protein; or had increases in 6) % body fat, 7) serum leptin, 8) coronary arteriole suppressor of cytokine signaling 3 (SOCS3) protein, and 9) coronary arteriole gp91phox protein. Higher endothelium dependent-vasodilation by ACh or leptin was abolished with incubation of NOS inhibitor NG-nitro-L-arginine-methyl ester (L-NAME) in LF-SED and HF-RUN groups. Further, impaired ACh-induced vasodilation in HF-SED was normalized by apocynin or TEMPOL to LF-SED and HF-RUN. These findings demonstrate multiple mechanisms (eNOS, leptin and redox balance) by which voluntary running opposes the development of impaired coronary arteriolar vasodilation during simultaneous high-fat feeding.

### **Biography**

Yoonjung Park is currently an Assistant Professor in the Department of Health & Human Performance at the University of Houston. He received PhD degree in Cardiovascular Exercise Physiology from Texas A&M University in the department of Health and Kinesiology and completed his postdoctoral training in the Departments of Internal Medicine, Division of Cardiovascular Medicine, at University of Missouri. He has a number of publications in the field of obesity, diabetes, aging and cardiovascular disease, especially microcirculation, and has an expertise in the effects of exercise/physical activity on vascular function.

[ypark10@uh.edu](mailto:ypark10@uh.edu)