

Cardiac Imaging Modalities in the Diagnosis of Coronary Artery Disease

Zhonghua Sun*

Discipline of Medical Imaging, Department of Imaging and Applied Physics, Curtin University, Australia

Abstract

Coronary artery disease is the leading cause of death in advanced countries. Early detection and diagnosis of coronary artery disease plays an important role in the identification of disease severity and prediction of disease outcome, consequently improving patient management. Diagnosis and management of coronary artery disease is increasingly dependent on less-invasive imaging modalities, including coronary CT angiography, cardiac magnetic resonance imaging, cardiac radionuclide imaging such as SPECT and PET modalities. Rapid developments of these imaging modalities have significantly improved the diagnostic performance of each imaging technique with high diagnostic accuracy achieved in both diagnostic and prognostic value in coronary artery disease. This editorial provides an overview of the diagnostic applications of a variety of less-invasive imaging modalities in the diagnosis of coronary artery disease. This special issue of "Arteriosclerotic Vascular Disease: Part II" in the journal of Clinical and Experimental Cardiology will give particular attention to contributions focusing on the clinical applications of these imaging modalities in the arteriosclerotic vascular disease, in particular, coronary artery disease.

Keywords: Coronary artery disease; Arteriosclerotic vascular disease; Diagnostic value; Imaging modalities

Coronary artery disease (CAD) is the leading cause of death in advanced countries and its prevalence is increasing among developing countries [1,2]. Various less-invasive imaging modalities are increasingly used in the diagnosis of CAD including coronary CT angiography, cardiac magnetic resonance imaging (MRI), and cardiac single photon emission computed tomography (SPECT), positron emission tomography (PET) and integrated SPECT/CT and PET/CT [3]. To improve early diagnosis and patient management, it is essential to have an overview of the diagnostic value of different imaging modalities in CAD. This editorial provides an overview of the diagnostic performance of these imaging modalities in CAD, with a focus on the advantages, limitations and future directions of the use of each imaging modality in the diagnosis of CAD.

Coronary CT angiography represents the most rapidly developed imaging modality in cardiac imaging with evolution from single slice CT to multislice CT, from early generation of 4- and 16-slice CT to 64- and 320-slice CT scanners, demonstrating excellent visualization of coronary anatomy and assessment of coronary artery disease [4-6]. In summary, diagnostic sensitivity of coronary CT angiography has been significantly improved with 64- or more slice CT scanners when compared to the early generations of 4- and 16-slice scanners, while, the negative predictive value remains consistently high (>90%), regardless of the type of CT scanners [7-11]. This indicates the main role of coronary CT angiography is to rule out significant CAD, thus reducing the need for invasive coronary angiography. The prime indication of coronary CT angiography is to diagnose patients with a low and intermediate probability of CAD as a simple non-invasive testing, while patients with a high probability of CAD will benefit from invasive coronary angiography [12].

In addition to the diagnostic value, coronary CT angiography allows for characterization of plaque components (calcified versus non-calcified plaques and shows potential prognostic value of disease extent and cardiac events [13,14]. Studies based on single center and multicenter clinical trials have shown that coronary CT angiography provides incremental prognostic value over clinical risk analysis in predicting major adverse cardiac events with absence of CAD leading to event free survival period, while presence of plaques associated with increased risk of cardiac events [15-19].

Radiation dose associated with coronary CT angiography is the main concern of this technique in cardiac imaging, and this has increased substantially over the last decade with the development of multislice CT scanners and widespread use of cardiac CT in routine clinical practice. This has raised a serious concern and it is a hot topic of debate in the literature. Various dose-saving strategies have been proposed and recommended in the past few years to lower radiation exposure to patients undergoing coronary CT angiography with tremendous progress having been achieved. Effective dose reduction has been accomplished by employing techniques with a radiation dose of less than 10 mSv to as low as 1 mSv in some studies [11,20,21], although much effort is still required to ensure that coronary CT angiography is safely performed in imaging patients with suspected coronary artery disease.

MRI provides excellent soft tissue contrast, with inherent 3D capabilities, and acquisition of images in any anatomical plane. Furthermore, MRI does not expose the patient to ionizing radiation, thus, the usefulness of MRI has been investigated widely. However, the diagnostic accuracy of cardiac MRI in CAD varies widely according to the literature, with sensitivity ranging from 38% to 83%, and specificity ranging from 57% to 95% due to variable scanning protocols used in the studies [22]. Recent technical developments in MRI, especially with the emergence of 3.0 T MR imaging system have been shown to be a promising technique for performing cardiac MRI, with significant improvement of diagnostic value for detection of CAD [23,24]. Despite these advantages, cardiac MRI is still limited in the visualization of distal coronary segments due to inferior spatial resolution, thus, it is

*Corresponding author: Zhonghua Sun, Associate Professor, Discipline of Medical Imaging, Department of Imaging and Applied Physics, Curtin University, GPO Box, U1987, Perth, Western Australia, 6845, Australia, Tel: +61-8-9266 7509; Fax: +61-8-9266 2377; E-mail: z.sun@curtin.edu.au

Received September 10, 2013; Accepted September 11, 2013; Published September 12, 2013

Citation: Sun Z (2013) Cardiac Imaging Modalities in the Diagnosis of Coronary Artery Disease. J Clin Exp Cardiol S6: e001. doi:10.4172/2155-9880.S6-e001

Copyright: © 2013 Sun Z. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

not as widely used as coronary CT angiography in the diagnosis of CAD.

Noninvasive evaluation for obstructive CAD is performed by gatekeeper tests that offer physiologic information of coronary stenosis (physiologic imaging) or the degree of stenosis (anatomic imaging). Coronary CT angiography serves as an excellent anatomic gatekeeper as it has a very high negative predictive value, while stress perfusion cardiac MRI is regarded as a physiologic gatekeeper. Stress perfusion cardiac MRI has been proved to be a robust and accurate diagnostic test for CAD when invasive coronary angiography is used as the reference standard [25-28]. Several systematic reviews and meta-analyses have shown that the sensitivity and specificity of stress perfusion MRI ranged from 89% to 91% and 76% to 81%, using invasive coronary angiography as the reference standard [26-28]. Desai and Jha recently conducted a meta-analysis of 12 studies regarding the cardiac stress perfusion MRI in the diagnosis of flow-limiting obstructive CAD using fractional flow reserve measured at invasive coronary angiography as the reference standard [29]. Their analysis shows that cardiac stress perfusion MRI has a sensitivity of 89.1% and 87.7% and a specificity of 84.9% and 88.6% on a patient-based and on a coronary territory-based analysis, respectively. Thus, cardiac stress perfusion MRI is an accurate test for the detection of flow-limiting stenosis.

Myocardial perfusion imaging (MPI) with stress gated SPECT has been widely used in the diagnosis of CAD and is a well-documented non-invasive method for risk stratification with high diagnostic accuracy when compared to coronary CT angiography [30,31]. The presence of ischemia could be used to classify the patients as having CAD and candidates for receiving aggressive medical therapy and management. Coronary CT angiography has limited accuracy for identifying the physiologic significance of perfusion defects in patients with intermediate or high pre-test likelihood of CAD when compared to MPI SPECT [32]. Thus, MPI SPECT offers additional functional information in the evaluation of coronary stenosis. MPI SPECT can be used as the gatekeeper to invasive coronary angiography. Bateman et al. showed that referral to invasive coronary angiography was 3.5%, 9%, and 60%, respectively, corresponding to normal to mild, moderately abnormal and severely abnormal perfusion scans [33]. A negative SPECT imaging has been confirmed to serve as an excellent prognostic indicator with an annual cardiac event rate of <1% for the general population, while an increasing cardiac events are associated with increasing severity of both fixed and reversible perfusion defects, regardless of the presence of non-obstructive coronary disease [34-36].

Cardiac PET imaging is another well-established tool for the evaluation of ischemia, blood flow quantification, myocardial viability and perfusion [37,38]. Cardiac PET utilizing ¹⁸F-FDG is considered the most sensitive modality for detecting hibernating viable myocardium and predicting left ventricular functional recovery post-coronary revascularization. PET has higher spatial and temporal resolution than SPECT due to more robust methods of attenuation correction, thus, PET allows quantification of resting and hyperemic regional myocardial perfusion. When PET was integrated into clinical patient management, a significant reduction in cardiac events was observed in patients with ¹⁸F-FDG PET-assisted management, according to randomized controlled trials [39,40]. PET images provide incremental prognostic information to the clinical and angiographic findings with regard to event-free survival. An increased extent and severity of perfusion defects with stress PET were reported to be associated with increased frequency of adverse cardiac events, thus, this indicates PET can be used to predict cardiac mortality [41,42].

Cardiac PET is not yet as widely available as SPECT imaging. Furthermore, experience in image interpretation and operation may vary widely. Cardiac PET will continue to play a key role in the investigation of myocardial viability and perfusion contributing more to available data.

Integrated SPECT/PET-multislice CT has huge potential for cardiac imaging. The incremental value of hybrid imaging lies in accurate spatial co-localization of myocardial perfusion defects and anatomic coronary arteries. This combined technology allows detection and quantification of the burden of calcified and non-calcified plaques, quantification of vascular activity and endothelial health, identification of flow-limiting coronary stenosis, and potentially identification of high-risk plaques in the coronary artery tree [43]. Combined SPECT/CT and PET/CT systems are today well established in clinical routine imaging with promising results reported [44-48], although more multicentre trials are needed to validate the diagnostic value of the hybrid imaging modalities. Combined PET/MRI represents another new integrated protocol, however, it is only limited to a few clinical centers for preclinical cardiac imaging with a focus on animal experiments [49,50].

In summary, this editorial briefly reviews the diagnostic applications of these less-invasive imaging modalities including coronary CT angiography, cardiac MRI, cardiac SPECT and cardiac PET in coronary artery disease. Advantages and limitations of each imaging modality in the detection of coronary artery disease are also highlighted. Researchers are encouraged to contribute both original and review papers to this special issue with the aim of delivering both educational and teaching message to clinicians with research interests in cardiac imaging.

References

1. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, et al. (2010) Executive summary: heart disease and stroke statistics--2010 update: a report from the American Heart Association. *Circulation* 121: 948-954.
2. Gaziano TA, Bitton A, Anand S, Abrahams-Gessel S, Murphy A (2010) Growing epidemic of coronary heart disease in low- and middle-income countries. *Curr Probl Cardiol* 35: 72-115.
3. Sun ZH, Cao Y, Li HF (2011) Multislice computed tomography angiography in the diagnosis of coronary artery disease. *J Geriatr Cardiol* 8: 104-113.
4. Sun Z, Jiang W (2006) Diagnostic value of multislice computed tomography angiography in coronary artery disease: a meta-analysis. *Eur J Radiol* 60: 279-286.
5. Nieman K, Oudkerk M, Rensing BJ, van Ooijen P, Munne A, et al. (2001) Coronary angiography with multi-slice computed tomography. *Lancet* 357: 599-603.
6. Achenbach S (2006) Computed tomography coronary angiography. *J Am Coll Cardiol* 48: 1919-1928.
7. Pugliese F, Mollet NR, Runza G, van Mieghem C, Meijboom WB, et al. (2006) Diagnostic accuracy of non-invasive 64-slice CT coronary angiography in patients with stable angina pectoris. *Eur Radiol* 16: 575-582.
8. Sun Z, Lin C, Davidson R, Dong C, Liao Y (2008) Diagnostic value of 64-slice CT angiography in coronary artery disease: a systematic review. *Eur J Radiol* 67: 78-84.
9. Vanhoenacker PK, Heijnenbroek-Kal MH, Van Heste R, Decramer I, Van Hoe LR, et al. (2007) Diagnostic performance of multidetector CT angiography for assessment of coronary artery disease: meta-analysis. *Radiology* 244: 419-428.
10. Abdulla J, Abildstrom Z, Gotzsche O, Christensen E, Kober L, et al. (2007) 64-multislice detector computed tomography coronary angiography as potential alternative to conventional coronary angiography: a systematic review and meta-analysis. *Eur Heart J* 28: 3042-3050.

11. Sun Z, Choo GH, Ng KH (2012) Coronary CT angiography: current status and continuing challenges. *Br J Radiol* 85: 495-510.
12. Sun Z, Aziz YF, Ng KH (2012) Coronary CT angiography: how should physicians use it wisely and when do physicians request it appropriately? *Eur J Radiol* 81: e684-687.
13. Sun Z, Dimpudus FJ, Nugroho J, Adipranoto JD (2010) CT virtual intravascular endoscopy assessment of coronary artery plaques: a preliminary study. *Eur J Radiol* 75: e112-119.
14. Sun Z (2012) Cardiac CT imaging in coronary artery disease: Current status and future directions. *Quant Imaging Med Surg* 2: 98-105.
15. Liu YC, Sun Z, Tsay PK, Chan T, Hsieh IC, et al. (2013) Significance of coronary calcification for prediction of coronary artery disease and cardiac events based on 64-slice coronary computed tomography angiography. *Biomed Res Int* 2013: 472347.
16. Min JK, Feignoux J, Treutenaere J, Laperche T, Sablayrolles J (2010) The prognostic value of multidetector coronary CT angiography for the prediction of major adverse cardiovascular events: a multicenter observational cohort study. *Int J Cardiovasc Imaging* 26: 721-728.
17. Schlett CL, Banerji D, Siegel E, Bamberg F, Lehman SJ, et al. (2011) Prognostic value of CT angiography for major adverse cardiac events in patients with acute chest pain from the emergency department: 2-year outcomes of the ROMICAT trial. *JACC Cardiovasc Imaging* 4: 481-491.
18. Hou ZH, Lu B, Gao Y, Jiang SL, Wang Y, et al. (2012) Prognostic value of coronary CT angiography and calcium score for major adverse cardiac events in outpatients. *JACC Cardiovasc Imaging* 5: 990-999.
19. Hadamitzky M, Achenback S, Al-Mallah M, Berman D, Budoff M, et al. (2013) Optimized Prognostic Score for Coronary Computed Tomographic Angiography: Results From the CONFIRM Registry (COronary CT Angiography Evaluation For Clinical Outcomes: An International Multicenter Registry). *J Am Coll Cardiol* 62: 468-476.
20. Sun Z (2010) Multislice CT angiography in coronary artery disease: Technical developments, radiation dose and diagnostic value. *World J Cardiol* 2: 333-343.
21. Sun Z, Ng KH (2011) Coronary computed tomography angiography in coronary artery disease. *World J Cardiol* 3: 303-310.
22. Danias PG, Roussakis A, Ioannidis JP (2004) Diagnostic performance of coronary magnetic resonance angiography as compared against conventional X-ray angiography: a meta-analysis. *J Am Coll Cardiol* 44: 1867-1876.
23. Sommer T, Hackenbroch M, Hofer U, Schmiedel A, Willinek WA, et al. (2005) Coronary MR angiography at 3.0 T versus that at 1.5 T: initial results in patients suspected of having coronary artery disease. *Radiology* 234: 718-725.
24. Yang Q, Li K, Liu X, Bi X, Liu Z, et al. (2009) Contrast-enhanced whole-heart coronary magnetic resonance angiography at 3.0-T: a comparative study with X-ray angiography in a single center. *J Am Coll Cardiol* 54: 69-76.
25. Greenwood JP, Maredia N, Younger JF, Brown JM, Nixon J, et al. (2012) Cardiovascular magnetic resonance and single-photon emission computed tomography for diagnosis of coronary heart disease (CE-MARC): a prospective trial. *Lancet* 379: 453-460.
26. Nandalur KR, Dwamena BA, Choudhri AF, Nandalur MR, Carlos RC (2007) Diagnostic performance of stress cardiac magnetic resonance imaging in the detection of coronary artery disease: a meta-analysis. *J Am Coll Cardiol* 50: 1343-1353.
27. Jaarsma C, Leiner T, Bekkers SC, Crijs HJ, Wildberger JE, et al. (2012) Diagnostic performance of noninvasive myocardial perfusion imaging using single-photon emission computed tomography, cardiac magnetic resonance, and positron emission tomography imaging for the detection of obstructive coronary artery disease: a meta-analysis. *J Am Coll Cardiol* 59: 1719-1728.
28. de Jong MC, Genders TS, van Geuns RJ, Moelker A, Hunink MG (2012) Diagnostic performance of stress myocardial perfusion imaging for coronary artery disease: a systematic review and meta-analysis. *Eur Radiol* 22: 1881-1895.
29. Desai RR, Jha S (2013) Diagnostic performance of cardiac stress perfusion MRI in the detection of coronary artery disease using fractional flow reserve as the reference standard: a meta-analysis. *AJR Am J Roentgenol* 201: W245-252.
30. Fallahi B, Beiki D, Gholamrezanezhad A, Mahmoudian B, Ansari Gilani K, et al. (2008) Single Tc99m Sestamibi injection, double acquisition gated SPECT after stress and during low-dose dobutamine infusion: a new suggested protocol for evaluation of myocardial perfusion. *Int J Cardiovasc Imaging* 24: 825-835.
31. Elhendy A, van Domburg RT, Sozzi FB, Poldermans D, Bax JJ, et al. (2001) Impact of hypertension on the accuracy of exercise stress myocardial perfusion imaging for the diagnosis of coronary artery disease. *Heart* 85: 655-661.
32. Min JK, Kang N, Shaw LJ, Devereux RB, Robinson M, et al. (2008) Costs and clinical outcomes after coronary multidetector CT angiography in patients without known coronary artery disease: comparison to myocardial perfusion SPECT. *Radiology* 249: 62-70.
33. Bateman TM, O'Keefe JH Jr, Dong VM, Barnhart C, Ligon RW (1995) Coronary angiographic rates after stress single-photon emission computed tomographic scintigraphy. *J Nucl Cardiol* 2: 217-223.
34. Hachamovitch R, Hayes SW, Friedman JD, Cohen I, Berman DS (2005) A prognostic score for prediction of cardiac mortality risk after adenosine stress myocardial perfusion scintigraphy. *J Am Coll Cardiol* 45: 722-729.
35. Borges-Neto S, Shaw LK, Tuttle RH, Alexander JH, Smith WT 4th, et al. (2005) Incremental prognostic power of single-photon emission computed tomographic myocardial perfusion imaging in patients with known or suspected coronary artery disease. *Am J Cardiol* 95: 182-188.
36. Leslie WD, Tully SA, Yogendran MS, Ward LM, Nour KA, et al. (2005) Prognostic value of automated quantification of 99mTc-sestamibi myocardial perfusion imaging. *J Nucl Med* 46: 204-211.
37. Berman DS, Hachamovitch R, Shaw LJ, Friedman JD, Hayes SW, et al. (2006) Roles of nuclear cardiology, cardiac computed tomography, and cardiac magnetic resonance: assessment of patients with suspected coronary artery disease. *J Nucl Med* 47: 74-82.
38. Machac J (2005) Cardiac positron emission tomography imaging. *Semin Nucl Med* 35: 17-36.
39. Beanlands RS, Nichol G, Huszti E, Humen D, Racine N, et al. (2007) F-18-fluorodeoxyglucose positron emission tomography imaging-assisted management of patients with severe left ventricular dysfunction and suspected coronary disease: a randomized controlled trial (PARR-2). *J Am Coll Cardiol* 50: 2002-2012.
40. Abraham A, Nichol G, Williams KA, Guo A, deKemp RA, et al. (2010) 18F-FDG PET imaging of myocardial viability in an experienced center with access to 18F-FDG and integration with clinical management teams: the Ottawa-FIVE substudy of the PARR 2 trial. *J Nucl Med* 51: 567-574.
41. Yoshinaga K, Chow BJ, Williams K, Chen L, deKemp RA, et al. (2006) What is the prognostic value of myocardial perfusion imaging using rubidium-82 positron emission tomography? *J Am Coll Cardiol* 48: 1029-1039.
42. Dorbala S, Hachamovitch R, Curillova Z, Thomas D, Vangala D, et al. (2009) Incremental prognostic value of gated Rb-82 positron emission tomography myocardial perfusion imaging over clinical variables and rest LVEF. *JACC Cardiovasc Imaging* 2: 846-854.
43. Di Carli MF, Murthy VL (2011) Cardiac PET/CT for the evaluation of known or suspected coronary artery disease. *Radiographics* 31: 1239-1254.
44. Namdar M, Hany TF, Koepfli P, Siegrist PT, Burger C, et al. (2005) Integrated PET/CT for the assessment of coronary artery disease: a feasibility study. *J Nucl Med* 46: 930-935.
45. Rispler S, Keidar Z, Ghersin E, Roguin A, Soli A, et al. (2007) Integrated single-photon emission computed tomography and computed tomography coronary angiography for the assessment of hemodynamically significant coronary artery lesions. *J Am Coll Cardiol* 49: 1059-1067.
46. Groves AM, Speechly-Dick ME, Kayani I, Pugliese F, Endozo R, et al. (2009) First experience of combined cardiac PET/64-detector CT angiography with invasive angiographic validation. *Eur J Nucl Med Mol Imaging* 36: 2027-2033.
47. Sato A, Nozato T, Hikita H, Miyazaki S, Takahashi Y, et al. (2010) Incremental value of combining 64-slice computed tomography angiography with stress nuclear myocardial perfusion imaging to improve noninvasive detection of coronary artery disease. *J Nucl Cardiol* 17: 19-26.
48. Al Moudi M, Sun Z, Lenzo N (2011) Diagnostic value of SPECT, PET and PET/CT in the diagnosis of coronary artery disease: A systematic review. *Biomed Imaging Interv* 7: e9.

49. Catana C, Procissi D, Wu Y, Judenhofer MS, Qi J, et al. (2008) Simultaneous in vivo positron emission tomography and magnetic resonance imaging. Proc Natl Acad Sci U S A 105: 3705-3710.
50. Büscher K, Judenhofer MS, Kuhlmann MT, Hermann S, Wehrl HF, et al. (2010) Isochronous assessment of cardiac metabolism and function in mice using hybrid PET/MRI. J Nucl Med 51: 1277-1284.

Citation: Sun Z (2013) Cardiac Imaging Modalities in the Diagnosis of Coronary Artery Disease. J Clin Exp Cardiol S6: e001. doi:[10.4172/2155-9880.S6-e001](https://doi.org/10.4172/2155-9880.S6-e001)

This article was originally published in a special issue, [Coronary Heart Disease](#) handled by Editor(s). Dr. José G. Díez, Texas Heart Institute, USA