


9th International Conference on

GENOMICS & PHARMACOGENOMICS

June 15-16, 2017 London, UK



Henry M Sobell

University of Rochester, USA

The centers of premelton signals, the beginning and ends of genes

Premeltons are examples of emergent structures (i.e., structural solitons) that arise spontaneously in DNA due to the presence of nonlinear excitations in its structure. They are of two kinds: B-B (or A-A). Premeltons form at specific DNA-regions to nucleate site-specific DNA melting. These are stationary and, being globally non-topological, undergo breather motions that allow drugs and dyes to intercalate into DNA. B-A (or A-B) premeltons, on the other hand, are mobile, and being globally topological, act as phase-boundaries transforming B- into A- DNA during the structural phase-transition. They are not expected to undergo breather-motions. A key feature of both types of premeltons is the presence of an intermediate structural-form in their central regions (proposed as being a transition-state intermediate in DNA-melting and in the B- to A- transition), which differs from either A- or B- DNA. The so called beta-DNA, this is both metastable and hyperflexible—and contains an alternating sugar-puckering pattern along the polymer-backbone combined with the partial-un-stacking (in its lower energy-forms) of every other base-pair. Beta-DNA is connected to either B- or to A- DNA on either side by boundaries possessing a gradation of nonlinear structural-change, these being called the kink and the anti-kink regions. The presence of premeltons in DNA leads to a unifying theory to understand much of DNA physical-chemistry and molecular-biology. In particular, premeltons are predicted to define the 5' and 3' ends of genes in naked-DNA and DNA in active-chromatin, this is having important implications for understanding physical aspects of the initiation, elongation and termination of RNA-synthesis during transcription. For these and other reasons, the model will be of broader interest to the general audience working in these areas. The model explains a wide variety of data, and carries within it a number of experimental predictions—all readily testable – as will be described in my talk.

Biography

Henry M Sobell completed his studies at Brooklyn Technical High School (1948-1952), Columbia College (1952-1956), and University of Virginia School of Medicine (1956-1960). Instead of practicing clinical medicine, he then went to the Massachusetts Institute of Technology (MIT) to join Professor Alexander Rich in the Department of Biology (1960-1965), where, he had a Helen Hay Whitney Post-doctoral Fellowship where he learned the technique of single crystal X-ray analysis. He then joined the Chemistry department at University of Rochester.

sobell@localnet.co

Notes: