

2nd World Biotechnology Congress

December 04-05, 2017 | Sao Paulo, Brazil

***Bauhinia* stem extracts as a possible new treatment for breast cancer and metastasis: Inhibition of migration, invasion and of the activity of matrix metalloproteinases**

Santos K M¹, Gomes I N F¹, Romão W¹, Ribeiro R I M A¹, Silva-Oliveira R J², Pinto F E³, Oliveira B G³ and Reis R M V⁴

¹Federal University of São João del Rei, Brazil

²Barretos Cancer Hospital, Brazil

³Federal University of Espírito Santo, Brazil

⁴University of Minho, Portugal

Metastasis is the main cause of cancer-related death and requires the development of effective treatments with reduced toxicity and effective activity. The breast cancer is the most common among women and the second most prevalent type in the entire population. Thus the search for new sources of antitumor and anti-metastatic therapies, such as plants, is very important. In this work, we showed the antitumor and antimetastatic activities of four fractions (ID7, IID10, IA19 and IIIA32) of the stems of *Bauhinia* species on the murine breast cancer line 4T1. These fractions were used because they completely inhibited the MMP-2 and MMP-9 activity. The viability assay (MTT, Trypan blue) showed that all fractions studied decreased the viability of 4T1 cells, being the ID7 fraction the most selective. The fluorescence microscopy assay with acridine orange and propidium iodide showed that fractions increased the apoptotic cells percent. The wound closure and trans well assays were used to evaluate the migration cell, and the trans well assay with Matrigel was used to evaluate the invasion cell, wherein all fractions inhibited the 4T1 wound closure, IID10, IA19 fractions inhibited trans well migration, and ID7, IID10, and IIIA32 decreased invasion cell. Furthermore, all fractions increased the 4T1 adhesion to basement membrane components and decreasing the MMP-2 activity in the 4T1 cells supernatant. In the in vivo assay, this fraction decreased the volume and weight of the tumor extracted from mice induced with 4T1 and treated with ID7, in addition to decreasing the number of lung metastases. The ID7 ESI-MS(-) characterization suggesting the presence of fatty acids, phenolic acids, and diterpene. Thus, it was found that the *Bauhinia* fractions tested, exhibited selective antitumor and antimetastatic activity of breast cancer.

Biography

Santos K M is a Biomedical from José do Rosário Vellano University (2010). She has completed her Master's degree in Biotechnology from the Federal University of São João del Rei (2013). Currently, she is a PhD student in Biochemistry and Molecular Biology by Multicentral Program of the Brazilian Society of Biochemistry and Molecular Biology, working on the main themes: medicinal plants (*Bauhinia*) fractionation and identification of compounds, cancer and metastasis (metalloproteinases), Alzheimer (acetylcholinesterase inhibition), cryopreservation and culture of tumor cells and depression (behavioral assays). She is also Professor at the Biomedicine and Physiotherapy courses at the José do Rosário Vellano University and at the University Center of Formiga.

kamillamonteiro07@gmail.com

Notes: