6th International Conference on

EPIDEMIOLOGY & PUBLIC HEALTH

October 23-25, 2017 | Paris, France

EMERGING DRUG RESISTANCE SALMONELLA STRAINS FROM CHICKEN MEAT

Nahed Ghoneim^a and MonaKadry^a [°]Cairo University, Egypt

This work was carried out to study the occurrence of salmonella infections in chicken and human suffering from gastroenteritis in Egypt. A total of 349 chicken meat and 40 stool specimens of children were collected, samples were subjected to bacteriological examination and identified biochemically as salmonella then subjected to serological identification, also were tested for their antibiotic susceptibility by disc diffusion method Also, genotyping by PCR to detect *salmonella enterotoxine gene (stn)* and their expression by infant mouse assay. the study revealed that 14 (4.01%) and 2 (5%) were positive for *Salmonella species* in chicken meat and children respectively and serological identification were (*Salmonella Infantis, Salmonella Typhi, Salmonella Kentucky, Salmonella Rubislaw, Salmonella Poona, Salmonella Enteritidis*) in chicken meat and children isolates respectively. Disc diffusion method showed that 3 (21.4%) in chicken meat isolates and 2 (100%) in children isolates were multidrug resistant in which S. Kentucky have resistance to ciprofloxacin, the drug of choice for treating salmonellosis in children. Also, genotyping showed that 9 (64.28%) and 2 (100%) isolates confirmed to be enterotoxigenic strains in chicken meat and children respectively and this (stn) gene have been expressed (100%) by infant mouse assay. Sequencing and phylogenetic tree of 4 studied isolates resulted in assessment the relations between different isolation sources. Special attention must be paid to antibiotics that are used exclusively in poultry farms, appropriate measures must be taken to control the spread of resistant bacteria to human.

Notes: