

14th World Congress on

TOXICOLOGY AND PHARMACOLOGY

March 12-14, 2018 Singapore

The antioxidant effect of resveratrol in cisplatin induced oxidative stress

Florin Muselin¹, Eugenia Dumitrescu¹, Zeno Garban², Alexandru O Doma¹, Alexandra Trif¹ and Romeo T Cristina¹¹Banat University of Agricultural Sciences and Veterinary Medicine, Romania²Romanian Academy, Romania

Statement of the Problem: Cisplatin is one of the most used cytostatic with a broad antitumor spectrum. The uses of different cytostatic are followed by release of reactive oxygen species (ROS) responsible for the adverse effect of the chemotherapy. The purpose of this study was to determine if the resveratrol administration can reduce the cisplatin induced oxidative stress.

Methodology: The study was conducted on 24 Wistar rats divided in four groups: C - the control group receiving 1 ml of physiological saline I.P., CR - control blank group receiving 20 mg/kg resveratrol I.P., CP - receiving cisplatin 10 mg/kg I.P. and CP+R receiving combination of 10 mg/kg cisplatin and 20 mg/kg resveratrol I.P. At the end of experiment were analyzed the biomarkers of oxidative stress enzymes: Glutathione (GSH), glutathione reductase (GSH-r), glutathione peroxidase (GSH-px), catalase (CAT), superoxide dismutase (SOD) and malondialdehyde (MDA).

Findings: Administration of cisplatin was followed by significant decrease of GSH (-29.44%, $p < 0.01$), GSH-r (-31.88%, $p < 0.0001$), CAT (-52.28%, $p < 0.0001$) and increase of GSH-Px (+31.25%, $p < 0.001$), SOD (+11.27%, $p > 0.05$) and (+24.05%, $p < 0.05$) comparative to control group. In case of cisplatin combined with resveratrol administration, the majority of biomarker enzymes of oxidative stress presented not significant ($p > 0.05$) differences comparative to control group (GSH: -4.04%, GSH-r: -1.96%, GSH-px: +1.97%, SOD: -1.92%, MDA: +10.31%), exception in case of CAT which remains significantly lower than control (-21.62%, $p < 0.01$).

Conclusion: Analyzing the dynamic of enzymes biomarkers of oxidative stress we can say that administration of resveratrol can reduce the ROS formation and has a good effect as antioxidant in case of cisplatin administration.

References

1. Kursvietiene L, Staneviciene I, Mongirdiene A, Bernatoniene J (2016) Multiplicity of effects and health benefits of resveratrol. *Medicina*; 52(3): 148-155.
2. Kuhad A, Pilkhwal S, Sharma S, Tirkey N, Chopra K (2007) Effect of curcumin on inflammation and oxidative stress in cisplatin-induced experimental nephrotoxicity. *J Agric Food Chem*; 55(25): 10150-5.
3. Rjiba-Touati K, Boussema IA, Belarbia A, Achour A, Bacha H (2011) Protective effect of recombinant human erythropoietin against cisplatin-induced oxidative stress and nephrotoxicity in rat kidney. *Int J Toxicol*; 30(5): 510-517.
4. Tuan BT, Visacri MB, Amaral LS, Baldini D, Ferrari GB, Quintanilha JCF, Pincinato EC, Mazzola PG, Lima CSP, Moriel P (2016) Effects of high-dose Cisplatin chemotherapy and conventional radiotherapy on urinary oxidative and nitrosative stress biomarkers in patients with head and neck cancer. *Basic and Clin Toxicol and Pharmacol*; 118(1): 83-86.
5. Valdecantos MP, Pérez-Matute P, Quintero P, Martínez JA (2010) Vitamin C, resveratrol and lipoic acid actions on isolated rat liver mitochondria: all antioxidants but different. *Redox Rep*; 15(5): 207-216.

Biography

Florin Muselin has his expertise in heavy metals reproductive toxicology, oxidative stress and medicinal and poisonous plants. He is an Associate Professor at Faculty of Veterinary Medicine from BUASMV Timisoara, Romania.

florin.muselin@gmail.com




Figure-1: Schematic reaction of the cisplatin administration, production of reactive oxygen species and resveratrol.