

## De Novo Urgency: A Review of the Literature

Lleberia J\*, Pubill J, Mestre M, Aguiló O, Serra L and Canet Y

Department of Obstetrics and Gynecology, Corporation Park Tauli Sabadell, Spain

### Summary

De novo urgency can be defined as the appearance of urge urinary incontinence after an anti-incontinence surgery that persists after six months. This term can only be used when the patient did not have any urgency preoperative symptoms. The incidence of de novo urgency varies between 3.1% and 25.9% and although its pathophysiology is unclear, it is suggested that an obstruction (absolute or relative) is caused during the procedure that induces a reaction in the detrusor muscle. The evaluation of de novo urgency includes a thorough anamnesis, physical examination, residual urine evaluation, urinalysis, and urodynamic testing. When there is evidence of a bladder outlet obstruction, its treatment is urethrolisis or urethral dilatation. For patients without obstruction, a more conservative approach including pelvic floor exercises and antimuscarinics is recommended. A meticulous surgical technique is the key to try preventing this pathology. As excellent practise for incontinence techniques cannot prevent all cases must continue deepen their study of anatomical and functional factors.

**Keywords:** De novo urgency; Tape complications; Physiopatology

### Introduction

Pelvic floor muscle training is the first therapy option for stress incontinence with as close as to 50% efficacy [1]. In case of failure or when this therapy is not possible due to concomitant presence of Prolapse or limitations of the patient (physical or mental) the next step is, nowadays, anti-incontinence surgery such as tension-free vaginal tape [2-4]. Since the introduction of this technique in 1995, it has won popularity and is at present one of the most used methods for the treatment of stress incontinence in developed countries [5]. The reported success rates for this procedure are high, results reported at seventeen year's follow up indicate a 90% objective cure, however we will focus on one of its mid-long term complications: de novo urgency with or without incontinence [6-8].

Post-surgical follow up have reported that the incidence of the novo urgency varies between 3.1% and 25.9% [9-13]. Despite a lack of a generally agreed definition, according to the series it is clear that a prerequisite is that the patient had no urgency symptoms preoperatively. The term of the novo urgency implies the postoperative development of urge urinary incontinence (defined as the involuntary leakage from the urethra synchronous with the sensation of a sudden, compelling desire to void that is difficult to defer) and its persistence after six months [14,15].

### Pathophysiology

The etiology and mechanisms for the development of de novo urgency are unclear. However it is suggested that, as it is related to a surgical urethral sphincter support, a bladder outlet obstruction (absolute or relative) causes changes in the voiding function which are responsible of the onset of the symptoms [16]. Generally, it has been shown that women with bladder outlet obstruction present a higher grade of detrusor over activity than women without it. Furthermore, both Pope and Cardozo research studies have demonstrated that anti-incontinence surgery results in an increased urethral resistance during micturition [17-20]. Because detrusor pressure generated during voiding is largely dependent on outflow resistance, one would expect that an increase in urethral resistance should be met by an increase in detrusor pressure. By 12 months post surgery, mean void pressure in these women increased from  $26 \pm 8$  cm H<sub>2</sub>O to  $57 \pm 11$  cm H<sub>2</sub>O, average flow rates diminished and urethral resistance increased [17]. From these results it was concluded that the establishment of some degree of urethral obstruction (with or without overactive bladder

symptoms) is inevitable in the process of restoring continence [19]. Moreover, there was no change in urethral resistance in those patients whose surgery failed [18]. Bombier et al. performed a postoperative bladder neck position and urethral compression evaluation using magnetic resonance that demonstrated bladder neck elevation and urethral compression to be associated with detrusor over activity [16]. Bump et al. described the key to successful anti-incontinence surgery as the establishment of optimal dynamic obstruction [20].

Although bladder outlet obstruction has commonly been used as one explanation for the novo urgency; in some patients no evidence obstruction has been detected. Therefore, it is logical to postulate alternative mechanisms for the development of this symptomatology. Falconer et al. performed a two year follow-up of Tension-free Vaginal Tape (TVT) and observed an increase of paraurethral connective tissue metabolism which was more pronounced in postmenopausal women [21,22].

Other authors suggest that damage on the autonomic innervation may be the onset of the syndrome. In vitro studies of detrusor smooth muscle from patients and animal models with detrusor over activity have revealed changes in physiologic properties consistent with denervation and supersensitivity [23]. Moreover, it has been found that postoperative dysfunctional voiding may occur due to altered innervations patterns to the pelvic floor [24]. Additionally, the incidence of postoperative the novo urgency is similar in current sling procedures and in classical abdominal operations [16,25,26]. Indeed, it has been suggested that detrusor over activity may have been caused by the damage inflicted on the autonomic innervation of the bladder by surgical dissection [9]. Therefore, a history of caesarean section could be considered as a risk factor for developing de novo urgency and undergoing a hysterectomy could worsen its symptoms [27].

Nevertheless, the majority of anti-incontinence procedures such as

**\*Corresponding author:** Judith Lleberia Juanos, Department of Obstetrics and Gynecology, Corporation Park Tauli Sabadell, Park Tauli 1, Sabadell, Barcelona 08191, Spain, Tel: 0034-937231010; E-mail: [lleberia@hotmail.com](mailto:lleberia@hotmail.com)

**Received** July 31, 2013; **Accepted** August 26, 2013; **Published** August 28, 2013

**Citation:** Lleberia J, Pubill J, Mestre M, Aguiló O, Serra L, et al. (2013) De Novo Urgency: A Review of the Literature. Gynecol Obstet 3: 166. doi:10.4172/2161-0932.1000166

**Copyright:** © 2013 Lleberia J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

TVT, are not at all related to dissection, most patients do not show evidence of obstruction nor have they undergone any gynaecological surgery, hence the logical hypothesis that there are important unknown etiological factors for de novo urgency and its multifactorial etiology [8].

## Symptoms

De novo urgency is the onset of urgency with or without leakage of urine after surgery that persists after six months. Irritative symptoms after surgery may be due to bladder adaptation to new urethral pressures and should not be diagnosed as de novo urgency. It is the symptoms persistence what defines this condition and even more important is how it diminishes the patient quality of life on the long-term.

One can understand that even after successful results curing stress incontinence a patient will be disappointment when one incontinence type transforms into the other. In general, the immediate and irresistible drive to micturate, nocturnal enuresis and pain make urgency a more bothersome condition than stress incontinence. The anamnesis should focus on these symptoms as well as symptoms that could suggest urine retention. The use of a disease-specific questionnaire such as ICIQ-SF, as well as a 3-day voiding diary will be helpful. Also, it is essential to know accurately the type of anti-incontinence procedure that the patient underwent [9].

## Physical Examination

A directed urogynecological and neurologic examination needs to be performed. Anterior vaginal wall is carefully examined with the help of a vaginal speculum paying close attention to the appearance of urethra and bladder neck. Scarification is also examined as it can give us information on an over tensioned tape whether because it is folded or too-tightly positioned. If that is unclear, urethra evaluation with a hyssop and Qtipstest can be useful. The presence of prolapse is also examined as it is suggested that there is an association between prolapse and urgency [28,29]. Then both vaginal and rectal digital exam must be performed and bulbocavernous reflex and perineal sensation assessed. Also, given the importance of differential diagnosis between urgency and urine retention any signs of said condition should be detected and an evaluation of post void residue performed either by catheterism or by perineal ultrasound.

## Clinical Exams

### Sediment - urine culture

The initial evaluation of any patient presenting urgency should include urine analysis to rule out occult infection or inflammatory condition. Thus, it would be appropriate to perform a dipstick urinalysis with microscopy seeking for leucocytes and bacteria and also obtain a urine culture.

### Ultrasound

There are three ultrasonographic parameters that can be helpful in the differential diagnosis of de novo urgency: residual urine assessment, sling location and bladder wall thickness.

Is possible to determine if urgency symptoms are caused by chronic urine retention both performing a catheterism as well as an abdominal or vaginal ultrasound. Haylen et al. reported that the bladder volume can be calculated by the formula  $\text{Volume} = 5.9 \times (\text{height} \times \text{depth}) - 14.6$  ml (95% confidence limits =  $\pm 37$  ml) [30]. Although there are other methods of measuring urine volume, this formula is currently the most used [31,32].

Ultrasonographic detection of the sling generally allows the assessment of its location. Indeed, several studies have reported data about specific distances to determine sling position [33-35]. It is logical that this information could be useful as a means of understanding the mechanism of postoperative bladder dysfunction. For instance, tapes can be located in the bladder or urethral lumen, or even when well located they can be too tightly pulled. Dietz et al. suggest that although position and mobility of the TVT vary markedly, variations in placement seem to have relatively little effect on symptoms. They only found association in case of over tensioned sling during valsalva and overactive bladder, urgency, increased frequency and bladder voiding dysfunctions [35,36].

The third sonographic parameter for the evaluation of this condition is bladder wall thickness evaluation. According to Latthe's last review (2010) [37], bladder wall thickness greater than 5 mm is useful in the diagnosis of overactive detrusor with a sensitivity between 40% and 84% and specificity between 78-89%. More recent studies have found a correlation between bladder wall thickness and urgency both clinical and urodynamic [38]. This evaluation can be logically suitable in cases of de novo urgency. Other exams such as magnetic resonance offer detailed pelvic anatomy iconography however the visualization of the sling is more difficult with this technique [39,40].

## Urodynamics

Urodynamic testing is an essential component of any evaluation of voiding dysfunction, and especially after anti-incontinence surgery. Given that obstruction may occur in the presence of normal emptying, a low post void residual in an initial evaluation should not falsely reassure the physician of the absence of obstruction. In cases of a certain and indubitable history of normal voiding before surgery and an unequivocal temporal relationship between surgery and the appearance of post void residual, urethrolysis can be performed immediately without the need of urodynamic testing however, most cases of bladder voiding dysfunction require it. A filling cystometrogram is an essential part of the urodynamic evaluation because patients with de novo urgency suffer from sensory urgency or detrusor overactivity alone or in combination. To determine the presence of detrusor overactivity one should carefully look for unstable detrusor contractions of any amplitude during the course of filling and attempt to obtain valsalva induced instability. As for sensory urgency, it is characterized by a reduction in cystometric capacity without a compliance loss or instability. Also, a filling curve for low bladder compliance may be observed which could indicate an increase of the bladder tone due to occult motor instability without uninhibited contractions of significant amplitude. Al Ghazo et al. studied the correlation between urgency symptoms and urodynamics findings in overactive bladder and concluded that 61.3% of patients who suffered urgency without leaking and 69.8% of patients with urge urinary incontinence presented overactive bladder in urodynamic evaluation. In addition, they found an association between the combination of nocturia and incontinence with overactive bladder [41].

## Cystoscopy

Cystoscopy is performed to rule out the presence of a foreign body into the bladder and it also allows the inspection of the bladder looking for any inflammatory changes indicative of chronic cystitis or occult neoplasm.

## Treatment

Treatment for the novo urge syndrome depends on the overall

symptoms of the patient combined with the most likely etiologic diagnosis objectively determined. Therefore, in the first place it should be decided whether the patient benefits from conservative treatment or if there is a bladder outlet obstruction that requires urethrolysis.

### Medical treatment

If the evaluation shows absence of obstruction one should start with conservative approach consisting of behavioural therapy combined with pharmacotherapy. It is generally recommended to start with a timed voiding schedule programme combined with limitations of excessive fluids and potential bladder irritants or diuretics such as caffeine or alcohol intake. Antimuscarinic therapy or adrenergic receptor beta 3 agonist therapy should be prescribed when there is urodynamic evidence of overactive bladder even in the presence of mild urine retention without significant post void residual. In case of failure or intolerance of antimuscarinic therapy one should proceed with current alternative therapies for urge urinary incontinence such as sacral or tibial neuromodulation or also the injection of botulinic toxin. These patients will indeed need a close follow-up to detect the development of occult urinary retention. In general, patients with urinary retention will eventually fail conservative therapy.

### Urethrolysis

Urethrolysis is the elective procedure for patients with symptomatic de novo urgency and evidence of obstruction. In these patients one should proceed with the loosening of the sling or with urethral dilatation. For patients with mild symptoms and minimal obstruction, urethrolysis will be performed after failure of conservative treatment with cure rates ranging between 65% and 92% although between 19% and 50% of these patients may develop recurrent stress urinary incontinence [9].

### Prevention

A meticulous surgical technique is the key to preventing obstruction. Urethral obstruction may occur if the sling is located too close to the urethra, or if it folds or buckle. In case of doubt, sonographic evaluation may be helpful in the early assessment (even intraoperatively) of sling location. Alternatively, in cases where a prolonged postoperative catheterism is needed one should determine the location of the tape and proceed to its immediate loosening if it is pulled too tightly.

The mechanism of action of tension-free vaginal tape procedure can be explained by a biomechanical model. Given that the biological reaction of the tape unpredictable, the only factor that can be controlled is the tension of the tape.

### Conclusions

The increasing use of tension-free tapes for the treatment of incontinence entails an increase of de novo urgency. Understanding the pathophysiological changes that occur in the vesicourethral function due to this procedure is of the utmost importance to reach an accurate diagnosis and for the appropriate management of de novo symptoms.

### References

1. Bo K, Talseth T, Holme I (1999) Single blind, randomized controlled trial of pelvic floor exercises, electrical stimulation, vaginal cones and no treatment in management of genuine stress incontinence in women. *BMJ* 318: 487.
2. Ulmsten U, Henriksson L, Johnson P, Varhos G (1996) An ambulatory surgical procedure under local anesthesia for treatment of female urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 7: 81-85.
3. de Leval J (2003) Novel surgical technique for the treatment of female stress urinary incontinence: transobturator vaginal tape inside-out. *Eur Urol* 44: 724-730.

4. Zullo MA, Plotti F, Calcagno M, Marullo E, Palaia I, et al. (2001) One-year follow-up of tension-free vaginal tape (TVT) and trans-obturator suburethral tape from inside to outside (TVT-O) for surgical treatment of female stress urinary incontinence: a prospective randomized trial. *Eur Urol* 51: 1376-1382.
5. Abouassaly R, Steinberg JR, Lemieux M, Marois C, Gilchrist LI, et al. (2004) Complications of tension-free vaginal tape surgery: a multi-institutional review. *BJU Int* 94: 110-113.
6. Serati M, Bauer R, Cornu JN, Cattoni E, Braga A, et al. (2013) TVT-O for the treatment of pure urodynamic stress incontinence: efficacy, adverse effects, and prognostic factors at 5-year follow-up. *Eur Urol* 63: 872-878.
7. Nilsson CG, Palva K, Aarnio R, Morcos E, Falconer C (2013) Seventeen years' follow-up of the tension-free vaginal tape procedure for female stress urinary incontinence. *Int Urogynecol J* 24: 1265-1269.
8. Holmgren C, Nilsson S, Lanner L, Hellberg D (2007) Frequency of de novo urgency in 463 women who had undergone the tension-free vaginal tape (TVT) procedure for genuine stress urinary incontinence--a long-term follow-up. *Eur J Obstet Gynecol Reprod Biol* 132: 121-125.
9. Kershen RT, Appell RA (2002) De novo urge syndrome and detrusor instability after anti-incontinence surgery: current concepts, evaluation, and treatment. *Curr Urol Rep* 3: 345-353.
10. Collinet P, Ciofu C, Costa P, Cosson M, Deval B, et al. (2008) The safety of the inside-out transobturator approach for transvaginal tape (TVT-O) treatment in stress urinary incontinence: French registry data on 984 women. *Int Urogynecol J Pelvic Floor Dysfunct* 19: 711-715.
11. Celebi I, Güngördük K, Ark C, Akyol A (2009) Results of the tension-free vaginal tape procedure for treatment of female stress urinary incontinence: a 5-year follow-up study. *Arch Gynecol Obstet* 279: 463-467.
12. Lee KS, Choo MS, Doo CY, Han DH, Lee YS, et al. (2008) The long term (5-years) objective TVT success rate does not depend on predictive factors at multivariate analysis a multicentre retrospective study. *Eur Urol* 53: 176-183.
13. Giberti C, Gallo F, Cortese P, Schenone M (2007) Transobturator tape for treatment of female stress urinary incontinence: objective and subjective results after a mean follow-up of two years. *Urology* 69: 703-707.
14. Bates P, Bradley WE, Glen E, Griffiths D, Melchior H, et al. (1979) The standardization of terminology of lower urinary tract function. *J Urol* 121: 551-554.
15. Haylen BT, de Ridder D, Freeman RM, Swift SE, Berghmans B, et al. (2010) An International Urogynecological Association (IUGA)/International Continence Society (ICS) Joint Report on the Terminology for Female Pelvic Floor Dysfunction. *Neurourol Urodynam* 21: 5-26.
16. Bombier L, Freeman RM, Perkins EP, Williams MP, Shaw SR (2002) Why do women have voiding dysfunction and de novo detrusor instability after colposuspension? *BJOG* 109: 402-412.
17. Pope AJ, Shaw PJR, Coptcoat MJ, Worth PHL (1990) Changes in bladder function following a surgical alteration in outflow resistance. *Neurourol Urodyn* 9: 503-508.
18. Klutke JJ, Klutke CG, Bergman J, Elia G (1999) Bladder neck suspension for stress urinary incontinence: how does it work? *Neurourol Urodyn* 18: 623-627.
19. Cardozo LD, Stanton SL, Williams JE (1979) Detrusor instability following surgery for genuine stress incontinence. *Br J Urol* 51: 204-207.
20. Bump RC, Hurt WG, Elser DM, Theofrastous JP, Addison WA, et al. (1999) Understanding lower urinary tract function in women soon after bladder neck surgery. Continence Program for Women Research Group. *Neurourol Urodyn* 18: 629-637.
21. Falconer C, Ekman-Ordeberg G, Malmström A, Ulmsten U (1996) Clinical outcome and changes in connective tissue metabolism after intravaginal slingplasty in stress incontinent women. *Int Urogynecol J Pelvic Floor Dysfunct* 7: 133-137.
22. Falconer C, Ekman-Ordeberg G, Ulmsten U, Westergren-Thorsson G, Barchan K, et al. (1996) Changes in paraurethral connective tissue at menopause are counteracted by estrogen. *Maturitas* 24: 197-204.
23. Mills IW, Greenland JE, McMurray G, McCoy R, Ho KM, et al. (2000) Studies of the pathophysiology of idiopathic detrusor instability: the physiological properties of the detrusor smooth muscle and its pattern of innervation. *J Urol* 163: 646-651.

24. Artibani W (1997) Diagnosis and significance of idiopathic overactive bladder. *Urology* 50: 25-32.
25. Kinn AC (1995) Burch colposuspension for stress urinary incontinence. 5-year results in 153 women. *Scand J Urol Nephrol* 29: 449-455.
26. Langer R, Lipshitz Y, Halperin R, Pansky M, Bukovsky I, et al. (2001) Long-Term (10-15 years) follow-up after Burch colposuspension for urinary stress incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 12: 323-326.
27. Neumann G, Olesen PG, Hansen V, Lauszus FF, Ljungström B, et al. (2004) The short-term prevalence of de novo urinary symptoms after different modes of hysterectomy. *Int Urogynecol J Pelvic Floor Dysfunct* 15: 14-19.
28. Boer TA, Salvatore s, Cardozo L, Chapple C, Kelleher C, et al. (2010) Pelvic Organ Prolapse and Overactive Bladder. *Neurourol Urodyn* 29: 30-39.
29. Lleberia J, Pubill J, Mestre M, Garcia E, Grimau M, et al. (2011) Surgical treatment of mixed urinary incontinence: effect of anterior colpoplasty. *Int Urogynecol J* 22: 1025-1030.
30. Haylen BT, Frazer MI, Sutherst JR, West CR (1989) Transvaginal ultrasound in the assessment of bladder volumes in women. Preliminary report. *Br J Urol* 63: 149-151.
31. Beacock CJ, Roberts EE, Rees RW, Buck AC (1985) Ultrasound assessment of residual urine. A quantitative method. *Br J Urol* 57: 410-413.
32. Rageth JC, Langer K (1982) Ultrasonic assessment of residual urine volume. *Urol Res* 10: 57-60.
33. Chantarasorn V, Shek KL, Dietz HP (2011) Sonographic appearance of transobturator slings: implications for function and dysfunction. *Int Urogynecol J* 22: 493-498.
34. Long CY, Hsu CS, Lo TS, Liu CM, Chen YH, et al. (2008) Ultrasonographic assessment of tape location following tension-free vaginal tape and transobturator tape procedure. *Acta Obstet Gynecol Scand* 87: 116-121.
35. Dietz HP, Mouritsen L, Ellis G, Wilson PD (2004) How important is TVT location? *Acta Obstet Gynecol Scand* 83: 904-908.
36. Chene G, Cotte B, Tardieu AS, Savary D, Mansoor A (2008) Clinical and ultrasonographic correlations following three surgical anti-incontinence procedures (TOT, TVT and TVT-O). *Int Urogynecol J Pelvic Floor Dysfunct* 19: 1125-1131.
37. Latthe PM, Champaneria R, Khan KS (2010) Systematic review of the accuracy of ultrasound as the method of measuring bladder wall thickness in the diagnosis of detrusor overactivity. *Int Urogynecol J* 21: 1019-1024.
38. Kuhn A, Genoud S, Robinson D, Herrmann G, Güntherth A, et al. (2011) Sonographic transvaginal bladder wall thickness: does the measurement discriminate between urodynamic diagnoses? *Neurourol Urodyn* 30: 325-328.
39. Hubka P, Doumouchtsis SK, Berger MB, DeLancey JO (2012) Variation of distances from mid-urethra to the obturator foramen: an MRI study. *Int Urogynecol J* 23: 1075-1080.
40. Morris VC, Murray MP, Delancey JO, Ashton-Miller JA (2012) A comparison of the effect of age on levator ani and obturator internus muscle cross-sectional areas and volumes in nulliparous women. *Neurourol Urodyn* 31: 481-486.
41. Al-Ghazo MA, Ghalayini IF, Al-Azab R, Hani OB, Matani YS, et al. (2011) Urodynamic detrusor overactivity in patients with overactive bladder symptoms. *Int Neurourol J* 15: 48-54.