

Do We have Enough Direct Evidence to Postulate that Abnormally Shaped Red Blood Cells Impair Microvascular Blood Flow in Critical Conditions?

Sergey B Zaets*

1Wall Street, 5B, Fort Lee, NJ 07024, USA

Red blood cells (RBCs) play a key role in microcirculation and oxygen delivery to organ and tissues. This role is determined by their ability to deform and to pass through the vessels that are even smaller than RBC size. There is a close relationship between RBC deformability and its unique disc shape. Under different circumstances, RBC may change their shape and undergo echinocyte, stomatocyte or spherocyte transformation. Echinocytes or stomatocytes are potentially capable to restore the disc shape if they are put in favorable environment (e.g. fresh normal plasma), whereas the spherocyte transformation is considered to be irreversible [1]. *In-vitro* studies using micropore filtration technique demonstrated that normal RBCs (discocytes) have an optimum shape for the flow in microvessels. In contrast to discocytes, stomatocytes and echinocytes produced by adding chlorpromazine or sodium salicylate, respectively, to normal RBCs demonstrated altered passage through circulation [2]. Mechanisms of microcirculatory disorders in critical conditions such as trauma-hemorrhagic shock, sepsis or burn have been under investigation for decades. Numerous experimental and clinical studies have proven that these insults cause a significant decrease in RBC deformability [3-11]. Smaller number of publications report RBC shape changes as a result of critical conditions [5,11,12-15]. Experimental studies demonstrated that the number of abnormal (reversibly and irreversibly changed) RBCs after hemorrhagic shock was increased more than five times compared to control values and their proportion to normal discocytes exceeded 30% [5,13]. The similar phenomenon was observed in patients with severe trauma [14]. Linear regression analysis revealed a tight correlation between the percentage of abnormally shaped RBCs and their elongation index, the measure of deformability, in rats subjected to trauma-hemorrhagic shock [5]. The question comes if we have enough direct evidence to postulate that abnormally shaped RBCs impair microvascular blood flow in critical conditions?

Several studies demonstrated that RBCs artificially hardened by various chemicals can cause microcirculatory blockade. Simchon et al. performed exchange transfusions in rats with RBCs with reduced deformability or with sham RBCs [16]. Lowered RBC deformability was achieved by incubation with glutaraldehyde and confirmed by micropore filtration technique. Labeling of RBCs with isotopes made possible their detection in organs and tissues. Experiments showed that poorly deformed RBCs were trapped in spleen, lung, and liver. The degree of flow reduction in these organ compared to control animals (rats subjected to sham RBC transfusion) measured using radioactive microspheres was 77%, 69%, and 54%, respectively and correlated with the amount of trapped RBCs. However, the authors did not investigate the shape of transfused RBCs. Betticher et al. perfused isolated rabbit lungs with stiffened and abnormally shaped RBCs (stomatocytes) [17]. RBC transformation was achieved by adding chlorpromazine. Reduced deformability of RBCs and their shape abnormalities were confirmed by micropore filtration technique and light microscopy, respectively. The authors demonstrated a decrease in pulmonary oxygen diffusion capacity and an increase in the mean pulmonary arterial circulating pressure (both by 18%) as a result of perfusion with abnormal RBCs. It should be stated that in these two aforementioned experiments,

abnormal RBCs were produced *in-vitro* but not as a result of a real critical condition. The conclusions received in these two studies were confirmed in "pure" *in-vivo* experiments by Machiedo et al. [18]. The authors performed exchange transfusion in rats with blood obtained animals rats subjected to trauma-hemorrhagic shock or to sham shock. The reduced RBC deformability in rats that experienced trauma-hemorrhagic shock and normal RBC deformability in those who were subjected to sham shock was confirmed by laser ektactometry. However, the authors did not investigate RBC shape in transfused blood. It was demonstrated that microvascular blood flow in the lungs, spleen, and intestine measured by radioactive microsphere technique was significantly lower in rats subjected to the infusion of trauma-hemorrhagic shock blood than in those that were infused with sham blood.

Our current knowledge suggests that the hypothesis of the important role of critical insult-induced RBC shape changes in the development of microcirculatory disorders in organ and tissues is most likely true. However, there is only indirect evidence. Experimental studies that prove this hypothesis have certain limitations. In some of them, RBCs altered *in-vitro* but not RBCs that became abnormal as a result of a real critical condition modulated *in-vivo* were used to cause a decrease in microvascular blood flow. Other experiments, where blood from animals subjected to trauma-hemorrhagic shock was transfused to recipient animals to induce microcirculatory disorders, did not evaluate RBC morphology in infused blood. Thus additional experiments are still needed to prove direct contribution of RBC shape alterations caused by critical conditions to the development of post-insult microcirculatory disorders. If this hypothesis is confirmed, the next step should be determination of the proportion between reversibly and irreversibly changed RBCs that may cause clinically relevant microvascular blood flow abnormalities.

References

1. Bessis M (1972) Red cell shapes. An illustrated classification and its rationale. *Nouv Rev Fr Hematol* 12: 721-745.
2. Reinhart HR, Chien S (1986) Red cell rheology in stomatocyte-echinocyte transformation: roles of cell geometry and cell shape. *Blood* 67: 1110-1118.
3. Poraicu D, Mogoseanu A, Tomescu N, Bota C, Menessy I (1983) Decrease of red blood cell filterability seen in intensive care I. The correlation of low erythrocyte filterability with mortality and its return to normal values in critically ill patients under parenteral nutrition. *Resuscitation* 10: 291-303.

*Corresponding author: Sergey Zaets, 1Wall Street, 5B, Fort Lee, NJ 07024, USA, Tel: 1-201-873-8901; Fax: 1-609-419-3006; E-mail: zaets001@yahoo.com

Received August 02, 2013; Accepted August 05, 2013; Published August 08, 2013

Citation: Zaets SB (2013) Do We have Enough Direct Evidence to Postulate that Abnormally Shaped Red Blood Cells Impair Microvascular Blood Flow in Critical Conditions? *Anatom Physiol* 3: e122. doi: [10.4172/2161-0940.1000e122](https://doi.org/10.4172/2161-0940.1000e122)

Copyright: © 2013 Zaets SB. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

4. Langenfeld JE, Livingston DH, Machiedo GW (1991) Red cell deformability is an early indicator of infection. *Surgery* 110: 398-403.
5. Zaets SB, Berezina TL, Morgan C, Kamiyama M, Spolarics Z, et al. (2003) Effect of trauma-hemorrhagic shock on red blood cell deformability and shape. *Shock* 19: 268-273.
6. Tatarishvili J, Momtselidze N, McHedlishvili G (2004) Blood rheological abnormalities in the microcirculation during experimental traumatic shock. *Clin Hemorheol Microcirc* 30: 407-410.
7. Hurd TC, Dasmahapatra KS, Rush BF Jr, Machiedo GW (1988) Red blood cell deformability in human and experimental sepsis. *Arch Surg* 123: 217-220.
8. Baskurt OK, Gelmont D, Meiselman HJ (1998) Red blood cell deformability in sepsis. *Am J Respir Crit Care Med* 157: 421-427.
9. Reggiori G, Occhipinti G, De Gasperi A, Vincent JL, Piagnerelli M (2009) Early alterations of red blood cell rheology in critically ill patients. *Crit Care Med* 37: 3041-3046.
10. Bekyarova G, Yankova T (1998) Alpha-Tocopherol and reduced glutathione deficiency and decreased deformability of erythrocytes after thermal skin injury. *Acta Physiol Pharmacol Bulg* 23: 55-59.
11. Zaets SB, Berezina TL, Xu DZ, Lu Q, Ricci J, et al. (2003) Burn-induced red blood cell deformability and shape changes are modulated by sex hormones. *Am J Surg* 186: 540-546.
12. Poraicu D, Sandor S, Menessy I (1983) Decrease of red blood cell filterability seen in intensive care. II. Red blood cell crenelation "in vivo" as morphological evidence of increased red blood cell viscosity in low flow states. *Resuscitation* 10: 305-316.
13. Berezina TL, Zaets SB, Kozhura VL, Novoderzhkina IS, Kirsanova AK, et al. (2001) Morphologic changes of red blood cells during hemorrhagic shock replicate changes of aging. *Shock* 15: 467-470.
14. Berezina TL, Zaets SB, Machiedo GW (2004) Alterations of red blood cell shape in patients with severe trauma. *J Trauma* 57: 82-87.
15. Piagnerelli M, Boudjeltia KZ, Brohee D, Piro P, Carlier E, et al. (2003) Alterations of red blood cell shape and sialic acid membrane content in septic patients. *Crit Care Med* 31: 2156-2162.
16. Simchon S, Jan KM, Chien S (1987) Influence of reduced red cell deformability on regional blood flow. *Am J Physiol* 253: H898-H903.
17. Betticher DC, Reinhart WH, Geiser J (1995) Effect of RBC shape and deformability on pulmonary O₂ diffusing capacity and resistance to flow in rabbit lungs. *J Appl Physiol* 78: 778-783.
18. Machiedo GW, Zaets SB, Berezina TL, Xu DZ, Feketova E, et al. (2009) Trauma-hemorrhagic shock-induced red blood cell damage leads to decreased microcirculatory blood flow. *Crit Care Med* 37: 1000-1010.