

Effect of Cadmium Toxicity on Aquatic Macrophyte *Pistia Stratiotes* (L.)Rolli N.M^{1*}, Suvarna khandi SS¹, Gadi SB², Mulagund GS³ and Taranath TC³¹BLDEA's Degree College Jamkhandi, Karnataka, India²Dept of Botany JSS College Dharwad, Karnataka, India³Department of Botany Karnataka University Dharwad, India**Abstract**

The present study focuses on cadmium toxicity on morphology and selected biochemical parameters of *Pistia*. The laboratory experiments were conducted for the assessment of morphological index parameters (MIP), biochemical parameters and accumulation of cadmium in plants at its various concentrations (0.1, 0.5, 1.0, 1.5 and 2.0 ppm) at the regular interval for 12 days exposure. *Pistia* showed visible symptoms like withering of roots, chlorosis and necrosis at higher concentration (2.0 ppm), however, the plant showed normal growth at lower concentration (0.1 ppm). The estimation of biochemical parameters (Total chlorophyll, Protein and Carbohydrate) of test plants showed a significant increase at lower concentration (0.1 ppm) of cadmium. The biochemical parameters decrease with increase in exposure concentration and duration. The toxic effect of cadmium is directly proportional to its concentration and duration. The accumulation of cadmium by *Pistia stratiotes* was maximum at 4 days exposure duration and marginal at subsequent concentrations and exposure duration. With respect to biochemical parameters the concentrations are significant. However, metal accumulation is significant at concentrations and exposure duration.

Keywords: Accumulation; Biochemical parameters; Cadmium; Toxicity

Introduction

Heavy metal contamination in the water bodies is increasing at an alarming rate due to industrial and anthropogenic activities [1]. Heavy metals cannot be degraded and therefore, are persistent. Efforts are being made to develop technologies that use biological materials effectively in removing heavy metals from the environment. Plants possess the capacity to accumulate heavy metals. Higher plant species possess physiological traits that enable them to tolerance and hyper accumulate metals [2,3]. The metal tolerance of plants may be attributed to different enzymes, stress proteins and phytochelatin [4]. The accumulation of metals at higher concentrations causes retardation of growth, biochemical activities and also generation of -SH group containing enzymes [5].

The aim of this study was to determine Cd toxicity on morphological index parameters (MIP), contents of total chlorophyll, protein and carbohydrates in submerged aquatic macrophyte, *Pistia stratiotes*.

Materials and Methods

Pistia stratiotes, a free floating aquatic plant from unpolluted water bodies and maintained in cement pots (1 m diameter) under natural conditions at a temperature 28-30°C. About 20 g of young healthy *Pistia* is acclimatized for two weeks in Arnon and Hoagland nutrient solution [6] maintaining pH between 7.1-7.4. The concentrations of Cd in the polluted waters are in the range of 0.1, 0.5, 1.0, 1.5 and 2.0 mg/l and tap water as a control. Morphological Index Parameters (MIP) viz; root length, leaf length and breadth were observed for 12 days at interval of 4 days. Photographs of *Pistia* treated with different concentrations of cadmium were taken by using Canon's Power Shot G2 digital camera. For the further study the plants were harvested at the end of 4, 8 and 12 days exposure and are thoroughly washed with distilled water and used for the estimation of total chlorophyll, protein and carbohydrate and also for morphological observations. Plants harvested after 48 hrs were dried at 80°C for 2 days for metal extraction.

The fresh plant sample of one g is macerated in 100 ml of 80% (v/v) chilled acetone by using pestle and mortar. The extract was centrifuged

and supernatant was used for the estimation of total chlorophyll by standard method [7] using 652 nm against the solvent (80% acetone as a blank). The protein was estimated by Lowry's method [8] using Bovine Serum Albumin (BSA) as a standard using 660 nm and carbohydrates by phenol sulphuric acid method [9] using glucose as standard at 490 nm. Morphological characteristics with the help of photographs using Canons Power Shot G₂-digital camera.

The estimation of metal (Cd) in the test plant was carried out by using standard method [10]. The dried and powdered 1 g plant material was digested by using mixed acid digestion method in Gerhardt digestion unit. The digested samples were diluted with double distilled water and filtered through Whatman filter paper No-44. The estimation of Cd was done by AAS (GBC 932 Plus Australia) with air acetylene oxidizing flame and metal hollow cathode lamp at 217.00 nm wavelength. Working standards (SISCO-Chem-Bombay Lab) were used for the calibration of instrument.

Statistical analysis

Data are presented as mean values \pm SE from two independent experiments with three replicates each. Data were subjected to Two-way ANOVA to know significance between concentrations and between exposure duration for the accumulation of heavy metal (Cd). Further, Dunet's test is also applied for multiple comparisons between control and other concentrations. Two-way ANOVA test is also extended to know the significance between concentration and duration for biochemical parameters.

***Corresponding author:** Dr. N.M. Rolli, Associate Professor, BLDEA's Degree College, Jamkhandi, 587301, Karnataka, India, Tel: 9448896839; Fax: (08353)220183; E-mail: drnmrolli@rediffmail.com

Received November 29, 2013; **Accepted** December 23, 2013; **Published** December 27, 2013

Citation: Rolli NM, Suvarna khandi SS, Gadi SB, Mulagund GS, Taranath TC (2013) Effect of Cadmium Toxicity on Aquatic Macrophyte *Pistia Stratiotes* (L.). J Environ Anal Toxicol 4: 201. doi: [10.4172/2161-0525.1000201](http://dx.doi.org/10.4172/2161-0525.1000201)

Copyright: © 2013 Rolli NM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Results and Discussions

Morphological toxicity

Morph metric assay is one of the quantitative tools for the assessment of toxicants was measured by using Morphological Index Parameter (MIP). The rate of inhibition of growth in the root and leaf is directly proportional to the concentration of Cd. Two-way ANOVA test states that the concentrations are significantly toxic at 5% level but duration is not significant. MCA test is also represent maximum deviation is at higher concentration compared to control (Table 1). The plant showed normal growth. Similar observation was made by Garg et al. [11] in *Limnanthemum cristatum* at 1 ppm concentration of Pb, Zn and Cr. The higher concentration of Cd (0.5 to 2.0 ppm) exhibited toxicity symptoms like chlorosis and leaf fall were observed, then brownish was occurred being more marked in old leaves especially at 2.0 ppm Cd concentration such adverse effect of Pb on the growth has been reported by Dogan and Saygideger [12] and Kopittke et al. [13]. Similar observations were made by Yongpisanphop et al. [14] in *Salvinia natanus* and Wolff et al. [15] in *Salvinia auriculata*.

Biochemical toxicity

Chlorophyll content is a parameter that is sensitive to heavy metal toxicity [16]. Cadmium at 0.1 mg/l found to augment chlorophyll synthesis and directly proportional to the duration, the content increased by 0.79 %, 1.04% and 1.30 % at 4, 8 and 12 days exposure compared to control pond. Similar observation has been reported by Singh et al. [2] in *Hydrilla verticillata* and Dhir and Srivastava [17] in *Salvinia natanus* at 10 mg/l of Cu, Fe, Zn, Co, Cr. The stimulations of chlorophyll synthesis may be due to phytochelatin (PCs) which plays role in detoxification [18]. However, the higher concentration of Cd found to inhibit the chlorophyll synthesis. The inhibition at 2.0 mg/l Cd by 20.05 %, 31.49 % and 39.58 % (significant at P>0.95) at 4, 8 and 12 days exposure respectively compared to control. Two-way ANOVA represents biochemical toxicity to the test plant, concentrations are significant at P>0.01 level but duration is not significant. Two-way ANOVA represents, cadmium toxicity is at P>0.01 level significant towards but duration is not significant (Figure 1A).

Higher metal concentration, Pb 20 mg/l and 5.0 mg/l Cd showed decrease in total chlorophyll [2]. The decline in chlorophyll content in plants exposed to Cd due to i) inhibition of important enzymes associated with chlorophyll biosynthesis ii) peroxidation of chloroplast membranes resulting from heavy metal induced oxidative stress and iii) formation of metal substituted chlorophyll [19].

Carbohydrate content of *Pistia* at 0.1 ppm increased to the extent of 8.82 %, 11.42 % and 13.15 % respectively at 4, 8 and 12 days exposure

compared to control. It is believed that Cd and Pb are inducers for phytochelatin synthesis and definite role of detoxification of Cd and Pb and hence increase in the carbohydrate content at lower (0.1 ppm) concentration [20]. However, the high concentration of Cd found to inhibit carbohydrate. The inhibition at 2 mg/l of Cd by 47.05 %, 31.49 % and 39.58 % respectively at 4, 8 and 12 days exposure compared to control.




Figure 1: Biochemical effects of Cadmium on *Pistia*. (A) Total Chlorophyll (B) Carbohydrate (C) Protein.

Conc (ppm)	Exposure Duration (in days)									
	4			8			12			
	Root length			Leaf size			Leaf size			
	Length	Breadth	Length	Breadth	Length	Breadth	Length	Breadth	Length	Breadth
Control	6.1 ± 0.04	6.16 ± 0.07	6.3 ± 0.12	1.4 ± 0.04	1.33 ± 0.02	1.46 ± 0.02	1.36 ± 0.02	1.47 ± 0.02	1.37 ± 0.04	1.37 ± 0.04
0.1	7.0 ± 0.072	7.13 ± 0.03	7.33 ± 0.02	1.66 ± 0.02	1.533 ± 0.02	1.66 ± 0.02	1.53 ± 0.04	1.73 ± 0.02	1.66 ± 0.02	1.66 ± 0.02
0.5	6.36 ± 0.07	6.26 ± 0.07	6.1 ± 0.047	1.5 ± 0.047	1.333 ± 0.027	1.5 ± 0.047	1.33 ± 0.02	1.33 ± 0.02	1.33 ± 0.02	1.11 ± 0.02
1.0	4.76 ± 0.04	4.6 ± 0.08	4.29 ± 0.04	1.4 ± 0.047	1.333 ± 0.027	1.4 ± 0.047	1.33 ± 0.02	1.3 ± 0.02	1.3 ± 0.02	1.21 ± 0.02
1.5	4.53 ± 0.07	4.46 ± 0.15	3.24 ± 0.04	1.33 ± 0.02	1.066 ± 0.720	1.2 ± 0.00	1.06 ± 0.07	1.0 ± 0.00	1.0 ± 0.00	1.1 ± 0.027
2.0	4.0 ± 0.355	3.16 ± 0.02	2.9 ± 0.35	1.2 ± 0.047	0.833 ± 0.027	1.0 ± 0.00	0.81 ± 0.07	0.7 ± 0.00	0.7 ± 0.00	0.6 ± 0.072

Values are expressed in cms
Mean-values ± standard Error

Table 1: Morphological response of *Pistia stratiotes* to Cadmium.




Figure 2: Accumulation profile of Cadmium by *Pistia*.

62.85% and 74.35% (significant at $P > 0.05$) at 4, 8 and 12 days exposure respectively compared to control. Two way statistical ANOVA represents biochemical toxicity to the test plant concentrations are significant at $P > 0.01$ level but duration is not at significant level (Figure 1B).

Many studies show that the protein content of many aquatic macrophytes was increased by accumulation of Pb at lower toxicity concentration [21]. The protein content increase marginally at 0.1 ppm at 2.22%, 8.33% and 12.24% respectively at 4, 8 and 12 days exposure compared to control. The stimulation of protein synthesis at lower concentration of Cd (0.1ppm) may be attributed to the synthesis of stress proteins. The phytochelatin (PC) and phytochelatin synthetase bind and regulate the Cd and sequester the toxicity in the plants and thus, shows metal tolerance [18,22]. The reduction in protein content was observed with progressive increase in Cd concentration. Protein content in *Pistia* decreased by 24.44 %, 45.83% and 59.81% respectively at 4, 8 and 12 days exposure at 2.0 ppm concentration compared to control ($P > 0.01$) (Figure 1C). The Cd induces oxidative stress by generating reactive oxygen species (ROS). These disrupt cellular homeostasis, thus, enhances the production of ROS. These ROS reacts with proteins, lipids, nucleic acids causing membrane damage and enzyme inactivation [11,23]. The decrease in protein content of macrophyte may be due to the above reasons.

Metal Accumulation

Figure 2 shows the concentration of Cd accumulated in *Pistia*. There was increase in the accumulation may be due to availability of increased number of binding sites for the complexation of heavy metal ions leading to the increased absorption. In the test plant the accumulation of Cd is directly proportional to exposures and concentrations. The *Pistia* grown in experimental pond containing 0.1 ppm accumulate 112.50 µg/gm, 130.75 µg/gm and 133.75 µg/gm and accumulation at high concentration (2.0) was 1270.0 µg/gm, 1375.25 µg/gm and 1381.00 µg/gm during 4, 8 and 12 days exposures respectively. It was observed that the rate of accumulation is maximum at 4 days exposure irrespective of concentrations; however, at remaining durations it is marginal. Similar observations were made by Bendra et al. [24] in *Cladophora glomerata* at the concentration 0.1 M solution Cd Initial increase in the accumulation may be due to the availability of increased number of binding sites for the complexation of heavy metal ions leading to the increased absorption, however, slow accumulation may be attributed to binding almost all ions to the plants and establishment of equilibrium status between adsorbate and adsorbent [3,25].

Two way ANOVA represents concentration and exposure duration are significant at $P < 0.01$ and Dunet's test reveals the concentrations are significantly differ with control.

Acknowledgement

The authors are thankful to the Principal, BLDEA's Degree College Jamkhandi (Karnataka) and also to the Chairman, PG Department Botany, Karnataka University, Dharwad for providing necessary facilities to carry out research work.

References

1. Meagher RB (2000) Phytoremediation of toxic elemental and organic pollutants. *Curr Opin Plant Biol* 3: 153-162.
2. Singh A, Kumar CS, Agarwal A (2011) Phytotoxicity of Cadmium and Lead in *Hydrilla verticillata* (L.F) Royle. *Journal of Physiology* 3: 01-04.
3. Sbihi K, Cherifi O, El gharmali A, Oudra B, Aziz F (2012) Accumulation and toxicological effects of cadmium, copper and zinc on the growth and photosynthesis of the fresh water diatom *Planorhodium lanceolatum* (Brebisson) Lange-Bertalot: A laboratory study. *J Mater Environ Sci* 3: 497-506.
4. Van Asche F, Clijsters H (1990) Effects of metals on enzyme activity in plants. *Plant Cell Environ* 13: 195-206.
5. Vestena S, Cambraia J, Ribeiro C, Oliveira JA, Oliva MA (2011) Cadmium induced Oxidative stress and antioxidative enzyme response in water hyacinth and *Salvinia*. *Braz J Plant Physiol* 23: 131-139.
6. Arnon DI, Hoagland DR (1940) Crop production in artificial culture solutions and in soil with special reference to factor influencing yields and absorption of inorganic nutrients. *Soil Sci* 50: 463-483.
7. Arnon DI (1949) Copper enzymes in isolated chloroplasts. polyphenoloxidase in *Beta vulgaris*. *Plant Physiol* 24: 1-15.
8. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193: 265-275.
9. Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F (1956) Colorimetric method for determination of sugars and related substances. *Anal Chem* 28: 350-356.
10. Allen SE, Grimshaw HM, Parkinson JA, Quarmby C (1974) Chemical analysis of ecological materials. Blackwell Scientific Publications, Oxford.
11. Garg P, Chandra P, Devi S (1994) Chromium (VI) induced morphological changes in *Limnanthemum cristatum* Griseb: A possible bioindicator. *Phytomorphology* 44: 201-206.
12. Dogan M, Saygideger SD, Colak U (2009) Effect of lead toxicity on aquatic macrophyte *Elodea canadensis* Michx. *Bull Environ Contam Toxicol* 83: 249-254.
13. Kopitke PM, Asher CJ, Kopitke RA, Menzies NW (2007) Toxic effects of Pb²⁺ on growth of cowpea (*Vigna unguiculata*). *Environ Pollut* 150: 280-287.
14. Yongpisanphop J, Kruatrachue M, Pokethitoyook P (2005) Toxicity and accumulation of lead and chromium in *Hydrocotyle umbellata*. *J Environ Biol* 26: 79-89.
15. Wolff G, Pereira GC, Castro EM, Louzada J, Coelho FF (2012) The use of *Salvinia auriculata* as a bioindicator in aquatic ecosystems: biomass and structure dependent on the cadmium concentration. *Braz J Biol* 72: 71-77.
16. Gupta P, Chandra P (1996) Response of Cadmium to *Ceratophyllum demersum* L., a rootless submerged plant. *Waste Manag* 116: 335-337.
17. Dhir B, Srivastava S (2013) Heavy metal tolerance in metal hyperaccumulator plant, *Salvinia natans*. *Bull Environ Contam Toxicol* 90: 720-724.
18. Rolli NM, Suvamkhandi SS, Mulgund GS, Ratageri RH, Taranath TC (2010) Biochemical responses and accumulation of cadmium in *Spirodela polyrhiza*. *J Environ Biol* 31: 529-532.
19. Patsikka E, Kairavuo M, Sersen F, Aro EM, Tyystjarvi E (2002) Excess copper predisposes photosystem II to photoinhibition in vivo by outcompeting iron and causing decrease in leaf chlorophyll. *Plant Physiol* 129: 1359-1367.
20. Bhattacharya M, Choudhary MA (1995) Heavy metal (Pb²⁺ and Cd²⁺) stress induced damages in *Vigna* seedlings and possible involvement of phytochelatin like substances in mitigation of heavy metal stress. *Ind J Exp Biol* 33: 236-238.
21. Mohan BS, Hosetti BB (1997) Potential phytotoxicity of lead and cadmium to *Lemna minor* grown in sewage stabilization ponds. *Environ Pollution* 98: 233-238.

22. Steffens JC (1997) The heavy metal binding peptides of plants. Ann Rev Plant Physiol Plant mol Biol 41: 553-575.
23. Romero-Puertas MC, Corpas FJ, Rodriguez-Seranno M, Gornez M, Del Rio LA, et al. (2007) Differential expression regulation of antioxidative enzymes by cadmium in pea plants. J Plant Physiol 164: 1346-1357.
24. Mc Hardy BM, George JJ (1990) Bioaccumulation and toxicity of Zn in the green alga, *Cladophora glomerata*. Environ Pollut 66: 55-66.
25. Rai AK, Kumar S (1999) Removal of Cr (VI) by low cost dust adsorbents. Applied Microbiol Biotechnol 39: 661-667.

Citation: Rolli NM, Suvarna khandi SS, Gadi SB, Mulagund GS, Taranath TC (2013) Effect of Cadmium Toxicity on Aquatic Macrophyte *Pistia Stratiotes* (L.). J Environ Anal Toxicol 4: 201. doi: [10.4172/2161-0525.1000201](https://doi.org/10.4172/2161-0525.1000201)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission/

