

Case Report

Open Access

First Detection of Hypercholesterolemia Causing ApoB-100 R3527Q Mutation in a Family in Greece

Chatzistefanidis D^{1,2*}, Markoula S¹, Vartholomatos G³, Milionis HJ⁴, Miltiadous G⁴, Georgiou I², Elisaf MS⁴ and Kyritsis AP¹

¹Department of Neurology, Medical School, University of Ioannina, Ioannina, Greece

²Medical Genetics and Assisted Reproduction, Medical School, University of Ioannina, Ioannina, Greece

³Department of Haematology, University Hospital of Ioannina, Greece

⁴Department of Internal Medicine, University of Ioannina, Ioannina, Greece

Abstract

Familial defective apolipoprotein B (FDB) is an autosomal dominant genetic disorder causing hypercholesterolemia in affected patients. It is occurring due to mutation of apoB gene leading to a decreased low density lipoprotein (LDL) particles clearance. The R3527Q mutation is one of the disease's causative mutations. Data support mutation's origin 6000-7000 years ago in Central Europe and its prevalence is decreasing in relation to the distance from where initially occurred. Until now, the presence of R3527Q mutation had not been confirmed in Greece. This is the first report of FDB due to R3527Q mutation in a family in Northwest Greece.

Keywords: ApoB-100; Dyslipidemia; Familial defective apolipoprotein B; R3527Q

Introduction

Familial hypercholesterolemia (FH) is a common autosomal dominant genetic disorder leading to plasma accumulation of low-density lipoprotein (LDL) cholesterol in affected patients. It is also characterized by tendon xanthomatosis and premature atherosclerosis associated with cardiovascular disease. Symptoms are mainly a result of abnormal cholesterol deposition due to ineffective LDL clearance. LDL particles are removed from circulation through binding to LDL receptors at liver cells. Apolipoprotein B-100 (ApoB-100) is the protein component of LDL particles which serves as a ligand to LDL receptors. Mutations of the LDL receptor or ApoB-100 (familial defective apolipoprotein B or FDB) may lead to defective LDL clearance and subsequent FH [1]. Recently, mutations of proprotein convertase subtilisin/kexin type 9 (PCSK9) genes have also been associated with FH [2].

Various mutations of ApoB-100 have been related with FDB: R3480W [3], R3527Q [4], R3500W [5], R3531C [6] and H3543Y [7]. The glutamine to arginine substitution at position 3527 (R3527Q, previously R3500Q) can be found among FDB patients of Caucasian origin with noteworthy geographical spread. The R3527Q mutation prevalence is higher in Central Europe (Rhein-Main area of Germany, Switzerland and Belgium) [8-11] and becomes less common as distance from Central Europe increases, suggesting a common ancestor in this area. Data regarding its prevalence in Eastern and Southeast Europe were scarce. Recently, the R3527Q mutation was found in Bulgaria [12] but it was absent from subjects screened in Turkey [13-15]. A study from Greece [16] also failed to detect R3527Q mutation.

In this report, we present the first family carrying the R3527Q mutation found in Greece, further expanding the R3527Q mutation spread in Europe, with the description of the phenotype.

Clinical Report

A 42-years-old patient, attending the Outpatient Lipid Clinic of University Hospital of Ioannina, Northwest Greece and receiving lipid lowering treatment, was diagnosed with FH, with his sons, 2 and 5 years old respectively, found with dyslipidemia as well. The patient

was additionally attending the epilepsy outpatient Clinic because of epilepsy.

The patient was further tested for genetic characterization of his disease and was found to be heterozygous for the R3527Q mutation of ApoB-100 protein. Consequently, we further studied all available family members to determine mutation distribution in the family. Table 1 shows their lipid profile. All procedures were in accordance with Helsinki Declaration. Whole blood (EDTA-anticoagulated) was collected for genetic analysis from all subjects who gave their consent. Genomic DNA was extracted from whole blood. CVD StripAssay™ of ViennaLab Diagnostics GmbH was used for genotyping. PCR amplification was done by using biotinylated primers and hybridization of amplification products by a test strip containing allele specific oligonucleotide probes immobilized as an array of parallel lines. Bound biotinylated sequences are detected using streptavidin-alkaline phosphatase and color substrated. Analysis of clinical chemistry parameters of serum was carried out on an Olympus AU2700 analyzer (Olympus Diagnostica, Hamburg, Germany) by standard procedures. Total cholesterol and triglycerides were determined enzymatically and HDL-cholesterol by a direct assay (Olympus Diagnostica, Hamburg, Germany).

As seen in Figure 1, patient's mother was also detected as heterozygous for the R3527Q mutation. Mutation was also found in one patient's brother and one sister, who were screened, as well as in patient's both sons. Rest of the family members (marked with a "?" in Figure 1) were not screened as either we were not able to reach them or they denied to be screened. All known heterozygous family

***Corresponding author:** Chatzistefanidis Dimitrios, Department of Neurology, Medical School, University of Ioannina, Ioannina, Greece, Tel: 00306970051261; Fax: 00302651099630; E-mail: dimihat@otenet.gr

Received June 15, 2013; Accepted July 01, 2013; Published July 04, 2013

Citation: Chatzistefanidis D, Markoula S, Vartholomatos G, Milionis HJ, Miltiadous G, et al. (2013) First Detection of Hypercholesterolemia Causing ApoB-100 R3527Q Mutation in a Family in Greece. J Genet Syndr Gene Ther 4: 155. doi:10.4172/2157-7412.1000155

Copyright: © 2013 Chatzistefanidis D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

members are followed up because of dyslipidemia. Affected patients are effectively treated with statins. They were asked to inform other family members for regular lipid blood testing and/or genotyping in case of dyslipidemia.

Patient and his family live in a region in Northwest Greece, near

Albania and FYROM (Figure 2), where the mutation has been detected.

Discussion

ApoB-100 protein is a very large molecule consisting of 4536 aminoacids with a total molecular mass of about 500 kDa. Gene

	Mother	Patient	Son 1	Son 2	Brother	Sister
Total Cholesterol	212 mg/dL	268 mg/dL	243 mg/dL	240 mg/dL	262 mg/dL	246 mg/dL
Triglycerides	206 mg/dL	171 mg/dL	140 mg/dL	42 mg/dL	193 mg/dL	69 mg/dL
HDL	38 mg/dL	47 mg/dL	42 mg/dL	42 mg/dL	44 mg/dL	55 mg/dL
LDL	133 mg/dL	186,8 mg/dL	173 mg/dL	190 mg/dL	179 mg/dL	177 mg/dL
Non-HDL	N/A	221 mg/dL	201 mg/dL	N/A	N/A	N/A
ApoA1	N/A	122 mg/dL	103 mg/dL	N/A	N/A	N/A
ApoB	N/A	165 mg/dL	151 mg/dL	N/A	N/A	N/A
Lp(a)	N/A	72 mg/dL	59 mg/dL	N/A	N/A	N/A

N/A: Not Available

Table 1: Lipid profile of R3527Q affected patients attending Outpatient Lipid Clinic.




Figure 1: Family tree of Q3527R affected patient.





Figure 2: Geographical location of family affected from Q3527R mutation in Greece.



encoding apoB-100 resides in chromosome 2 (2p24-p23) and consists of 28 introns and 29 exons. As shown in Figure 3, a large part of the protein is coded by exon 26, where most common mutations are to be found [17]. The R3527Q mutation leads to a glutamine-for-arginine substitution at position 3527 of ApoB-100 protein. Mutation is located at position 10708 in exon 26 of apoB gene [4].

Familial hypercholesterolemia is characterized by high plasma concentrations of LDL cholesterol. This lipoprotein consists of neutral lipids and a shell of phospholipids, cholesterol and apoB-100 protein. LDL clearance from circulation is achieved by binding to LDL receptors in liver via apoB-100. While FH has been found to be a result of defective variants or lack of LDL-receptors, FDB is a result of lower affinity of ApoB-100 for the LDL-receptor [18]. The glutamine-for-arginine substitution caused by R3527Q mutation is located in LDL receptor-binding domain of apoB-100 protein [4]. Findings support an altered secondary or tertiary structure of apoB-100 receptor-binding domain resulting to lower affinity and decreased LDL clearance [19].

The R3527Q mutation is almost exclusively found in Caucasian population. Its prevalence varies among different populations ranging from 1:71 [10] to 1:1250 [1]. The fact that almost all affected families have the mutation on the same haplotype, characterized as 194, lead Myant et al. [8] to suggest that this mutation originate even before 6000-7000 years. Other authors [20,21], using mutation's prevalence, expanded this theory and supported that initial mutation arose in Celtic populations of Central Europe and followed their expansion and migration across Europe. This is supported by mutation's decreasing prevalence with increasing distance from Central Europe. Especially in South Europe, this mutation has a very low prevalence in Italy [22] and Spain [23]. It is of interest that mutation prevalence in Spain seems to be greater in Galicia, a region in northwest Spain, known to have been inhabited by Celts during the 11th century BC [23]. Reports from

Eastern Europe support the mutation's spread toward this area [24-28]. Recently, a report from Bulgaria supported mutation's existence in Southeast Europe. Authors reported an estimated prevalence of 1:451 in adult Bulgarians [12].

The mutation is almost universally absent from non-European populations around Europe. Researchers failed to detect the mutation in populations from Turkey [13-15], Lebanon [29], Morocco [30] and Iran [31]. Overseas, mutation can be found mainly in subjects of Caucasian origin. Of interest is a report regarding a very high prevalence of R3527Q mutation in the Older Order Amish of Lancaster County, Pennsylvania? As they are descendants of approximately 300 founders from Switzerland, data support a strong founder effect [32].

Regarding Greece, data are limited in only one report with molecular characterization of FH in 100 patients. Authors failed to detect R3527Q mutation [16] in the Greek population. To our knowledge, this is the first report for known R3527Q carriers in Greece; further supporting the theory of mutation's spread in European populations after its initial appearance in Central Europe. Our report fills the gap between mutation's presence in Bulgarian and absence from Turkish population and is a good example of how genetics can help understanding population movements around Europe.

Conclusion

The R3527Q mutation is a known mutation responsible for FDB. Its prevalence supports that mutation first arose in Central Europe and was further spread among European populations. Until now, data regarding its presence in Greek population were missing. To our knowledge, this is the first report confirming its presence in subjects of Greek origin. Further, large scale studies are needed to confirm its prevalence in various parts of Greece.

Conflict of Interest

Every author declares no conflict of interest.

References

- Hansen PS (1998) Familial defective apolipoprotein B-100. *Dan Med Bull* 45: 370-382.
- Abifadel M, Varret M, Rabès JP, Allard D, Ouguerram K, et al. (2003) Mutations in PCSK9 cause autosomal dominant hypercholesterolemia. *Nat Genet* 34: 154-156.
- Borén J, Ekström U, Agren B, Nilsson-Ehle P, Innerarity TL (2001) The molecular mechanism for the genetic disorder familial defective apolipoprotein B100. *J Biol Chem* 276: 9214-9218.
- Soria LF, Ludwig EH, Clarke HR, Vega GL, Grundy SM, et al. (1989) Association between a specific apolipoprotein B mutation and familial defective apolipoprotein B-100. *Proc Natl Acad Sci U S A* 86: 587-591.
- Gaffney D, Reid JM, Cameron IM, Vass K, Caslake MJ, et al. (1995) Independent mutations at codon 3500 of the apolipoprotein B gene are associated with hyperlipidemia. *Arterioscler Thromb Vasc Biol* 15: 1025-1029.
- Pullinger CR, Hennessy LK, Chatterton JE, Liu W, Love JA, et al. (1995) Familial ligand-defective apolipoprotein B. Identification of a new mutation that decreases LDL receptor binding affinity. *J Clin Invest* 95: 1225-1234.
- Soufi M, Sattler AM, Maerz W, Starke A, Herzum M, et al. (2004) A new but frequent mutation of apoB-100-apoB His3543Tyr. *Atherosclerosis* 174: 11-16.
- Myant NB, Forbes SA, Day IN, Gallagher J (1997) Estimation of the age of the ancestral arginine3500->glutamine mutation in human apoB-100. *Genomics* 45: 78-87.
- Kotze MJ, Peeters AV, Langenhoven E, Wauters JG, Van Gaal LF (1994) Phenotypic expression and frequency of familial defective apolipoprotein B-100 in Belgian hypercholesterolemic. *Atherosclerosis* 111: 217-225.
- Fisher E, Schrnagl H, Hoffmann MM, Kusterer K, Wittmann D, et al. (1999)

- Mutations in the apolipoprotein (apo) B-100 receptor-binding region: detection of apo B-100 (Arg3500-->Trp) associated with two new haplotypes and evidence that apo B-100 (Glu3405-->Gln) diminishes receptor-mediated uptake of LDL. *Clin Chem* 45: 1026-1038.
11. Miserez AR, Laager R, Chiodetti N, Keller U (1994) High prevalence of familial defective apolipoprotein B-100 in Switzerland. *J Lipid Res* 35: 574-583.
 12. Horvath A, Savov A, Kirov S, Karshelova E, Paskaleva I, et al. (2001) High frequency of the ApoB-100 R3500Q mutation in Bulgarian hypercholesterolaemic subjects. *J Med Genet* 38: 536-540.
 13. Tamer L, Tanriverdi K, Ercan B, Unlu A, Sucu N, et al. (2004) Apolipoprotein B gene polymorphisms in people in the east Mediterranean area of Turkey. *East Mediterr Health J* 10: 125-130.
 14. EroÄŸlu Z, Selvi N, Kosova B, Biray C, Kumral E, et al. (2008) Absence of apolipoprotein B-3500 mutation in Turkish patients with coronary and cerebrovascular atherosclerosis. *Anadolu Kardiyol Derg* 8: 7-9.
 15. SÄŸzen MM, Whittall R, Oner C, Tokatli A, KalkanoÄŸlu HS, et al. (2005) The molecular basis of familial hypercholesterolaemia in Turkish patients. *Atherosclerosis* 180: 63-71.
 16. Dedoussis GV, Genschel J, Bochow B, Pitsavos C, Skoumas J, et al. (2004) Molecular characterization of familial hypercholesterolemia in German and Greek patients. *Hum Mutat* 23: 285-286.
 17. Blackhart BD, Ludwig EM, Pierotti VR, Caiati L, Onasch MA, et al. (1986) Structure of the human apolipoprotein B gene. *J Biol Chem* 261: 15364-15367.
 18. Rader DJ, Cohen J, Hobbs HH (2003) Monogenic hypercholesterolemia: new insights in pathogenesis and treatment. *J Clin Invest* 111: 1795-1803.
 19. Lund-Katz S, Innerarity TL, Arnold KS, Curtiss LK, Phillips MC (1991) 13C NMR evidence that substitution of glutamine for arginine 3500 in familial defective apolipoprotein B-100 disrupts the conformation of the receptor-binding domain. *J Biol Chem* 266: 2701-2704.
 20. Miserez AR, Muller PY (2000) Familial defective apolipoprotein B-100: a mutation emerged in the mesolithic ancestors of Celtic peoples? *Atherosclerosis* 148: 433-436.
 21. Fisher E, Gross W, März W (2000) High prevalence of FDB3500 mutation in the Swiss population. *Atherosclerosis* 153: 519-521.
 22. Cefalù AB, Barbagallo CM, Sesti E, Caldarella R, Polizzi F, et al. (2001) Italian familial defective apolipoprotein B patients share a unique haplotype with other Caucasian patients. *Clin Exp Med* 1: 151-154.
 23. Castillo S, Tejedor D, Mozas P, Reyes G, Civeira F, et al. (2002) The apolipoprotein B R3500Q gene mutation in Spanish subjects with a clinical diagnosis of familial hypercholesterolemia. *Atherosclerosis* 165: 127-135.
 24. Kalina A, Czeizel E, Romics L, Pados G, Reiber I, et al. (1998) [Incidence of familial defective apolipoprotein B-100 in cases of patients diagnosed with familial hypercholesterolemia]. *Orv Hetil* 139: 755-759.
 25. GÅrski B, Kubalska J, Naruszewicz M, LubiÅski J (1998) LDL-R and Apo-B-100 gene mutations in Polish familial hypercholesterolemias. *Hum Genet* 102: 562-565.
 26. Ceska R, Vrablík M, Horínek A (2000) Familial defective apolipoprotein B-100: a lesson from homozygous and heterozygous patients. *Physiol Res* 49 Suppl 1: S125-130.
 27. Gasparovic J, Basistová Z, Fábryová L, Wsólóv L, Vohnout B, et al. (2007) Familial defective apolipoprotein B-100 in Slovakia: are differences in prevalence of familial defective apolipoprotein B-100 explained by ethnicity? *Atherosclerosis* 194: e95-107.
 28. Meshkov AN, Malyshev PP, Kukharchuk VV (2009) [Familial hypercholesterolemia in Russia: genetic and phenotypic characteristics]. *Ter Arkh* 81: 23-28.
 29. Sabbagh AS, Daher RT, Otrock ZK, Khalek RN, Zaatari GS, et al. (2007) ApoB-100 R3500Q mutation in the Lebanese population: prevalence and historical review of the literature. *Mol Biol Rep* 34: 267-270.
 30. Chater R, Ait Chihab K, Rabès JP, Varret M, Chabraoui L, et al. (2006) Mutational heterogeneity in low-density lipoprotein receptor gene related to familial hypercholesterolemia in Morocco. *Clin Chim Acta* 373: 62-69.
 31. Farrokhi E, Shayesteh F, Asadi Mobarakeh S, Roghani Dehkordi F, Ghatreh Samani K, et al. (2011) Molecular characterization of Iranian patients with possible familial hypercholesterolemia. *Indian J Clin Biochem* 26: 244-248.
 32. Shen H, Damcott CM, Rampersaud E, Pollin TI, Horenstein RB, et al. (2010) Familial defective apolipoprotein B-100 and increased low-density lipoprotein cholesterol and coronary artery calcification in the old order amish. *Arch Intern Med* 170: 1850-1855.

Citation: Chatzistefanidis D, Markoula S, Vartholomatos G, Milionis HJ, Miltiadous G, et al. (2013) First Detection of Hypercholesterolemia Causing ApoB-100 R3527Q Mutation in a Family in Greece. *J Genet Syndr Gene Ther* 4: 155. doi:10.4172/2157-7412.1000155

This article was originally published in a special issue, **Genetic Disorders** handled by Editor(s). Dr. Xiaobo Li, Yeshiva University, USA

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/omicsgroup/>