

Getting off the Mat: Lumbar Stabilization Training in Standing Using Ultrasound Imaging

Todd Watson* and Sue McPherson

Department of Physical Therapy, Western Carolina University, USA

The abdominal drawing-in maneuver (ADIM) has been explained as the optimal method for activation of the transversus abdominus (TrA) muscle[1], whose importance in controlling motion (stabilizing) of the lumbar spine is well-known[2,3]. The ADIM is boilerplate physiotherapy exercise in a traditional stabilization program for low back pain (LBP) patients and is often one of the first exercises instructed to patients in a lumbar stabilization exercise program prior to exercise progression in non-weight bearing and then in weight bearing[4]. Biofeedback has been recommended for low back pain patients with impaired activation of the TrA[5-7]. Ultrasound imaging (USI) visualizes the “corset-like action” or lateral sliding and thickening of the TrA muscle during activation of TrA[8]. With USI utilization in physiotherapist practice, patients are able to see changes in their TrA cross-sectional area, and clinicians are able to assess the muscle recruitment pattern[9]. The USI visualization is important for us, because in our experience many clients initially believe they are proficient with “core activation” - until they are able to visualize their incorrect muscle activation pattern. This training of the TrA has been shown to improve pain and function in patients with chronic LBP[10-12]. The success of ADIM training programs in healthy adult populations during functional loaded tasks has developed since such exercising may also lead to injury prevention of LBP[6]. Acquiring correct lumbar stabilization is typically verified by eliminating the visual USI feedback while doing the same exercise maneuvers to evaluate skill retention, or another task to assess skill transfer.

Historically, supine hook lying postures and exercises were chosen for initial spinal position during ADIM training and skill acquisition. This was so as to enhance ease of performance and acquisition. However, both the TrA and the internal oblique muscles are noted to increase in thickness in response to the weight-bearing tasks over non-weight bearing tasks[13].

Recently, a number of studies have begun to examine USI training of subjects acquiring ADIM proficiency during performance of upright standing, functional tasks so as to provide in situ training for those postures typically involved in injury to the lumbar spine. Performance reproducibility has been established for an entire experimental protocol involving USI measurements of TrA activation during loaded standing and functional activities[14]. Additionally, acceptable reliability has been shown for USI measurement of TrA muscle thickness, and thickness changes taken during loaded and functional activities[15]. This is important as reliable USI measures of TrA muscle thickness changes meet the continuing necessity for clinicians' knowledge of correct and appropriately intense muscle activation in the clinical setting to assist progression of lumbar stabilization exercises.

Considering that adults on a daily basis typically perform tasks such as an extended forward reach with a weighted object, which proves to be problematic to those with clinical lumbar instability, we included this task in our recent experimental protocol of ADIM acquisition in healthy subjects[16]. Minimal time was required for the ADIM training, and the ADIM technique, taught in the supine position, translated into use of this technique during upright loaded functional tasks (e.g. lifting a weighted box, holding a weighted object with arms extended) for

at least five months without any monitoring of compliance. In terms of cost effective injury prevention strategies, it seems that our ADIM program is viable. Injury prevention training in this area will continue to grow, along with advances in the quality of USI equipment in terms of portability, cost, user software, and quality of images. For example, USI software is now providing higher-quality resolutions of images of TrA muscle activation in real time and easier retrieval of past performances.

These images provide a wide array of various forms of augmented feedback. The client views the thickening of his/her TrA during or immediately following performances of functional tasks, and the therapist tells the client the number of successful TrA contractions following performance of the tasks. USI also contributes to goal setting whereby the client and therapist collaborate on strategies to increase TrA thickness while viewing and not viewing the visual display of his/her TrA. Additionally, both goal setting and augmented feedback may be combined. Here for example, the client is provided with a goal to increase TrA contraction by a specific amount and estimates whether or not this goal was met following task performances. Taken together, these tactics may expedite motor learning and facilitate clinical outcomes.

Based on our findings of upright training of ADIM in healthy subjects, we recommend moving beyond the traditional non-weight bearing initialization of stabilization exercises as soon as possible for the patient; commencing an upright stabilization, functional activity approach as soon as symptoms allow for LBP stabilization patients.

We expect both rehabilitation and injury prevention programs to grow in their use of USI as feedback, albeit slowly as current use in physiotherapy remains relatively small[17]. Also, we found those individuals to be highly motivated to learn how to activate their TrA muscle during tasks, because TrA muscle feedback via USI was new and challenging to them, akin to video game play. Notably, the application of motor learning principles and use of retention and transfer of learning testing will be vital in design and analysis when creating injury prevention programs for healthy adults[18,19].

Finally, we believe training programs that adhere to motor learning principles may help to alleviate compliance issues if enough task appropriate practice is incorporated in training and community programs. This will help to ensure clients reach the point where ADIM technique is incorporated automatically in their activities of daily living.

*Corresponding author: Todd Watson, Professor of Physical Therapy, Department of Physical Therapy, Western Carolina University, USA, Tel: 828-227-2126; E-mail: twatson@email.wcu.edu

Received August 15, 2013; Accepted December 09, 2013; Published December 12, 2013

Citation: Watson T, McPherson S (2013) Getting off the Mat: Lumbar Stabilization Training in Standing Using Ultrasound Imaging. J Yoga Phys Ther 3: 147. doi:10.4172/2157-7595.1000147

Copyright: © 2013 Watson T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

References

1. Urquhart DM, Hodges PW, Allen TJ, Story IH (2005) Abdominal muscle recruitment during a range of voluntary exercises. *Man Ther* 10: 144-153.
2. Hodges P, Kaigle Holm A, Holm S, Ekstrom L, et al. (2003) Intervertebral stiffness of the spine is increased by evoked contraction of transversus abdominis and the diaphragm: in vivo porcine studies. *Spine* 28: 2594-2601.
3. Hodges PW, Sapsford R, Pengel LH (2007) Postural and respiratory functions of the pelvic floor muscles. *NeuroUrol Urodyn* 26: 362-371.
4. Richardson C, Hodges P, Hides J (2004) *Therapeutic Exercise for Lumbopelvic Stabilization: A Motor Control Approach for the Treatment and Prevention of Low Back Pain*. Edingurgh, Churchill Livingstone, Australia.
5. Henry SM, Westervelt KC (2005) The use of real-time ultrasound feedback in teaching abdominal hollowing exercises to healthy subjects. *J Orthop Sports Phys Ther* 35: 338-345.
6. Anderson Worth SG, Henry SM, Bunn JY (2007) Real-time ultrasound feedback and abdominal hollowing exercises for people with low back pain. *NZ J Physiother* 35: 4-11.
7. Hodges PW, Pengel LH, Herbert RD, Gandevia SC (2003) Measurement of muscle contraction with ultrasound imaging. *Muscle Nerve* 27: 682-692.
8. Mannion AF, Pulkovski N, Gubler D, Gorelick M, O'Riordan D, et al. (2008) Muscle thickness changes during abdominal hollowing: an assessment of between-day measurement error in controls and patients with chronic low back pain. *Eur Spine J* 17: 494-501.
9. Richardson CA, Snijders CJ, Hides JA, Damen L, Pas MS, et al. (2002) The relation between the transversus abdominis muscles, sacroiliac joint mechanics, and low back pain. *Spine* 27: 399-405.
10. O'Sullivan PB, Phytz GD, Twomey LT, Allison GT (1997) Evaluation of specific stabilizing exercise in the treatment of chronic low back pain with radiologic diagnosis of spondylolysis or spondylolisthesis. *Spine* 22: 2959-2967.
11. Hides JA, Richardson CA, Jull GA (1996) Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine* 21: 2763-2769.
12. O'Sullivan P, Twomey L, Allison G, Sinclair J, Miller K (1997) Altered patterns of abdominal muscle activation in patients with chronic low back pain. *Aust J Physiother* 43: 91-98.
13. Hides JA, Wong I, Wilson SJ, Belavy DL, Richardson CA (2007) Assessment of abdominal muscle function during a simulated unilateral weight-bearing task using ultrasound imaging. *J Orthop Sports Phys Ther* 37: 467-471.
14. McPherson SL, Watson T (2012) Reproducibility of ultrasound measurement of transversus abdominis during loaded, functional tasks in asymptomatic young adults. *PM R* 4: 402-412.
15. Watson T, McPherson S, Fleeman S (2011) Ultrasound measurement of transversus abdominis during loaded, functional tasks in asymptomatic individuals: rater reliability. *PM R* 3: 697-705.
16. McPherson SL, Watson T (2013) Training of transversus abdominis activation in supine with ultrasound biofeedback translated to increased transversus abdominis activation during upright loaded functional tasks. *PM R*: In Press.
17. Jedrzejczak A, Chipchase LS (2008) The availability and usage frequency of real time ultrasound by physiotherapists in South Australia: an observational study. *Physiother Res Int* 13: 231-240.
18. Magill R, Anderson D (2013) *Motor learning and control: Concepts and applications*. (10th edn), McGraw-Hill, Boston, USA.
19. Schmitz RA, Lee TD (2011) *Motor control and learning: A behavioral emphasis*. (5th edn), Human Kinetics, Champaign, IL.