

## Case Report

## Open Access

# HPV6 Infection of an Infant's Penile Condyloma at the Urethral Meatus

Vageli DP<sup>1\*</sup>, Doukas SG<sup>1</sup> and Markou A<sup>2</sup>

<sup>1</sup>Department of Pathology, Medical School, University of Thessaly, Greece

<sup>2</sup>Department of Surgical Pediatric, General Hospital of Larissa, Larissa, Greece

### Abstract

Penile warts in infants are rare and condylomata acuminata at the urethral meatus are rarer. This is a first report of a 2.5 years infant with a condyloma acuminatum at the urethral meatus of penis that was found positive for HPV 6, by PCR analysis. Infant's mother presented warts on her hands, suggesting a possible transmission of virus from mother to infant via extragenital contact, without excluding a vertical transmission.

### Introduction

Genital warts in children are infrequent while penile warts are even rarer [1-5]. The youngest documented case of this type was a 7-month-old circumcised Caucasian boy presented with a rapidly progressing giant penile condyloma acuminatum that was surgically removed followed by a fatal course of *Neisseria meningitidis*. This case serves as a warning, concerning the course of condyloma in infants, since the infant died from meningococemia a few days after surgery [5]. Condyloma acuminatum of the urethral meatus is very rare while there is only a single report of a 5-year old child presented such a lesion without HPV infection history [5]. Here we present a HPV infected penile condyloma acuminatum at the urethral meatus, of a 2.5 year old infant, with a possible virus transmission from mother to infant.

### Patient and Molecular Analysis

A 2.5 years infant, who presented a penile condyloma acuminatum at the urethral meatus, came to the Pediatric Surgery Clinic of General Hospital of Larissa (Figure 1). His mother mentioned that she had presented warts on her hands. The condyloma was surgically removed and two years after surgery the child was still healthy.

The condyloma mass was subdivided to a part that was taken for the standard histological diagnosis while the remaining tissue was immediately immersed into stabilization solution and frozen to -20°C. DNA was extracted from tissue using Puregene® Cell and Tissue extraction kit (Gentra Systems, Inc., USA) and a HPV detection molecular method was applied, as previously referred [6,7]. In brief a multiplex PCR (MPCR) was employed for detection and discrimination of HPV types 6, 11, 16, 18 and 33 (specific E6 gene) using a commercial kit (MPCR kit, Maxim Biotech) which includes specific primers for each HPV type and corresponding positive control samples. The results of HPV-positive sample were confirmed by a PCR analysis using specific primers for each HPV type only (Maxim Biotech). We used non-template negative control samples, in each PCR reaction to exclude false positive results. The PCR products were electrophorized through 2% agarose gels, stained with 0.5 mg/ml ethidium bromide and visualized on a UV light transilluminator. The detection of specific sizes bands indicated the presence of target sequences, corresponded to HPV type specific, in our specimen (Figure 2).

We performed a melting curve analysis of PCR products to confirm HPV 6 genotype. Melting Curve analysis was performed in Corbett RotorGene instrument as we have described previously [6,8], using SYBER Green (QuantiTect kit, Qiagen) and ramping 65-95°C (raising by 0.2°C each step).

### Results

PCR analysis for HPV detection and typing revealed that the DNA sample, derived from the penile condyloma acuminatum of the infant, was positive for HPV type 6 (Figure 2). Specifically, agarose gel electrophoresis of our specimen showed a 263-bp amplified PCR product, like HPV-6-positive control, corresponding to PCR amplified specific E6 gene of HPV type-6 (Figure 2). Confirmation of HPV6 genotype was done performing a melting curve analysis of penile condyloma and HPV6 positive control PCR products. Both samples showed melting peaks corresponded to HPV6 genotype (Figure 3).

### Discussion

We present a rare case of an infant with a penile condyloma acuminatum at the urethral meatus that was analyzed by PCR method and found positive for HPV type 6. In bibliography, it has been referred a single case of a condyloma acuminatum of the urethral meatus in an infant, without a history of HPV infection [5]. Genital HPV infection in children seemed to behave as a sexual transmitted via abuse [9] or vertically transmitted or extragenital contact transmitted [10,11]. In our reported case, the infant's mother mentioned that she had presented



**Figure 1:** A condyloma acuminatum at the urethral meatus of an infant's penis.

**\*Corresponding author:** Dimitra Vageli, Department of Pathology, Medical School, University of Thessaly, Larissa, Greece, Tel: 0030-2410685650; E-mail: [vagelidim@yahoo.gr](mailto:vagelidim@yahoo.gr)

**Received** June 13, 2013; **Accepted** July 02, 2013; **Published** July 04, 2013

**Citation:** Vageli DP, Doukas SG, Markou A (2013) HPV6 Infection of an Infant's Penile Condyloma at the Urethral Meatus. J Genet Syndr Gene Ther 4: 157. doi:10.4172/2157-7412.1000157

**Copyright:** © 2013 Vageli DP, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



warts on her hands. This could support a HPV 6 transmission from mother to the urethral meatus of infant's penis via extragenital contact, during infant's caring, without excluding a vertical transmission of virus.

### References

- Vardas E, Giuliano AR, Goldstone S, Palefsky JM, Moreira ED Jr, et al. (2011) External genital human papillomavirus prevalence and associated factors among heterosexual men on 5 continents. *J Infect Dis* 203: 58-65.
- Smith JS, Backes DM, Hudgens MG, Bailey RC, Veronesi G, et al. (2010) Prevalence and risk factors of human papillomavirus infection by penile site in uncircumcised Kenyan men. *Int J Cancer* 126: 572-577.
- Weaver MG, Abdul-Karim FW, Dale G, Sorensen K, Huang YT (1989) Detection and localization of human papillomavirus in penile condylomas and squamous cell carcinomas using in situ hybridization with biotinylated DNA viral probes. *Mod Pathol* 2: 94-100.
- Babich SB, Haber SD, Caviedes EY, Teplitsky P (2003) Condylomata acuminata in a boy. *J Am Dent Assoc* 134: 331-334.
- Gor RA, Schober JM (2009) Giant condyloma with demise secondary to meningococemia in an infant boy. *J Pediatr Urol* 5: 327-329.
- Vageli D, Sourvinos G, Ioannou M, Koukoulis GK, Spandidos DA (2007) High-risk human papillomavirus (HPV) in parotid lesions. *Int J Biol Markers* 22: 239-244.
- Mammas IN, Vageli D, Spandidos DA (2008) Geographic variations of human papilloma virus infection and their possible impact on the effectiveness of the vaccination programme. *Oncol Rep* 20: 141-145.
- Vageli D, Daniil Z, Dahabreh J, Karagianni E, Liloglou T, et al. (2006) Microsatellite instability and loss of heterozygosity at the MEN1 locus in lung carcinoid tumors: a novel approach using real-time PCR with melting curve analysis in histopathologic material. *Oncol Rep* 15: 557-564.
- Unger ER, Fajman NN, Maloney EM, Onyekwuluje J, Swan DC, et al. (2011) Anogenital human papillomavirus in sexually abused and nonabused children: a multicenter study. *Pediatrics* 128: e658-665.
- Handley JM, Maw RD, Bingham EA, Horner T, Bharucha H, et al. (1993) Anogenital warts in children. *Clin Exp Dermatol* 18: 241-247.
- Obalek S, Misiewicz J, Jablonska S, Favre M, Orth G (1993) Childhood condyloma acuminatum: association with genital and cutaneous human papillomaviruses. *Pediatr Dermatol* 10: 101-106.

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 250 Open Access Journals
- 20,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/omicsgroup/>

**Citation:** Vageli DP, Doukas SG, Markou A (2013) HPV6 Infection of an Infant's Penile Condyloma at the Urethral Meatus. *J Genet Syndr Gene Ther* 4: 157. doi:10.4172/2157-7412.1000157

This article was originally published in a special issue, **Cancer Genetics** handled by Editor(s). Dr. Ahmed M Malki, Alexandria University, Egypt