

¿ Is Natural Latex an Important Cause of Allergy ?

Carlos Enrique Cabrera Pivara*

Director de Educación e Investigación, Unidad Médica de Alta Especialidad, UMAE, Hospital de Especialidades, IMSS, Jalisco

Natural latex is an important cause of allergy in individuals with risk factors. Neuromuscular relaxants together with natural latex are among the most studied agents responsible for pre-operative anaphylaxis. Allergic reaction to latex represents, in most cases, an immune response regulated by IgE antibodies (type I hypersensitivity). This type of reaction is produced in individuals who have previously been in contact with latex and have formed sensitivity, the same as in other common forms of allergies. These types of reactions are also more frequent in certain risk groups, such as: health personnel, patients with congenital defects (spina bifida, urologic, myelomeningocele), those who have undergone multiple surgical interventions and history of repeated exposures -specially in serous and mucous- to materials such as latex. There is a known existence in patients with atrophy history, in rubber and plastic workers, janitors and patients with history of latex intolerance as: balls, balloons, gloves, preservatives, etc. There are cross reactions between latex and food antigens (bananas, cashews, avocado, celery, grenade, etc), these allergens have an antigenic action very similar to latex, clinically and immunologically [1,2].

Immunopathogenesis of allergy to latex is not completely understood. Several proteins have been identified from natural latex extracts that adhere to IgE. The main one being Hevein b 6, formed from prohevein (Hev b 6.01) and two Hevein post-transcriptional proteins (Hev b 6.02 and Hev b 6.03). These three components act like independent immunogens. In the original Hev b 6.01 protein are found regions responsible for the IgE binding, as well as the proliferative response of the T lymphocyte. Hev b 6.02 domain contains discontinuous epitopes that can recognize the B lymphocyte, while Hev b 6.03 is a better inducer of the proliferative response for having sequentiated unions to HLA-DR4 [3,4]. Though this it is possible to initiate an acquired immune response directed to immunogens, latex related proteins activated by allergenic-specific T CD+ lymphocytes. Identification of T cell dominant epitope is important and decisive for the development of a potential specific immunotherapy [5].

Clinical spectrum includes slight dermatitis cases that may or may not affect mucous, together with urticarial, pruritus and exanthema being the most frequent. Other forms in which it manifests are rhinitis respiratory profiles and bronchial obstruction. Only in rare occasions do lethal symptoms appear such as anaphylaxis, convulsions, frequently observed during anesthesia [3,4].

Pre-operative identification of high risk patients and preventive measures may decrease latex allergy and anaphylaxis incidences [5,7].

The severity of the immediate reaction depends on the level of sensitivity and the amount of latex in which the person comes in contact with. Greatest danger of suffering extreme reactions occurs when latex comes in contact with humid areas of skin of inner parts during surgery, due to the body absorbing more quickly a larger amount of allergen, not forgetting that latex may be suspended in the air and can cause respiratory symptoms. Latex proteins may adhere to powder used in latex gloves, in using these gloves these particles together with latex

allergens fly through the air, where they can be inhaled or come in contact with the nose or the eyes and can cause symptoms.

High levels of this type of allergen dust have been measured in operating rooms. For this reason the American Society of Anesthesiologists have recommended who patients that have been identified as allergic to latex be the first to use operating rooms, which is when the concentration of these airborne allergens (latex) is minimum, the use of non-powder or synthetic gloves is also recommended (vinyl or nitrile), reducing the risk of reaction, and the use of a physiologic solution on the gloves to remove the excess of dust [2,8-10].

For these reasons show the Natural latex is an important cause of allergy in individuals with risk factors and the most studied agents responsible for pre-operative anaphylaxis.

Reference

1. Escolano F, Sierra P (1996) [Allergic reactions during anesthesia]. *Rev Esp Anestesiol Reanim* 43: 17-26.
2. Hepner DL, Castells MC (2003) Latex allergy: an update. *Anesth Analg* 96: 1219-1229.
3. Banerjee B, Wang X, Kelly KJ, Fink JN, Sussman GL, et al. (1997) IgE from latex allergic patients binds to cloned and expressed B cell epitopes of prohevein. *J Immunol* 159: 5724-5732.
4. Heimsoth R, Rozynek P, Bruning T, Rihs HP (2004) Characterization of B and cell responses and HLA DR4 binding motifs of the latex allergen Hev b. *Allergy* 59: 724-733.
5. de Silva HD, Gardner LM, Drew AC, Beezhold DH, Rolland JM, et al. (2004) The hevein domain of the major latex-glove allergen Hev b 6.01 Contains dominant T cell reactive sites. *Clin Exp Allergy* 34: 611-618.
6. Mertes PM, Laxenaire MC, Alla F (2003) Anaphylactic and anaphylactoid reactions occurring during anesthesia in France in 1999-2000. *Anesthesiology* 99: 536-545.
7. Thong BY, Yeow Chan (2004) Anaphylaxis during surgical and interventional procedures. *Ann Allergy Asthma Immunol* 92: 619-628.
8. Laxenaire MC, Mertes PM (2001) Anaphylaxis during anaesthesia. Results of a two years survey in France. *Br J Anaesth* 87: 549-558.
9. Kemp SF, Lockey RF (2002) Anaphylaxis: a review of causes and mechanisms. *J Allergy Clin Immunol* 110: 341-348.
10. Kurek M, Michalska KG (2003) Anaphylaxis during surgical and diagnostic procedures. *Allergy Clin Immunol Int* 15:168-174.

Corresponding author: Carlos Enrique Cabrera Pivara, Director de Educación e Investigación, Unidad Médica de Alta Especialidad, UMAE, Hospital de Especialidades, IMSS, Jalisco, Tel: 36170060; E-mail: carlos.cabrera@imss.gob.mx

Received April 1, 2011; **Accepted** April 5, 2011; **Published** April 10, 2011

Citation: Pivara CEC (2011) ¿ Is Natural Latex an Important Cause of Allergy ? *J Aller Ther* 2:102e. doi:10.4172/2155-6121.1000102e

Copyright: © 2011 Pivara CEC. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.