

MicroRNAs Profiling Reveals a Potential Link between the SDG8 Methyltransferase and Brassinosteroid-Regulated Gene Expression in *Arabidopsis*

Gaofeng Dong and Jiaxu Li*

Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology, Mississippi State University, Mississippi State, USA

MicroRNAs are a class of short non-coding RNAs (17-27 nucleotides) found in animals and plants. MicroRNAs play important roles in post-transcriptional regulation of gene expression by complementing the target mRNAs and causing translational repression or target mRNA degradation [1]. Studies have shown that thousands of human protein-coding genes are regulated by microRNAs [2], impacting many important biological processes, from development to physiology to stress response. In plants, microRNAs have been implicated in multiple essential biological processes, such as leaf morphogenesis and polarity, flower development, hormone signaling and metabolism, and stress responses [3]. MicroRNA genes are transcribed by RNA polymerases II, generating precursors that undergo a series of cleavage events to form mature microRNA [4]. MicroRNA biogenesis and its expression regulation are highly complicated [5]. Although the biological importance of microRNAs is well demonstrated in a wide range of cellular processes, how microRNA expression and abundance are regulated is not fully understood.

Histone modifications play critical roles in regulation of gene expression in eukaryotes [6]. Acetylation of histones H3 and H4 is commonly associated with gene activation, whereas histone deacetylation generally leads to gene silencing. Trimethylation of histone H3 on lysine 4 (H3K4me3) is frequently associated with active transcription, while trimethylation of histone H3 on lysine 27 (H3K27me3) is often associated with gene repression. Trimethylation of histone H3 on lysine 36 (H3K36me3) is usually enriched in coding regions of actively transcribed genes [6], but H3K36me3 also correlates with gene silencing in facultative and constitutive heterochromatin [7]. Recent studies in mammalian cells indicate that histone acetylation and methylation affect microRNA expression. Scott et al. [8] first showed that histone deacetylase inhibition results in alteration of microRNA levels in a breast cancer cell line. Following this study, several reports have shown that histone deacetylase inhibition alters microRNA expression in human carcinomas [9-11]. Overexpression of histone deacetylases in chronic lymphocytic leukemia results in silencing of miR-15a, miR-16, and miR-29b, while histone deacetylase inhibition can partially restore the expression of miR-15a, miR-16, and miR-29b [12]. These results collectively show that histone acetylation and deacetylation play important roles in regulation of microRNA expression in human cells. Stable RNAi-mediated suppression of the H3K4me3 demethylase JARID1B in breast tumor cells caused increased expression of several members of the let-7 family of microRNAs, suggesting that H3K4me3 is required for up-regulation of the expression of these microRNAs [13]. Parallel sequencing analyses show that H3K27me3 is associated with repressed microRNA genes in mouse lymphocytes [14]. Taken together, these studies indicate that histone methylation can modulate microRNA expression.

The SET domain group 8 (SDG8) protein is the primary methyltransferase for global histone H3K36 trimethylation in *Arabidopsis* [15,16]. SDG8 is involved in a number of developmental processes such as shoot branching, ovule and anther development, and

flowering time [16-18]. To explore the potential role of histone H3K36 trimethylation in the regulation of microRNA expression, we compared the miRNA expression profile of the knockout mutant of the SDG8 methyltransferase (sdg8) with the wild-type (WT) using microRNA microarray technology. MicroRNA microarray analysis was carried out by LC sciences (Houston, Texas, USA) on μ Paraflo™ microfluidics chips containing 154 microRNA probes to *Arabidopsis* microRNAs (Sanger miRBase Version 9.2). Of the 154 microRNAs examined, 23 were differentially expressed in sdg8 knockout mutant. Twelve of 23 were overexpressed [\log_2 (sdg8/WT) range: 0.22 to 3.09], whereas 11 of 23 were underexpressed [\log_2 (sdg8/WT) range: -0.32 to -1.42]. All 23 differentially expressed microRNAs had a q-value <0.01, indicating that the observed changes were significantly different. Differentially expressed microRNAs in sdg8 knockout mutant with $|\log_2$ (sdg8/WT)| values greater than 1 are listed in Table 1. These results suggest that a small number of known microRNAs examined are regulated by the SDG8 methyltransferase.

Brassinosteroids are a class of plant hormones that regulate multiple aspects of physiological responses essential to growth and development [19]. MicroRNA expression profiling of sdg8 knockout mutant leaves revealed that 5 microRNAs were up-regulated more than 2-fold compared to wild type leaves (Table 1). Two of these, miR395a and

microRNA	\log_2 (sdg8/WT)	Fold change
ath-miR843	3.09	8.52
ath-miR395a	2.48	5.58
ath-miR395b	2.37	5.17
ath-miR854a	1.41	2.66
ath-miR156h	1.06	2.09
ath-miR824	-1.42	-2.68
ath-miR822	-1.32	-2.50
ath-miR391	-1.01	-2.01

Positive values indicate increased microRNA expression in the sdg8-4 knockout mutant compared to wild type (WT), whereas negative values indicate decreased microRNA expression in the sdg8-4 mutant

Table 1: Differentially expressed microRNAs in sdg8 mutant leaves identified by microarray analysis.

*Corresponding author: Jiaxu Li, Department of Biochemistry, Molecular Biology, Entomology and Plant Pathology, Mississippi State University, 32 Creelman Street, Mississippi State, MS 39762, USA, Tel: +1-662-325-1115; Fax: +1-662-325-8955; E-mail: jl305@bch.msstate.edu

Received December 23, 2013; Accepted December 26, 2013; Published December 29, 2013

Citation: Dong G, Li J (2013) MicroRNAs Profiling Reveals a Potential Link between the SDG8 Methyltransferase and Brassinosteroid-Regulated Gene Expression in *Arabidopsis*. J Data Mining Genomics Proteomics 4: e110. doi:10.4172/2153-0602.1000e110

Copyright: © 2013 Dong G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

microRNA	Target accession	Target description	Target regulation by brassinosteroid
ath-miR156h	AT3G57920	Squamosa promoter binding protein-like 15	up
ath-miR822	AT5G02330	Cysteine/Histidine-rich C1 domain family protein	up
ath-miR824	AT3G12470	Polynucleotidyltransferase	up
ath-miR843	At3g13840	GRAS family transcription factor	up
ath-miR854a	AT5G05090	Homeodomain-like superfamily protein	up
ath-miR391	AT3G02380	CONSTANS-like 2 protein	down

MicroRNA targets were predicted using psRNATarget program (<http://plantgn.noble.org/psRNATarget/>) with default parameter settings. Brassinosteroid-responsive genes were determined by the AtGenExpress database (*Arabidopsis* eFP Browser, <http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi?dataSource=Hormone>)

Table 2: Predicted microRNA targets that are brassinosteroid-responsive genes.

miR156h, have been shown to be up-regulated by 24-epibrassinolide (a highly active brassinosteroid) [20]. Taken together, these results suggest that the SDG8 methyltransferase plays a role in modulation of brassinosteroid-regulated microRNA gene expression.

Next we searched for potential target genes of the differentially expressed microRNAs in *sdg8* knockout mutant using the psRNATarget program [21]. Brassinosteroid-responsive genes were identified from the hormone response data using *Arabidopsis* eFP Browser [22]. The potential microRNA target genes being brassinosteroid-responsive are listed in Table 2. The data mining analysis reported here implies that there is a link between the SDG8 methyltransferase and brassinosteroid-regulated gene expression in *Arabidopsis*, as the SDG8 homologue in rice, SDG725, has been shown to be critical in modulating brassinosteroid-related gene expression [23].

In conclusion, our microarray data indicate that the H3K36 methyltransferase SDG8 can modulate the expression of certain microRNA genes in *Arabidopsis*. Data mining analysis suggest a link between the SDG8 methyltransferase and brassinosteroid-regulated gene expression in *Arabidopsis*.

Acknowledgment

This work was supported in part by an NIH grant (1R15GM074704-01) to J.L. We thank LC Sciences (Houston, Texas, USA) for microRNA microarray analysis.

References

- Bartel DP (2009) MicroRNAs: target recognition and regulatory functions. *Cell* 136: 215-233.
- Friedman RC, Farh KK, Burge CB, Bartel DP (2009) Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 19: 92-105.
- Jones-Rhoades MW, Bartel DP, Bartel B (2006) MicroRNAs and their regulatory roles in plants. *Annu Rev Plant Biol* 57: 19-53.
- Macfarlane LA, Murphy PR (2010) MicroRNA: Biogenesis, Function and Role in Cancer. *Curr Genomics* 11: 537-561.
- Rogers K, Chen X (2013) Biogenesis, turnover, and mode of action of plant microRNAs. *Plant Cell* 25: 2383-2399.
- Bannister AJ, Kouzarides T (2011) Regulation of chromatin by histone modifications. *Cell Res* 21: 381-395.
- Chantalat S, Depaux A, Hry P, Barral S, Thuret JY, et al. (2011) Histone H3 trimethylation at lysine 36 is associated with constitutive and facultative heterochromatin. *Genome Res* 21: 1426-1437.

- Scott GK, Mattie MD, Berger CE, Benz SC, Benz CC (2006) Rapid alteration of microRNA levels by histone deacetylase inhibition. *Cancer Res* 66: 1277-1281.
- Shin S, Lee EM, Cha HJ, Bae S, Jung JH, et al. (2009) MicroRNAs that respond to histone deacetylase inhibitor SAHA and p53 in HCT116 human colon carcinoma cells. *Int J Oncol* 35: 1343-1352.
- Bandres E, Agirre X, Bitarte N, Ramirez N, Zarate R, et al. (2009) Epigenetic regulation of microRNA expression in colorectal cancer. *Int J Cancer* 125: 2737-2743.
- Rhodes LV, Nitschke AM, Segar HC, Martin EC, Driver JL, et al. (2012) The histone deacetylase inhibitor trichostatin A alters microRNA expression profiles in apoptosis-resistant breast cancer cells. *Oncol Rep* 27: 10-16.
- Sampath D, Liu C, Vasan K, Sulda M, Puduvali VK, et al. (2012) Histone deacetylases mediate the silencing of miR-15a, miR-16, and miR-29b in chronic lymphocytic leukemia. *Blood* 119: 1162-1172.
- Mitra D, Das PM, Huynh FC, Jones FE (2011) Jumonji/ARID1 B (JARID1B) protein promotes breast tumor cell cycle progression through epigenetic repression of microRNA let-7e. *J Biol Chem* 286: 40531-40535.
- Kuchen S, Resch W, Yamane A, Kuo N, Li Z, et al. (2010) Regulation of microRNA expression and abundance during lymphopoiesis. *Immunity* 32: 828-839.
- Xu L, Zhao Z, Dong A, Soubigou-Tacconat L, Renou JP, et al. (2008) Di- and tri- but not monomethylation on histone H3 lysine 36 marks active transcription of genes involved in flowering time regulation and other processes in *Arabidopsis thaliana*. *Mol Cell Biol* 28: 1348-1360.
- Dong G, Ma DP, Li J (2008) The histone methyltransferase SDG8 regulates shoot branching in *Arabidopsis*. *Biochem Biophys Res Commun* 373: 659-664.
- Zhao Z, Yu Y, Meyer D, Wu C, Shen WH (2005) Prevention of early flowering by expression of FLOWERING LOCUS C requires methylation of histone H3 K36. *Nat Cell Biol* 7: 1256-1260.
- Ghini PE, Thorstensen T, Alm V, Vizcay-Barrena G, Windju SS, et al. (2009) The ASH1 HOMOLOG 2 (ASHH2) histone H3 methyltransferase is required for ovule and anther development in *Arabidopsis*. *PLoS One* 4: e7817.
- Clouse SD (2011) Brassinosteroid signal transduction: from receptor kinase activation to transcriptional networks regulating plant development. *Plant Cell* 23: 1219-1230.
- Lin LL, Wu CC, Huang HC, Chen HJ, Hsieh HL, et al. (2013) Identification of MicroRNA 395a in 24-Epibrassinolide-Regulated Root Growth of *Arabidopsis thaliana* Using MicroRNA Arrays. *Int J Mol Sci* 14: 14270-14286.
- Dai X, Zhao PX (2011) psRNATarget: a plant small RNA target analysis server. *Nucleic Acids Res* 39: W155-159.
- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, et al. (2007) An "Electronic Fluorescent Pictograph" browser for exploring and analyzing large-scale biological data sets. *PLoS One* 2: e718.
- Sui P, Jin J, Ye S, Mu C, Gao J, et al. (2012) H3K36 methylation is critical for brassinosteroid-regulated plant growth and development in rice. *Plant J* 70: 340-347.

Citation: Dong G, Li J (2013) MicroRNAs Profiling Reveals a Potential Link between the SDG8 Methyltransferase and Brassinosteroid-Regulated Gene Expression in *Arabidopsis*. J Data Mining Genomics Proteomics 4: e110. doi:10.4172/2153-0602.1000e110

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.editorialmanager.com/omicsgroup

