

Neural Mechanisms and Perspectives about the Therapeutic Exercises for Knee Osteoarthritis

Valdeci Carlos Dionisio^{1,2*}

¹Associate Professor, Federal University of Uberlândia, Minas Gerais, Brazil

²Master's Program in Health Sciences, Faculty of Medicine, Federal University of Uberlândia, Minas Gerais, Brazil

Osteoarthritis (OA) is the most common arthropathy in the worldwide [1], which is characterized by the degradation of articular cartilage and affects all joint structures such as the subchondral bone, fibrocartilage, ligament, capsule, and synovium [2]. The knee OA is highly prevalent in individuals of both sexes over 50 years old [3], and the high levels of pain lead the loss of functional capacity and quality of life [4].

The quadriceps weakness is a common finding in patients with OA, symptomatic or not [3]. The disuse due to pain lead for muscle atrophy, however, the neural inhibition also has been reported in individuals with OA [5], and it's has been called of arthrogenic muscle inhibition (AMI) [6]. This term describes the inability to contract a muscle completely, and is considered a reflex response to joint injury because it's beyond conscious and voluntary control, which can reduce about 60% of the quadriceps strength [7]. In the OA the AMI and quadriceps weakness would be linked to inflammation, pain, joint laxity and damaged structures [6] and therefore the IMA would be triggered by changes in the firing of sensory receptors due to damage in the knee joint [8], and changes in the excitability of spinal pathways [6], which could affect the performance of functional activities.

Although studies have reported differences in the muscle recruitment pattern and kinematic and kinetic changes in functional activities such as walking, going up and down stairs [9-11] and in simulations of muscle deficiency [12], other studies not found differences in these variables in the same conditions [13,14]. This discrepancy of results may be related to the speed of the movement in which the task is performed [9-11,13,14]. Individuals with OA tend to reduce the speed of movement because of the pain. Also, these differences can be related with subgroups of the individuals with OA [15].

The modification of the spinal pathways could be related to pain, which may be local, but also affect distant areas of the knee [16]. The pain can become chronic in patients with OA and the intense nociceptive information of the inflamed and painful structures lead to metabolic changes and reorganization of the spinal cord [17]. The term secondary hyperalgesia refers to an increased sensitivity of neurons in the spinal segments corresponding to the primary site of pain [16]. Imamura et al. [16] identified clinically hyperalgesia in disabled individuals with knee OA. Generalized hyperalgesia over superficial and deep structures suggests that central and peripheral nervous system may be involved in the maintenance of chronic pain. However, the relationship with therapeutic exercises has been few investigated.

These changes involving the nervous system and muscles could influence joint position sense, and proprioceptive acuity would be altered in patients with OA [18]. The age [19] and the presence of OA [5] affects the proprioception. The impaired proprioception may contribute to functional disability [20] and is related to weakness [21], but not with pain [18]. Hassan et al. [18] investigated the relation between pain and quadriceps function, proprioception and postural stability. Pain reduction resulted in improved function of the quadriceps, but not in proprioception and postural stability. On the other hand

Shakoor et al. [1] conducted a domiciliary program of quadriceps strengthening for eight weeks. They observed association between pain relief and muscle strength and correlation between improvement in pain and proprioception, but proprioceptive acuity was not different after training. Felson et al. [20] investigated through a longitudinal study the association between pain, proprioception and radiological OA. They found an association between proprioceptive acuity and pain, but not with the radiological findings. The authors conclude that poor proprioception by lower functional performance favors greater pain. It can be seen therefore that the relationship between pain, proprioception, and functional performance is not clear yet.

For those reasons there are several conservative treatments for knee OA. The fact that the knee OA has multifactorial causes makes it plausible that the treatment can also be multimodal [2]. However, therapeutic exercises are strongly recommended, especially quadriceps strengthening, aerobic and aquatic exercises [22]. The quadriceps strengthening appears to have greater efficacy in reducing symptoms [1]. The cause of this efficacy is the reduction of shear force and improves the distribution of forces on the knee [22], facilitating neuromuscular control [1]. There are not studies investigating the effect of the strengthening exercise in the mechanisms of peripheral and central pain, although the improvement in pain after strengthening exercise is frequently reported (assessed through questionnaires and scales). Furthermore, the characteristics of exercise, such as, how to make (uni or multi-joint exercise) and the application techniques (intensity, duration and frequency) more appropriate for treatment of knee OA is still a gap in the literature. Therefore, future research should be conducted to understand the mechanisms by which therapeutic exercise can help to reduce peripheral and central pain, in the neuromuscular control and its interaction with the other therapeutic modalities.

References

1. Shakoor N, Furmanov S, Nelson DE, Li Y, Block JA (2008) Pain and its relationship with muscle strength and proprioception in knee OA: results of an 8-week home exercise pilot study. *J Musculoskelet Neuronal Interact* 8: 35-42.
2. Miller LE, Block JE (2013) An 8-week multimodal treatment program improves symptoms of knee osteoarthritis: a real-world multicenter experience. *Pragmatic Obs Res* 4: 39-44.
3. Felson DT, Zhang Y (1998) An update on the epidemiology of knee and hip osteoarthritis with a view to prevention. *Arthritis Rheum* 41: 1343-1355.

*Corresponding author: Valdeci C Dionisio, Associate Professor, Federal University of Uberlândia, Minas Gerais, Rua Pirapuã, 70 – Itapema Sul, Uberlândia – MG – Brazil, CEP: 34.411-364, Tel/Fax: +55 (34) 3218-2910; E-mail: vc Dionisio@gmail.com

Received November 01, 2013; Accepted November 27, 2013; Published December 02, 2013

Citation: Dionisio VC (2013) Neural Mechanisms and Perspectives about the Therapeutic Exercises for Knee Osteoarthritis. *J Yoga Phys Ther* 3: 146. doi:10.4172/2157-7595.1000146

Copyright: © 2013 Dionisio VC. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

4. Picavet H, Hoeymans N (2004) Health related quality of life in multiple musculoskeletal diseases: SF-36 and EQ-5D in the DMC3 study. *Ann Rheum Dis* 63: 723-729.
5. Hurley MV, Scott DL (1998) Improvements in quadriceps sensorimotor function and disability of patients with knee osteoarthritis following a clinically practicable exercise regime. *Br J Rheumatol* 37: 1181-1187.
6. Rice DA, McNair PJ (2010) Quadriceps arthrogenic muscle inhibition: neural mechanisms and treatment perspectives. *Semin Arthritis Rheum* 40: 250-266.
7. Stevens JE, Mizner RL, Snyder-Mackler L (2003) Quadriceps strength and volitional activation before and after total knee arthroplasty for osteoarthritis. *J Orthop Res* 21: 775-779.
8. Gómez-Barrena E, Bonsfills N, Martín JG, Ballesteros-Massó R, Foruria A, et al. (2008) Insufficient recovery of neuromuscular activity around the knee after experimental anterior cruciate ligament reconstruction. *Acta Orthop* 79: 39-47.
9. Heiden TL, Lloyd DG, Ackland TR (2009) Knee joint kinematics, kinetics and muscle co-contraction in knee osteoarthritis patient gait. *Clin Biomech (Bristol, Avon)* 24: 833-841.
10. Astephen JL, Deluzio KJ, Caldwell GE, Dunbar MJ, Hubley-Kozey CL (2008) Gait and neuromuscular pattern changes are associated with differences in knee osteoarthritis severity levels. *J Biomech* 41: 868-876.
11. Liikavainio T, Bragge T, Hakkarainen M, Karjalainen PA, Arokoski JP (2010) Gait and muscle activation changes in men with knee osteoarthritis. *Knee* 17: 69-76.
12. Thompson JA, Chaudhari AM, Schmitt LC, Best TM, Siston RA (2013) Gluteus maximus and soleus compensate for simulated quadriceps atrophy and activation failure during walking. *J Biomech* 46: 2165-2172.
13. Lessi GC, da Silva Serrão PR, Gimenez AC, Gramani-Say K, Oliveira AB, et al. (2012) Male subjects with early-stage knee osteoarthritis do not present biomechanical alterations in the sagittal plane during stair descent. *Knee* 19: 387-391.
14. Hicks-Little CA, Peindl RD, Fehring TK, Odum SM, Hubbard TJ, et al. (2012) Temporal-spatial gait adaptations during stair ascent and descent in patients with knee osteoarthritis. *J Arthroplasty* 27: 1183-1189.
15. Saito I, Okada K, Nishi T, Wakasa M, Saito A, et al. (2013) Foot pressure pattern and its correlation with knee range of motion limitations for Individuals with medial knee osteoarthritis. *Arch Phys Med Rehabil* 94: 2502-2508.
16. Imamura M, Imamura ST, Kaziyama HH, Targino RA, Hsing WT, et al. (2008) Impact of nervous system hyperalgesia on pain, disability, and quality of life in patients with knee osteoarthritis: a controlled analysis. *Arthritis Rheum* 59: 1424-1431.
17. Coderre TJ, Katz J, Vaccarino AL, Melzack R (1993) Contribution of central neuroplasticity to pathological pain: review of clinical and experimental evidence. *Pain* 52: 259-285.
18. Hassan BS, Doherty SA, Mockett S, Doherty M (2002) Effect of pain reduction on postural sway, proprioception, and quadriceps strength in subjects with knee osteoarthritis. *Ann Rheum Dis* 61: 422-428.
19. Hurley MV, Rees J, Newham DJ (1998) Quadriceps function, proprioceptive acuity and functional performance in healthy young, middle-aged and elderly subjects. *Age Ageing* 27: 55-62.
20. Felson DT, Gross KD, Nevitt MC, Yang M, Lane NE, et al. (2009) The effects of impaired joint position sense on the development and progression of pain and structural damage in knee osteoarthritis. *Arthritis Rheum* 61: 1070-1076.
21. van der Esch M, Steultjens M, Harlaar J, Knol D, Lems W, et al. (2007) Joint proprioception, muscle strength, and functional ability in patients with osteoarthritis of the knee. *Arthritis Rheum* 57: 787-793.
22. Susko AM, Fitzgerald GK (2013) The pain-relieving qualities of exercise in knee osteoarthritis. *Open Access Rheumatol: Res Rev* 5: 81-91.