

Anticarcinogenic and Anti-Colitis Effects of the Japanese Vinegar Kurozu

Toru Shizuma* and Naoto Fukuyama

Department of Physiology, School of Medicine, Tokai University, Japan

Abstract

Kurozu (or Kurosu) is traditional Japanese black vinegar made by prolonged fermentation of unpolished rice and Kurozu Moromimatsu, its sediment, are both consumed in Japan as health foods or supplements.

Kurozu has established antioxidant activity that may confer the anti-inflammatory and anticarcinogenic effects reported in animal models. In this report, we summarized the studies demonstrating the anticarcinogenic and anti-colitis effects of Kurozu and Kurozu Moromimatsu.

Keywords: Kurozu or Kurosu; Kurozu Moromimatsu; Vinegar; Unpolished rice; Antioxidant activity; Anticarcinogenic effect; Anti-Colitis effect

Introduction

Kurozu (or Kurosu) is traditional Japanese black vinegar made by prolonged fermentation of unpolished rice that contains rice germ and rice bran [1]. While there are several preparation methods, one native to Kagoshima prefecture involves drying the fermented product in the sun for more than a year in earthenware jars. The liquid concentrate after drying is called Kurozu and the sediment at the bottom is called Kurozu Moromimatsu (Kurozu-M). Kurozu and Kurozu-M contain acetic acid as well as abundant organic materials, minerals, and amino acids not found in other vinegars. Kurozu-M is also rich in metabolites generated by lactobacillus and koji bacillus among other microbes. Kurozu has been reported to alleviate hyperlipidemia and hypertension [2,3]. Both Kurozu and Kurozu-M are widely consumed in Japan as health foods and supplements.

Oxidative stress is widely accepted as an important factor in the progression of inflammation and carcinogenesis. Kurozu has established antioxidant activity that may confer the anti-inflammatory and anticarcinogenic effects reported in animal models [4-6]; however, the active ingredients are unclear at present. In this report, we summarized the studies demonstrating the anticarcinogenic and anti-colitis effects of Kurozu and Kurozu-M.

Kurozu and Kurozu-M as Antioxidants

An ethyl acetate extract of Kurozu (EK) reportedly exhibits strong free radical-scavenging activity in both 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging system and linoleic acid autoxidation system [4-6]. In addition, EK suppressed lipid peroxidation in mouse skin more effectively than other vinegars and prevented dimethylbenzanthracene-induced skin carcinogenesis in mice [6]. These results suggest that the antitumor effects are partially due to the antioxidant properties of EK. Kurozu and Kurozu-M reduce nitrotyrosine levels, which is a marker of oxidative and nitrative stress [7,8]. Nitrotyrosine is produced through at least two pathways: reaction with peroxynitrite formed from superoxide (O_2^-) and nitric oxide (NO) and reaction of tyrosine with nitrite catalyzed by myeloperoxidase [9]. Black vinegar produced in Kagoshima prefecture, Izumi, has recently been reported to have potential antioxidant activity [10].

Anti-Colitis Effects of Kurozu on Experimental Animal Model

We demonstrated the protective effects of Kurozu in a rodent

model of dextran sulfate sodium (DSS)-induced colitis [8]. Oral administration of Kurozu significantly attenuated DSS-induced colitis and decreased nitrotyrosine levels in colonic epithelium, possibly by antioxidant activity. Acetic acid, the main component of Kurozu, has been reported to improve hyperlipidemia in humans and animal models [3]. However, our study indicated that acetic acid did not ameliorate DSS-induced colitis [8]. In another study, we chromatographically separated Kurozu into 4 molecular weight fractions and found active ingredients against DSS-induced colitis in rats within the 800-4,000 dalton fractions [11]. However, the specific anti-colitis factors remain to be identified.

Anticancer Effects of Kurozu *in vitro*

Nanda et al. [12] reported that EK inhibited the proliferation of several human cancer cell lines, including those derived from colon adenocarcinoma (Caco-2), lung carcinoma (A549), breast adenocarcinoma (MCF-7), bladder carcinoma (5637), and prostate carcinoma (LNCaP) [12]. Possible anticarcinogenic mechanisms in Caco-2 cells include induction of cell cycle arrest and apoptosis. This study also suggested that phenolic compounds in Kurozu are the active ingredients. In addition, Izumi reportedly inhibited the proliferation of the human squamous cell carcinoma cell line (HSC-5) *in vitro*, possibly by activating programmed necrosis (necroptosis) rather than apoptosis [13].

Therefore, apoptosis induction of Kurozu has not been unified among published studies; a possible reason being the differences in the manufacturing methods of the vinegar.

Anticancer Effect of Kurozu against Colon Cancer In Animal Models

Shimoji et al. [14] reported that EK inhibited the development of aberrant crypt foci-precursor lesions for colonic adenocarcinoma-in

*Corresponding author: Toru Shizuma, Department of Physiology, School of Medicine, Tokai University, Japan, Tel: +81-463-93-1121; Fax: +81-463-93-6684, E-mail: shizuma@is.icc.u-tokai.ac.jp

Received October 10, 2013; Accepted November 06, 2013; Published November 08, 2013

Citation: Shizuma T, Fukuyama N (2013) Anticarcinogenic and Anti-Colitis Effects of the Japanese Vinegar Kurozu. J Rice Res 1: 113. doi: [10.4172/jrr.1000113](https://doi.org/10.4172/jrr.1000113)

Copyright: © 2013 Shizuma T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the azoxymethane (AOM)-induced animal model. In this paper, EK inhibited the production of prostaglandin E2 (PGE2)-which induces hyper-proliferation-in colonic mucosa, and elevated glutathione S-transferase (GST) and quinone reductase activities in the liver [14]. These mechanisms may explain the anticarcinogenic effect of EK. Furthermore, they reported that oral administration of EK significantly inhibited the incidence and multiplicity of colon adenocarcinoma in the AOM-induced animal model [1]; thus, EK may effectively inhibit colon carcinogenesis possibly by suppressing cyclooxygenase (COX)-2.

The activation of certain matrix metalloproteinases (MMPs), particularly MMP-2 and -9, results in enhanced release of angiogenic factors into the extracellular matrix, which may promote and sustain angiogenesis and tumor growth [15]. We reported that oral administration of Kurozu-M inhibited the proliferation of the human colon adenocarcinoma cell line LoVo after subcutaneous implantation in nude mice and decreased both activated MMP-2 and -9 and nitrotyrosine levels in colonic epithelium; however, apoptosis was not induced [7]. Unlike Kurozu-M, Kurozu did not inhibit tumor growth in this study; this suggests that the antiangiogenic ingredient is contained in the sediment.

Kurozu-M Inhibition of Hepatocellular Carcinoma (HCC) In an Animal Model

We reported that oral administration of Kurozu-M significantly decreased the sizes of GST placental form-positive foci and prolonged survival in a diethylnitrosamine-induced HCC animal model [16]. HCC is typically hypervascular, and angiogenesis is considered to play an important role in its growth and progression. Therefore, one possible anti-HCC mechanism of Kurozu-M is reduced hepatic MMP-2 and -9 activities [16].

Conclusions

The anticarcinogenic and anti-colitis effects of Kurozu and Kurozu-M are demonstrated in several experimental animal models and *in vitro* studies. However, the active ingredients in the vinegar and sediment remain to be identified. At present, there is no clear evidence for anticancer or anti-colitis effects of Kurozu in humans, but clinical studies of this widely-used dietary supplement are warranted.

Acknowledgements

We would like to thank the present members of our department for their collaboration. We declare that there are no fundings.

References

1. Shimoji Y, Kohno H, Nanda K, Nishikawa Y, Ohigashi H, et al. (2004) Extract of Kurosu, a vinegar from unpolished rice, inhibits azoxymethane-induced colon carcinogenesis in male F344 rats. *Nutr Cancer* 49: 170-173.
2. Nishikawa Y, Takata Y, Nagai Y, Mori T, Kawada T, et al. (2001) Antihypertensive effect of Kurosu extract, a traditional vinegar produced from unpolished rice, in the SHR rats. *Nippon Shokuhin Kagaku Kogaku Kaishi* 48: 73-75.
3. Kondo T, Kishi M, Fushimi T, Ugajin S, Kaga T (2009) Vinegar intake reduces body weight, body fat mass, and serum triglyceride levels in obese Japanese subjects. *Biosci Biotechnol Biochem* 73: 1837-1843.
4. Murooka Y, Yamshita M (2008) Traditional healthful fermented products of Japan. *J Ind Microbiol Biotechnol* 35: 791-798.
5. Shimoji Y, Tamura Y, Nakamura Y, Nanda K, Nishidai S, et al. (2002) Isolation and identification of DPPH radical scavenging compounds in Kurosu (Japanese unpolished rice vinegar). *J Agric Food Chem* 50: 6501-6503.
6. Nishidai S, Nakamura Y, Torikai K, Yamamoto M, Ishihara N, et al. (2000) Kurosu, a traditional vinegar produced from unpolished rice, suppresses lipid peroxidation *in vitro* and in mouse skin. *Biosci Biotechnol Biochem* 64: 1909-1914.
7. Fukuyama N, Jujo S, Ito I, Shizuma T, Myojin K, et al. (2007) Kurozu moromimatsu inhibits tumor growth of LoVo cells in a mouse model *in vivo*. *Nutrition* 23: 81-86.
8. Shizuma T, Ishiwata K, Nagano M, Mori H, Fukuyama N (2011) Protective effects of Kurozu and Kurozu Moromimatsu on dextran sulfate sodium-induced experimental colitis. *Dig Dis Sci* 56: 1387-1392.
9. Radi R (2009) Peroxynitrite and reactive nitrogen species: the contribution of ABB in two decades of research. *Arch Biochem Biophys* 484: 111-113.
10. Nagashima M, Saito K (2010) Antioxidant activity of the new black vinegar "IZUMI". *J Nutr Health Aging* 14: 845-849.
11. Shizuma T, Nagano M, Fujii A, Mori H, Fukuyama N (2011) Therapeutic effects of four molecular-weight fractions of Kurozu against dextran sulfate sodium-induced experimental colitis. *Turk J Gastroenterol* 22: 376-381.
12. Nanda K, Miyoshi N, Nakamura Y, Shimoji Y, Tamura Y, et al. (2004) Extract of vinegar "Kurosu" from unpolished rice inhibits the proliferation of human cancer cells. *J Exp Clin Cancer Res* 23: 69-75.
13. Baba N, Higashi Y, Kanekura T (2013) Japanese black vinegar "Izumi" inhibits the proliferation of human squamous cell carcinoma cells via necroptosis. *Nutr Cancer* 65: 1093-1097.
14. Shimoji Y, Sugie S, Kohno H, Tanaka T, Nanda K, et al. (2003) Extract of vinegar "Kurosu" from unpolished rice inhibits the development of colonic aberrant crypt foci induced by azoxymethane. *J Exp Clin Cancer Res* 22: 591-597.
15. Roomi MW, Monterrey JC, Kalinovsky T, Rath M, Niedzwiecki A (2010) Comparative effects of EGCG, green tea and a nutrient mixture on the patterns of MMP-2 and MMP-9 expression in cancer cell lines. *Oncol Rep* 24: 747-757.
16. Shizuma T, Ishiwata K, Nagano M, Mori H, Fukuyama N (2011) Protective effects of fermented rice vinegar sediment (Kurozu moromimatsu) in a diethylnitrosamine-induced hepatocellular carcinoma animal model. *J Clin Biochem Nutr* 49: 31-35.

Citation: Shizuma T, Fukuyama N (2013) Anticarcinogenic and Anti-Colitis Effects of the Japanese Vinegar Kurozu. J Rice Res 1: 113. doi: [10.4172/jrr.1000113](https://doi.org/10.4172/jrr.1000113)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submition/>