



Coke production from heavy fractions and Algerian oil residues

Nourredine BENTAHAR

Department of the chemical and pharmaceutical processes, M'hamed Bougara University, Algeria

Abstract:

This research was performed in order to allow the study of the chemical composition influence of the coking process load on the efficiency and the quality of coke. For this reason, the coking of the following loads was realized: Atmospheric residue (RAT), vacuum Residue (RSV) and catalytic Residue of cracking (RCC). (The residues are obtained from an Algerian crude oil). As the oil residues are rich for their strongly polar composition, such as the asphaltene resins, and complex structures units (SCU), which has a role in the formation of coke, and as the dispersion of these latter improves the quality of coke, a study on the stability of aggregation was carried out by the addition of one stabilizer (oil Extract) in the coking process load. The Compounding (Extracted from /RCC oil) has been driven to the best efficiency of coke. The study consists of the influence..., this is characterized by the analyses Infra-red (IR) and x-ray diffraction (XRD).



Biography: Nourredine BENTAHAR, Department of the chemical and pharmaceutical processes, Faculty of hydrocarbons and chemistry, M'hamed Bougara University Boumerdes Avenue de l'Indépendance - Algeria

Publications:

1. Phytohormones and plant responses to salinity stress: a review
2. A biochar application protects rice pollen from high-temperature stress
3. Drought coping strategies in cotton: increased crop per drop
4. Diazotrophs-assisted phytoremediation of heavy metals: a novel approach

[World Congress on Oil and Gas Engineering, Webinar, September 28-29, 2020](#)

Abstract Citation: [Nourredine BENTAHAR, Coke production from heavy fractions and Algerian oil residues, Oil Gas 2020, World Congress on Oil and Gas Engineering, Webinar , September 28-29,2020](#)