

Editorial on Hepatitis E Virus Infection

Lene Harritshoj*

Department of Clinical Immunology, Copenhagen University Hospital, Copenhagen, Denmark

*Corresponding author: Lene Harritshoj, Department of Clinical Immunology, Copenhagen University Hospital, Copenhagen, Denmark; E-mail: lane.helm.harritshoej@regionh.dk

Received date: November 2, 2021; **Accepted date:** November 16, 2021; **Published date:** November 23, 2021

Citation: Harritshoj L (2021) Editorial on Hepatitis E virus infection. J Cytokine Biol 6: 68.

Copyright: © 2021 Harritshoj L. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Description

Hepatitis E virus infection produces a variety of extrahepatic symptoms in addition to liver symptoms. Hepatitis E virus infection has been linked to symptoms in the neurological system, kidneys, cryoglobulinemia, haematological system, reproductive system, autoimmune, and pancreas in multiple investigations. As a result, rather from being only a liver illness, Hepatitis E virus infection should be regarded a systemic disease. Distinct genotypes of Hepatitis E virus cause different extrahepatic symptoms. Even asymptomatic Hepatitis E virus infection can trigger and cause systemic disorders, and the severity of these diseases does not always correlate with the severity of Hepatitis E virus infection. Hepatitis E virus infection usually results in self-limited acute viral hepatitis, but persistent infection and extrahepatic symptoms have become a major health concern. Although genetic and serological evidence suggests that numerous other animal species may act as Hepatitis E virus hosts in industrialised countries, domestic pigs and wild boars are the principal reservoirs of Hepatitis E virus genotype 3 and genotype 4 for human infections. Hepatitis E virus infection is a rapidly spreading infection that poses a serious public health threat, particularly to immunocompromised people, pregnant women, and HBV-coinfected people. Depending on the HEV-genotype, Hepatitis E virus is transferred via faecal/oral or zoonotic transmission. Hepatitis E virus, the pathogen that causes the disease, is a little-studied but important infection. Hepatitis E virus usually produces self-limited acute viral hepatitis, but chronic infection with neurological and other extrahepatic symptoms is becoming a major clinical issue. Following the discovery of swine Hepatitis E virus in pigs and proof of its zoonotic potential, Hepatitis E virus strains from more than a dozen different animal species were genetically identified. Hepatitis E virus

strains from pigs, rabbits, deer, camels, and rats have been demonstrated to infect humans and overcome species boundaries. Abnormal liver function tests are prevalent during pregnancy. While hepatic damage during pregnancy usually has a minor impact on mother and foetal outcomes, it can occasionally result in serious maternal and foetal morbidity, and even death. We look at the epidemiology, clinical features, diagnosis, and treatment of hepatitis during pregnancy caused by less common pathogens like Epstein-Barr virus, cytomegalovirus, herpes simplex viruses, dengue fever, malaria, leptospirosis, Q fever, typhoid fever, and other uncommon infections, as well as the implications for infants' breastfeeding. Infection with the hepatitis E virus can induce persistent infections in solid organ transplant recipients, as well as liver damage and cirrhosis. Because infection is typically asymptomatic, under diagnosis is a big issue. Systematic screening may allow for early detection and treatment of liver disease. Infection with the hepatitis E virus is a serious public health issue. In India, Hepatitis E virus infection has been recognised as a leading source of enterically transmitted acute sporadic hepatitis, particularly among adults. Hepatitis E virus is hyperendemic in India, where outbreaks and instances of acute sporadic viral hepatitis are both common. The majority of these outbreaks can be linked to human faeces pollution of drinking water systems. In the recent decade, our understanding of the virus's epidemiology, clinical features, diagnostic procedures, therapeutic choices, and the necessity for immunisation has changed dramatically. In a group of immunocompromised patients, the prevalence of ongoing, chronic, and previous hepatitis E virus infections was studied. The link between transfusion-transmitted Hepatitis E virus and Hepatitis E virus seroprevalence was investigated, and the Hepatitis E virus seroprevalence was compared to that of healthy blood donors.