

# Eighty Six Years Duration of Type 1 Diabetes Mellitus

Joseph Fovenyi\*

Department of Diabetes Outpatient, Péterfy Sándor Hospital-Clinic and Manninger Jenő National Institute for Traumatology, Budapest, Hungary

\*Correspondence to: Joseph Fovenyi, Department of Diabetes Outpatient, Péterfy Sándor Hospital-Clinic and Manninger Jenő National Institute for Traumatology, Budapest, Hungary, Tel: 36302979686; E-Mail: j.fovenyi@gmail.com

Received date: August 03, 2021; Accepted date: August 17, 2021; Published date: August 24, 2021

Citation: Fovenyi J (2021) Eighty Six Years Duration of Type 1 Diabetes Mellitus. J Clin Diabetes, Vol.5 Iss.5 No:121.

Copyright: © 2021 Fovenyi J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

The patient was born in 1927 to a wealthy middle class family from Buda. Before she was born, her maternal uncle died of type 1 diabetes, probably of diabetic ketoacidosis at the age of 22. Her family members were not long-lived. She was diagnosed with diabetes in 1932 at the age of five. Her condition was immediately treated with rapid-acting ("regular") insulin, which was administered three times a day. In the mid-thirties, her father purchased a Zeiss polarimeter and the urine collected in three fractions was polarized three times daily and the dose of insulin was adjusted to the urine glucose levels obtained. In the early forties, they switched to a once daily mixture of protamine zinc and regular insulin. Her subsequent insulin therapy will be discussed below. In the thirties until the late forties, she maintained the usual ketogenic diet at the time with up to 60-70 grams of carbohydrates.

**Keywords:** Diabetes; Mellitus; Polarimeter

## Introduction

As for self-testing, she switched from polarimetric glucose self-monitoring until the mid-sixties to the then available urinary glucose assay using Diastix strips. She started using a home blood glucose self-monitoring in 1982 until her death [1].

She lived like other healthy children except for her diabetes, did several sports, such as cycling, horse riding, tennis, volleyball, swimming, and skating. She maintained regular daily exercise for the rest of her life. She had a lean body. Subcutaneous fat was almost completely absent, except for the one palm-sized area on the anterior surface of the thighs where insulin was always administered [2].

She married in 1952. Her husband adored her and basically devoted his life to her. Before his death in the early 2000s, he said he had been sleeping half-awake for decades holding his wife's hand watching out for the signs of hypoglycemia, which had occurred almost every night due to a then daily overdose of insulin. The patient was so used to being pampered since childhood that she has always insisted on her - often unrealistic - ideas, even in the face of her caring doctor [3].

The author was in his third postgraduate year as a doctor at St. John's Hospital, Budapest, II. Department of Internal Medicine, when he first met the patient at the end of 1962. She immediately selected him as her treating and caring physician and this relationship was retained for more than 55 years until the patient's death. She was admitted to St. John's Hospital once in 1967 during 15 years. She then injected a total of 60 units of insulin daily, half as rapid-acting and half as protamine zinc insulin. She later switched to crystalline pork, then to human zinc insulin. At that time no diabetic complications were detected in the patient. From 1978 until 2001 her treatment continued in the "B" Department of Internal Medicine at the Budapest Péterfy Sándor Hospital. Following the retirement of the author, she was treated at the Diabetes Outpatient Department. She was hospitalized

once at Péterfy Hospital in 1988. Two C-peptide tests were performed at that time with negative results [4].

She was the first in Hungary to receive the Joslin 75-Year Medal and award for her achievement in living with diabetes without serious diabetic complications for more than 75 years, which she was very proud of [5].

In 2012, she underwent bilateral cataract surgery, but her vision continued to deteriorate due to the progression of age-related dry macular degeneration. (No signs of diabetic retinopathy were observed during the eye examinations.) The deterioration of her vision later made the self-management of her diabetes more and more difficult.

It is very important to discuss her insulin therapy in more detail. Except for the last two and a half months of her life, she has not given permission to the use of any form of advanced insulin treatment. The reason for the consent of her doctor was that the 24-hour duration of action was clear for the once-daily administration of premixed rapid-acting plus NPH insulin which replaced crystalline zinc insulin. However, she checked her blood glucose 5-10 times a day, and corrected each jump with 1-2 units of insulin lispro injection. Her daily dose of insulin has been around 22 units for two decades.

This brings us to her dietary issues. When her fasting blood glucose was around 8-10 mmol/l, she would delay her breakfast up to 11 o'clock checking her glucose every hour. She rarely ate cooked food, she consumed most of her daily 150 g of carbohydrates in the form of buns, eating very little meat, vegetables and fruits. So, besides her absolute non-physiological and out-of-date insulin therapy, her diet was extremely deficient, and certainly not healthy.

Nonetheless, all her HbA1c values since 1993 have been around 7% (53 mmol/mol), sometimes dropping below 6% (42 mmol/mol). Meanwhile, her impaired vision and worsening neuropathy, which has not improved with alpha lipoic acid and pregabalin administration, as her only diabetic complication caused severe lower extremity

complaints and made the self-management of her diabetes more difficult.

Serious problems with her treatment/care developed in the fall of 2017. The reason for this was that although she measured her blood glucose values 8-10 times a day (she did not feel the finger prick pain due to neuropathy), her earlier stable blood glucose fluctuations of 4-10 mmol/l started to fluctuate strongly. This was due to the fact that her eyesight was impaired, so she had repeatedly made errors in her blood glucose measurement, and also often had her pen needles bent at 90 degrees, which she had not noticed either and did not actually inject insulin. This has worsened until 2018 to such an extent that - due to severe hypoglycemia - she had to be hospitalized 7 times in 3 months in the II. Department of Internal Medicine of our hospital. So she spent a total of 57 days in hospital in the spring before she died

Her ECG was normal. Her laboratory parameters measured at this time were: RBC: 3,4 million, Hb: 102 g/l, WBC: 7,400, Creatinine: 64 µmol/l, eGFR: 79 ml/min, Total cholesterol: 4.2 mmol/l, HDL cholesterol: 1.5 mmol/l, Triglycerides: 0.8 mmol/l. CRP: 57, HbA1c: 6.8% (49 mmol/mol), Urinary glucose excretion: 42 mg/24 hours. Thus, apart from anemia and inflammation, no other differences suggestive of diabetic complications could be detected.

In May, the patient returned home and agreed to 24-hour care at her apartment following very strong persuasion. With the presence of a nurse trained in blood glucose control, insulin administration and proper diet, the patient's condition began to improve rapidly, her appetite increased and she exercised regularly on a daily basis. Her blood sugar levels stabilized at 5-7 mmol/l and the hypoglycemic episodes were abolished with degludec and glulisine insulin given for two months. Her final insulin doses proved to be extremely low, with 8 units of degludec and 3x2 units of glulisine insulin.

She sent away the 24-hour nurse a month later, saying that if she was able to look after herself for 86 years, she would continue to be able to do so. Her high-level intellect has proved to be completely wrong this time. After the nurse left, she spent more and more time in bed with hypostatic pneumonia and bronchitis, refused hospital admission, collapsed at home and died 23 days later, two months after she was 91 years old.

Myocardial infarction localized to the interventricular septum was reported as the main cause of death, dilated cardiomyopathy, hypostatic pneumonia, bronchitis, and acute pulmonary edema were reported as additional findings during autopsy. The calcareous plaques at some places reduced the coronary arteries to a pinpoint lumen. Extreme levels of arteriosclerosis have been found throughout her arterial system.

## Conclusion

The life expectancy of type 1 diabetics is lower than that of their non-diabetic counterparts, but this has dramatically improved over the past century, primarily due to the discovery and introduction of insulin. Nevertheless some individuals who had diabetes since childhood may have a longer duration of diabetes and thus may live for longer, although the number of these individuals is not high. On the basis of available data, we cannot find an explanation for the very long diabetes duration.

Although there is no example of longevity in the patient's family, a special constellation of the genes may be responsible. Another important fact is the patient's resilience, namely her extremely positive outlook on life, her life-affirming attitude, and above all, her tremendous pride in herself and her performance. If we add that in the last 55 years of patient's life she has been cared for by a single, attentive and helpful doctor it weighs heavily, but only in addition to what was mentioned above.

Summary of the "clinical pearls" exemplified by this case

- Extreme long duration of Type 1 diabetes
- No advanced insulin treatment during 86 years
- Great blood glucose fluctuations during the first 60 years of the disease
- No severe late complications except neuropathy
- No exact explanation for the patient's longevity
- Patient was treated/cared by a single doctor during the last 55 years

## References

1. Miller RG, Secrest AM, Sharma RK, Songer TJ, Orchard TJ, et al. (2012) Improvements in the Life Expectancy of Type 1 Diabetes. The Pittsburgh Epidemiology of Diabetes Complications Study Cohort. *Diabetes*. 61:2987-2992.
2. Mameli C, Mazzantini S, Ben Nasr M, Fiorina P, Scaramuzza AE, et al. (2015) Explaining the increased mortality in type 1 diabetes *World J Diabetes* 6: 889-895.
3. Sebastiani P, Solovieff N (2012) Genetic Signatures of Exceptional Longevity in Humans. *Plos one*; .
4. Borch-Johnsen K, Kreiner S, Deckert T (1986) Mortality of type 1 (insulin-dependent) diabetes mellitus in Denmark: a study of relative mortality in 2930 Danish type 1 diabetic patients diagnosed from 1933 to 1972. *Diabetologia* 29:767-772.
5. Sartor G, Nystrom L, Dahlquist G (1991) The Swedish Childhood Diabetes Study: a seven-fold decrease in short-term mortality? *Diabet Med* 8:18-21.