

## Fertility Sparing Options in Gynecologic Oncology

Fabio Martinelli<sup>\*</sup>, Antonino Ditto, Giorgio Bogani, Mauro Signorelli and Francesco Raspagliesi

Department of Gynecologic Oncology, IRCCS National Cancer Institute, Milan, Italy

<sup>\*</sup>Corresponding author: Fabio Martinelli, Department of Gynecologic Oncology, IRCCS National Cancer Institute, Milan, Italy, Tel: 00390223903322; E-mail: [fabio.martinelli@istitutotumori.mi.it](mailto:fabio.martinelli@istitutotumori.mi.it)

Received: Dec 18, 2015; Accepted: Dec 21, 2015; Published: Dec 28, 2015

Copyright: © 2015, Martinelli F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Introduction

Traditionally, and according to actual guidelines, [1] cancers affecting female genital organs are treated with radical surgery, chemotherapy, radiation therapy or a combination of the above mentioned, all bearing a permanent damage of the female reproductive tract. Anyway, facing with a continuous trend in postponing childbearing and the early detection of gynecologic cancers, there has been a vast increase in the number of patients seeking fertility preserving options for the treatment of gynecologic malignancies [2]. Of the 6.6 million cancers occurring worldwide in female population, 1.09 million (16%) affects the female genital organs and up to 20% will be diagnosed in women of reproductive age. [3,4] As many as 15-45% of cervical cancers, 5-29% of endometrial cancers, and 12-34% of primary ovarian malignancies will be found in women eligible for fertility preservation. It is not surprising so the increase interest in oncofertility discipline. Even if not yet considered as "standard treatments", there are continuously emerging data on fertility sparing options available in gynecologic oncology. In the present paper we briefly review current options for women desiring fertility-sparing treatment of gynecologic malignancies [5].

### Cervical Cancer

For women with stage IA1 cervical cancer (non-visible lesion invading < 3 mm), coldknife-conization, loop-electrosurgical-excision-procedure (LEEP), or CO2-conization are all reasonable options. Recurrence after these procedures are <1% assuming negative margins are achievable.

For stages IA2 (non-visible lesion invading 3-5 mm) or IB1 (non-visible lesion invading > 5 mm or any visible lesion < 4 cm in size) disease, options vary depending on tumor size, histology, depth of invasion, and the presence or absence of lymph-vascular space invasion. Radical trachelectomy (via open, laparoscopic, robotically-assisted or vaginal approaches) plus pelvic lymphnode dissection is a largely accepted treatment for young women with early cervical cancer desiring fertility-sparing surgery. Typically this procedure is offered to women with squamous or adenocarcinomas less than 2 cm in size, however many centers are undertaking larger lesions with successful outcomes. The 4% recurrence rate after radical trachelectomy is seemingly equivalent to recurrence after radical hysterectomy and pregnancy rates of 41-79% have been reported. There is a slight increase in second trimester miscarriage but the majority of pregnant women deliver after 34 weeks. To overcome the increased risk of pre-term birth and second trimester miscarriage, less radical procedures such as cervical conization or simple trachelectomy with pelvic lymphadenectomy or neoadjuvant-chemotherapy followed by cervical conization and lymphadenectomy are also being explored for stages IA2/IB1 disease. Even if largely limited to small series, comparable

results in terms of oncological and obstetrical outcome seem achievable [6-8].

### Endometrial Cancer

Conservative therapy using oral/injectable progesterone therapy or a progesterone eluting IUD for uterine cancer should be limited to grade 1 endometrioid adenocarcinomas without myometrial invasion. Although evaluating myometrial invasion is difficult, pretherapeutic MRI/ Transvaginal ultrasound are often obtained to rule out significant invasion. Exclusion of adnexal masses is mandatory being the risk of adnexal involvement or presence of synchronous tumors up to 25%. In assessing outcomes, the published literature is largely composed of small case series with reported response rates of 75-80% and recurrence rates of 20-40%. Pregnancy rates as high as 35% have been achieved. Pretreatment dilatation and curettage ± hysteroscopy for accurate assessment of cell histology and grade is a necessity as well as close treatment monitoring with frequent endometrial sampling [9-11].

### Ovarian Cancer

Ovarian cancer is a heterogeneous group encompassing various types of diseases with different histologies, treatments and prognosis.

Due to their exquisite sensitivity to chemotherapy, germ-cell tumors of the ovary are highly amenable to fertility sparing therapy. Even when gross tumor on the bilateral ovaries and uterus are encountered, performance of a unilateral salpingo-oophorectomy (largely for diagnostic purposes) with retention of the diseased contralateral ovary and uterus followed by chemotherapy is a widely accepted practice. Overall germ-cell tumors have a cure rate of 90-95%, and a pregnancy rate of 35%.

Borderline tumors of the ovary as well as grossly apparent stage I sex-cord stromal tumors can also be managed with a unilateral salpingo-oophorectomy only or even cystectomy if feasible. Data over 2500 borderline tumor patients treated conservatively are available with a pregnancy rate as high as 55%.

For invasive epithelial ovarian cancer grossly limited to one ovary, conservative staging sparing one ovary and uterus ± adjuvant chemotherapy is an option in well consented patients. To date, more than 800 cases of early stage epithelial ovarian cancer managed with unilateral salpingo-oophorectomy have been reported with a recurrence rate ranging from 8 to 17% and 36% of patients achieving pregnancy. Apart patients with FIGO stage IA G1-2 for whom fertility sparing treatment is largely accepted; only few data for higher stages or aggressive histologies are available. Anyway comparable oncologic and obstetrical results seem achievable [12-17].

## Oncofertility

Major exciting advances have been made in reproductive-endocrinology and infertility with respect to embryo, oocyte, and even ovarian tissue preservation in young women with a diagnosis of cancer at any site (not just gynecologic malignancies). Those new technologies extremely appealing especially when dealing with malignancies other than that of the female genital tract can find a place even in a gynecologic oncologists' practice. It is well known that endometrial cancer patients as well as patients suffering from borderline tumors frequently have a history of infertility. To date ovarian stimulation after conservative treatment for cervical, endometrial and even borderline ovarian cancer is considered acceptable and is under investigation in other gynecological malignancies [18].

## Conclusions

In the last 20 years, gynecologic oncologists have developed a wide variety of fertility-sparing therapies for young women with gynecologic malignancies. These non-standard treatments are best administered and managed in referral centers and on-study when appropriate. Careful patient selection and extensive discussion and counseling of risks of these approaches are of the utmost importance. Ideally, a multidisciplinary team of gynecologic oncologists, psychologists, reproductive-endocrinologists and maternal-fetal-medicine doctors will partake in the care of these women. Ongoing prospective protocols will hopefully clarify risks and benefits of conservative treatment in gynecologic cancers.

## References

1. Infectious Diseases Committee and Reproductive Endocrinology and Infertility Committee of Society of Obstetricians and Gynecologists of Canada; Canadian HIV Pregnancy Planning Guideline Development Team Core Working Group, Loutfy MR, Margolese S, Money DM, Gysler M, et al. (2012) Canadian HIV Pregnancy Planning Guidelines: No. 278, June 2012. *Int J Gynaecol Obstet* 119: 89-99.
2. Hamilton BE, Martin JA, Osterman MJ, Curtin SC (2015) Births: Preliminary Data for 2014. *Natl Vital Stat Rep* 64: 1-19.
3. Siegel RL, Miller KD, Jemal A (2015) Cancer statistics. *CA Cancer J Clin* 65: 5-29.
4. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J (2015) Global cancer statistics, 2012. *CA Cancer J Clin* 65: 87-108.
5. Eskander RN, Randall LM, Berman ML, Tewari KS, Disaia PJ, et al. (2011) Fertility preserving options in patients with gynecologic malignancies. *Am J Obstet Gynecol* 205: 103-110.
6. Ditto A, Martinelli F, Bogani G, Fischetti M, Di Donato V (2015) Fertility-sparing surgery in early-stage cervical cancer patients: oncologic and reproductive outcomes. *Int J Gynecol Cancer* 25: 493-497.
7. Pareja R, Rendon GJ, Sanz-Lomana CM, Monzon O, Ramirez PT (2013) Surgical, oncological, and obstetrical outcomes after abdominal radical trachelectomy - a systematic literature review. *Gynecol Oncol* 131: 77-82.
8. Pareja R, Rendón GJ, Vasquez M, Echeverri L, Sanz-Lomana CM, et al. (2015) Immediate radical trachelectomy versus neoadjuvant chemotherapy followed by conservative surgery for patients with stage IB1 cervical cancer with tumors 2cm or larger: A literature review and analysis of oncological and obstetrical outcomes. *Gynecol Oncol* 137: 574-580.
9. Gunderson CC, Fader AN, Carson KA, Bristow RE (2012) Oncologic and reproductive outcomes with progestin therapy in women with endometrial hyperplasia and grade 1 adenocarcinoma: a systematic review. *Gynecol Oncol* 125: 477-482.
10. Gallos ID, Yap J, Rajkhowa M, Luesley DM, Coomarasamy A, et al. (2012) Regression, relapse, and live birth rates with fertility-sparing therapy for endometrial cancer and atypical complex endometrial hyperplasia: a systematic review and metaanalysis. *Am J Obstet Gynecol* 207: e1-12.
11. Koskas M, Yazbeck C, Walker F, Delorme P, Azria E, et al. (2012) [Fertility sparing management of endometrial adenocarcinoma and atypical hyperplasia: a literature review]. *Bull Cancer* 99: 51-60.
12. Vasconcelos I, de Sousa Mendes M (2015) Conservative surgery in ovarian borderline tumours: a meta-analysis with emphasis on recurrence risk. *Eur J Cancer* 51: 620-631.
13. Solheim O, Tropé CG, Rokkones E, Kærn J, Paulsen T, et al. (2015) Fertility and gonadal function after adjuvant therapy in women diagnosed with a malignant ovarian germ cell tumor (MOGCT) during the "cisplatin era". *Gynecol Oncol* 136: 224-229.
14. Satoh T, Hatae M, Watanabe Y, Yaegashi N, Ishiko O, et al. (2010) Outcomes of fertility-sparing surgery for stage I epithelial ovarian cancer: a proposal for patient selection. See comment in PubMed Commons below *J Clin Oncol* 28: 1727-1732.
15. Ditto A, Martinelli F, Bogani G, Lorusso D, Carcangiu M, et al. (2015) Long-term safety of fertility sparing surgery in early stage ovarian cancer: comparison to standard radical surgical procedures. *Gynecol Oncol* 138: 78-82.
16. Ditto A, Martinelli F, Lorusso D, Haeusler E, Carcangiu M, et al. (2014) Fertility sparing surgery in early stage epithelial ovarian cancer. *J Gynecol Oncol* 25: 320-327.
17. Somigliana E, Peccatori FA, Filippi F, Martinelli F, Raspagliesi F, et al. (2014) Risk of thrombosis in women with malignancies undergoing ovarian stimulation for fertility preservation. *Hum Reprod Update* 20: 944-951.
18. Dursun P, Doan NU, Ayhan A (2014) Oncofertility for gynecologic and non-gynecologic cancers: fertility sparing in young women of reproductive age. *Crit Rev Oncol Hematol* 92: 258-267.