

## Is the Indication of Liver Resection for Hepatocellular Carcinoma Expanding with the Application of Laparoscopic Approach?

Norihiro Kawabe\*, Zenichi Morise, Masashi Isetani, Satoshi Arakawa, Jin Kawase, Hidetoshi Nagata and Hirokazu Tomishige

Department of Surgery, Fujita Health University School of Medicine, Japan

\*Corresponding author: Kawabe Norihiro, Department of Surgery, Fujita Health University School of Medicine, Banbuntane Houtokukai Hospital, 3-6-10, Otobashi Nakagawa-ku, Nagoya, Aichi 454-8509, Japan, Tel: +81-52-321-8171; Fax: +81-52-323-4502; E-mail: zmorise@aol.com

Received date: July 18, 2014, Accepted date: Aug 26, 2014, Published date: Sep 03, 2014

Copyright: © 2014 Kawabe N, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

Liver resection (LR) for the patients of hepatocellular carcinoma, often with chronic liver disease, have high risks of developing significant postoperative complications and multicentric metachronous repeat lesions with the need of repeat treatments. Reduction of surgery-induced parenchymal injury and destruction of the collateral blood/lymphatic flow, which leads to less post-operative ascites production, and facilitation of repeat LR with less adhesion and improved vision/manipulation between adhesions are among the advantages of laparoscopic LR. These characteristics of laparoscopic LR may lead to expanding indication of LR.

**Keywords:** Liver resection; Laparoscopic hepatectomy; Hepatocellular carcinoma; Chronic liver disease; Repeat hepatectomy

### Introduction

Hepatocellular carcinoma (HCC) is the fifth most common primary cancer and the third most common cause of cancer-related deaths worldwide [1,2]. The treatment options for HCC include liver resection (LR) [3], liver transplantation [4], transarterial chemoembolization (TACE), and local ablation therapy [5]. Among them, LR and liver transplantation are thought to be the best hopes for cure. Although liver transplantation should be adapted to patients with deteriorating liver function and HCC within the Milan criteria [6], LR should be considered as a primary therapy in the other patients [7,8]. However, most patients with HCC have underlying chronic liver disease (CLD). Therefore, there are high risks of developing significant postoperative complications and also multicentric metachronous repeat lesions occurring in chronically injured liver after LR with the need of repeat treatments. When considering the treatment of HCC in patients with CLD, the degree of invasive surgical stress, especially to the impaired liver, should be considered in addition to the oncological therapeutic effects. Patients with severe CLD have various (overt and preliminary) symptoms, such as: (1) deteriorations of protein synthesis and metabolism; (2) GI tract congestion, ascites, pancytopenia due to portal hypertension and hypersplenism; and (3) susceptibility to infectious diseases and hepatopulmonary syndrome (hypoxaemia) due to increased shunt vessels [9]. Cirrhotic patients have high morbidity and mortality following anesthesia and surgery [10] and the risk of abdominal operations increases according to the preoperative Child-Pugh class of the patients [11]. Even limited small volume LR for severe cirrhotic patients often develops refractory ascites, which leads to fatal complications [12,13].

Since the first successful report in 1992 [14], laparoscopic LR is thought to be a less invasive procedure than conventional open LR for the treatment of hepatic lesions [15]. In a comprehensive meta-analysis study, laparoscopic LR was compared to open in 1678 patients across 26 studies [16]. While it is associated with longer operating times and no differences in oncological outcomes, it is advantageous in

several aspects. Laparoscopic LR is associated with reduced blood loss, decreased portal clamp time, decreases in overall and liver-specific complications, and shorter post-operative hospital stays. Furthermore, recent technological development of devices and accumulation of experiences have facilitated the expansion of the indication of the procedure [17,18]. For those patients with HCC and CLD who are not able to undergo usual treatment modalities due to liver function, tumor size and/or localization, "less invasive" pure laparoscopic hepatectomy may provide a good option, especially after repeat treatments for the disease with multicentric metachronous lesions occurring in chronically injured liver.

Under open surgery, even limited LR for severe CLD patients often develops refractory ascites leading to fatal complications [12,13]. Laparoscopic LR may be particularly advantageous for those patients, given the potential for lower levels of parietal and hepatic injury and the preservation of venous and lymphatic collateral circulation in cirrhotic patients. The safety and feasibility of the laparoscopic approach and its short-term benefits for HCC patients with CLD have been demonstrated by several series. The studies have investigated the major differences between laparoscopic and open LR [19-29]. Favorable short-term results, including fewer incidences of ascites and liver failure, and shorter postoperative hospital stay correlate with the laparoscopic procedure. Tranchart et al. reported laparoscopic LR of HCC for selected patients resulted in better postoperative outcomes without long-and short-term oncologic consequences [30]. Also in our experience [31], we reported the favorable perioperative course of the patients with severe CLD (Child B/C and ICG R15 of 40% or above) who underwent pure laparoscopic LR, which is comparable to that of the patients with mild/moderate CLD (Child A and ICG R15 of 10.1-27.4%). This study showed that laparoscopic LR has the advantage of minimal ascites [31] due to preservation of venous and lymphatic collateral circulation, which leads to lower risk of disturbance in water and/or electrolyte balance and hypoproteinemia that could trigger fatal liver failure. This feature could be the most remarkable specific advantage for laparoscopic LR. Patients who undergo LR are exposed three different types of stresses: 1) general, whole-body surgical stress, 2) reduced liver function due to resected liver volume, 3) surgery-

induced injuries for liver parenchyma and environment around the liver, such as parenchymal injury by compression of the liver and destruction of the collateral blood/lymphatic flow by laparotomy plus mobilization of the liver. Reduction of the third mentioned stress by laparoscopic LR should lower the risk for HCC patients with severe CLD. Among these patients with severe CLD in our series, one underwent living-related liver transplantation 20 months after hepatectomy [32]. The procedure could also be an advantageous option in bridging therapy to liver transplantation for certain HCC patients with severe CLD. Furthermore, our experience showed laparoscopic LR also results in improved vision and manipulation in a small operative field under several conditions, such as repeat hepatectomy with adhesions [32]. In addition to smaller extend of adhesion after laparoscopic than open LR, this feature of laparoscopic LR facilitates the repeat LR for the patients with multicentric metachronous lesions occurring in chronically injured liver. Also,

some novel approach of LR (for example, caudal, not anterior as open, approach for oncologically appropriate posterior sectorectomy and right hepatectomy [33]) is also possible laparoscopically, using the laparoscope-specific view/approach (such as good view from caudal and dorsal sides of the liver) and by changing position of the patients for the use of the gravity to dissection and handling large-volumed liver/tumors.

These characteristics of laparoscopic LR listed above may indicate it is a superior method when compared to open LR under certain conditions and its application may lead to expanding indication of LR. Table 1 is summary of recent reports of comparative studies of laparoscopic LR and open LR for HCC, which show favorable results for laparoscopic LR. However, there is no controlled study comparing laparoscopic LR and open LR, liver transplantation, ablation therapies etc. to date. Further investigations are needed.

Study		Ascites		Liver failure		Hospital stay (days, mean ± SD)		Mortality	
Author	Year Ref No.	LLR	OLR	LLR	OLR	LLR	OLR	LLR	OLR
Lai	2009 [34]	NA	NA	NA	NA	NA	NA	0/25	1/33
Aldrighetti	2010 [35]	0/16	1/16	NA	NA	6.3 ± 1.7	9 ± 3.8	0/16	0/16
Tranchart	2010 [30]	3/42	11/42	0/36	4/53	6.7 ± 5.9	9.6 ± 3.4	1/42	1/42
Kim	2011 [36]	0/26	1/29	NA	NA	11.08 ± 4.96	16.07 ± 10.697	0/26	0/29
Lee	2011 [37]	0/33	2/50	NA	NA	NA	NA	0/33	0/50
Truant	2011 [38]	5/36	12/53	NA	NA	6.5 ± 2.7	9.5 ± 4.8	0/36	4/53
Ker	2011 [27]	2/116	26/208	NA	NA	6.2 ± 3	12.4 ± 6.8	0/116	6/208

**Table 1:** Recent reports of laparoscopic (LLR) and open liver resection (OLR) comparative studies for HCC

## References

- El-Serag HB, Rudolph KL (2007) Hepatocellular carcinoma: epidemiology and molecular carcinogenesis. *Gastroenterology* 132: 2557-2576.
- Schütte K, Bornschein J, Malfertheiner P (2009) Hepatocellular carcinoma--epidemiological trends and risk factors. *Dig Dis* 27: 80-92.
- Capussotti L, Ferrero A, Viganò L, Polastri R, Tabone M (2009) Liver resection for HCC with cirrhosis: surgical perspectives out of EASL/AASLD guidelines. *Eur J Surg Oncol* 35: 11-15.
- Hwang S, Lee SG, Belghiti J (2010) Liver transplantation for HCC: its role: Eastern and Western perspectives. *J Hepatobiliary Pancreat Sci* 17: 443-448.
- Lau WY, Leung TW, Yu SC, Ho SK (2003) Percutaneous local ablative therapy for hepatocellular carcinoma: a review and look into the future. *Ann Surg* 237: 171-179.
- Mazzafferro V, Regalia E, Doci R, Andreola S, Pulvirenti A, et al. (1996) Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 334: 693-699.
- Ryder SD (2003) British Society of Gastroenterology Guidelines for the diagnosis and treatment of hepatocellular carcinoma (HCC) in adults. *Gut* 52 Suppl 3: iii1-8.
- Cha CH, Ruq L, Fong Y, Jarnagin WR, Shia J, et al. (2003) Resection of hepatocellular carcinoma in patients otherwise eligible for transplantation. *Ann Surg* 238: 315-321.
- Hoeper MM, Krowka MJ, Strassburg CP (2004) Portopulmonary hypertension and hepatopulmonary syndrome. *Lancet* 363: 1461-1468.
- Ziser A, Plevak DJ, Wiesner RH, Rakela J, Offord KP, et al. (1999) Morbidity and mortality in cirrhotic patients undergoing anesthesia and surgery. *Anesthesiology* 90: 42-53.
- Mansour A, Watson W, Shayani V, Pickleman J (1997) Abdominal operations in patients with cirrhosis: still a major surgical challenge. *Surgery* 122: 730-735.
- Belghiti J, Hiramatsu K, Benoit S, Massault P, Sauvanet A, et al. (2000) Seven hundred forty-seven hepatectomies in the 1990s: an update to evaluate the actual risk of liver resection. *J Am Coll Surg* 191: 38-46.
- Lai EC, Fan ST, Lo CM, Chu KM, Liu CL, et al. (1995) Hepatic resection for hepatocellular carcinoma. An audit of 343 patients. *Ann Surg* 221: 291-298.
- Gagner M, Rheault M, Dubuc J. (1992) Laparoscopic partial hepatectomy for liver tumor. *Surg Endosc* 6: 97-98
- Kaneko H, Tsuchiya M, Otsuka Y, Yajima S, Minagawa T, et al. (2009) Laparoscopic hepatectomy for hepatocellular carcinoma in cirrhotic patients. *J Hepatobiliary Pancreat Surg* 16: 433-438.
- Mirnezami R, Mirnezami AH, Chandrakumaran K, Abu Hilal M, Pearce NW, et al. (2011) Short- and long-term outcomes after laparoscopic and open hepatic resection: systematic review and meta-analysis. *HPB (Oxford)* 13: 295-308.
- Buell JF, Cherqui D, Geller DA, O'Rourke N, Iannitti D, et al. (2009) The international position on laparoscopic liver surgery: The Louisville Statement, 2008. *Ann Surg* 250: 825-830.
- Tsuchiya M, Otsuka Y, Tamura A, Nitta H, Sasaki A, et al. (2009) Status of endoscopic liver surgery in Japan: a questionnaire survey conducted by the Japanese Endoscopic Liver Surgery Study Group. *J Hepatobiliary Pancreat Surg* 16: 405-409.

19. Morise Z. (2012) Pure laparoscopic hepatectomy for HCC patients. In: Lau WY. *Hepatocellular Carcinoma - Clinical Research*. Zagreb, Croatia: InTech, 183-196.
20. Chen HY, Juan CC, Ker CG (2008) Laparoscopic liver surgery for patients with hepatocellular carcinoma. *Ann Surg Oncol* 15: 800-806.
21. Dagher I, Lainas P, Carloni A, Caillard C, Champault A, et al. (2008) Laparoscopic liver resection for hepatocellular carcinoma. *Surg Endosc* 22: 372-378.
22. Kaneko H, Takagi S, Otsuka Y, Tsuchiya M, Tamura A, et al. (2005) Laparoscopic liver resection of hepatocellular carcinoma. *Am J Surg* 189: 190-194.
23. Cherqui D, Laurent A, Tayar C, Chang S, Van Nhieu JT, et al. (2006) Laparoscopic liver resection for peripheral hepatocellular carcinoma in patients with chronic liver disease: midterm results and perspectives. *Ann Surg* 243: 499-506.
24. Shimada M, Hashizume M, Maehara S, Tsujita E, Rikimaru T, et al. (2001) Laparoscopic hepatectomy for hepatocellular carcinoma. *Surg Endosc* 15: 541-544.
25. Sarpel U, Hefti MM, Wisniewsky JP, Roayaie S, Schwartz ME, et al. (2009) Outcome for patients treated with laparoscopic versus open resection of hepatocellular carcinoma: case-matched analysis. *Ann Surg Oncol* 16: 1572-1577.
26. Hu BS, Chen K, Tan HM, Ding XM, Tan JW (2011) Comparison of laparoscopic vs open liver lobectomy (segmentectomy) for hepatocellular carcinoma. *World J Gastroenterol* 17: 4725-4728.
27. Ker CG, Chen JS, Kuo KK, Chuang SC, Wang SJ, et al. (2011) Liver Surgery for Hepatocellular Carcinoma: Laparoscopic versus Open Approach. *Int J Hepatol* 2011: 596792.
28. Endo Y, Ohta M, Sasaki A, Kai S, Eguchi H, et al. (2009) A comparative study of the long-term outcomes after laparoscopy-assisted and open left lateral hepatectomy for hepatocellular carcinoma. *Surg Laparosc Endosc Percutan Tech* 19: e171-174.
29. Nguyen KT, Marsh JW, Tsung A, Steel JJ, Gamblin TC, et al. (2011) Comparative benefits of laparoscopic vs open hepatic resection: a critical appraisal. *Arch Surg* 146: 348-356.
30. Tranchart H, Di Giuro G, Lainas P, Roudie J, Agostini H, et al. (2010) Laparoscopic resection for hepatocellular carcinoma: a matched-pair comparative study. *Surg Endosc* 24: 1170-1176.
31. Morise Z, Sugioka A, Kawabe N, Umemoto S, Nagata H, et al. (2011) Pure laparoscopic hepatectomy for hepatocellular carcinoma patients with severe liver cirrhosis. *Asian J Endosc Surg* 4: 143-146.
32. Morise Z, Kawabe N, Kawase J, Tomishige H, Nagata H, et al. (2013) Pure laparoscopic hepatectomy for hepatocellular carcinoma with chronic liver disease. *World J Hepatol* 5: 487-495.
33. Tomishige H, Morise Z, Kawabe N, Nagata H, Ohshima H, et al. (2013) Caudal approach to pure laparoscopic posterior sectionectomy under the laparoscopy-specific view. *World J Gastrointest Surg* 5: 173-177.
34. Lai EC, Tang CN, Ha JP, Li MK (2009) Laparoscopic liver resection for hepatocellular carcinoma: ten-year experience in a single center. *Arch Surg* 144: 143-147.
35. Aldrighetti L, Guzzetti E, Pulitanò C, Cipriani F, Catena M, et al. (2010) Case-matched analysis of totally laparoscopic versus open liver resection for HCC: short and middle term results. *J Surg Oncol* 102: 82-86.
36. Kim HH, Park EK, Seoung JS, Hur YH, Koh YS, et al. (2011) Liver resection for hepatocellular carcinoma: case-matched analysis of laparoscopic versus open resection. *J Korean Surg Soc* 80: 412-419.
37. Lee KF, Chong CN, Wong J, Cheung YS, Wong J, et al. (2011) Long-term results of laparoscopic hepatectomy versus open hepatectomy for hepatocellular carcinoma: a case-matched analysis. *World J Surg* 35: 2268-2274.
38. Truant S, Bouras AF, Hebbar M, Boleslawski E, Fromont G, et al. (2011) Laparoscopic resection vs. open liver resection for peripheral hepatocellular carcinoma in patients with chronic liver disease: a case-matched study. *Surg Endosc* 25: 3668-3677.