



Journal of Infectious Disease and Pathology

Akira Ito

Department of Parasitology, Asahikawa Medical University, Asahikawa 078-8510, Japan

Taeniasis caused by two species, the pork tapeworm, *Taenia solium* and the beef tapeworm, *Taenia saginata*, and 'Asian *Taenia*', *Taenia saginata asiatica*, are briefly overviewed.

Among these tapeworms, *T. solium* is unique and has serious public health importance, since it causes cysticercosis, especially neurocysticercosis, the most potentially lethal helminthic disease in humans, when tapeworm carriers and others get egg ingestion. Cysticercosis is crucially different from taeniasis, since the former is transmitted from human (tapeworm carriers) to humans including the tapeworm carriers themselves. These tapeworm infections, especially *T. solium* and *T. saginata asiatica* are exclusively endemic in remote areas where local people defecate out-doors and pigs have free access to human feces. Pigs are contaminated with eggs of these tapeworms in human feces and harbor huge number of the larval stage, cysticerci of *T. solium*.

This In contrast, although pig is only confirmed domesticated animal harboring cysticerci of *T. saginata asiatica* in the liver, the cysticerci have never been fully developed. The endemic areas are remote or rural areas in developing countries in Asia where the owners of these domesticated animals sacrifice them in the backyard of their houses and serve uncooked or undercooked meat and viscera without meat inspections at all. The endemic areas for both *T. solium* and *T. saginata asiatica* are often crucially different from *T. saginata*. It is affected from the religion. In Muslim societies, *T. solium* taeniasis is not endemic. In non-Muslim societies, *T. solium* is more common than *T. saginata*. Even though taeniasis are endemic in remote areas, CC including NCC is becoming cosmopolitan distribution. Such crucial difference between taeniasis and CC in Asia is overviewed.

Urinary tract Infections (UTI) speak to 35–40% of each nosocomial illness and have the greatest idea in the general HAIs arrangement. The prevalence of crisis facility acquired UTIs is constantly extending comprehensive. The market is most likely going to create at a CAGR of 9% during the measurement time period. SSI area is creating at a reliable rate and is presumably going to create at an equivalent pace during the figure time period

The prevalence of therapeutic centre acquired pneumonia (HAP) and related passings are growing at an upsetting rate far and wide. Thusly, there is an extended emphasis on constraining the occasion of HAP in restorative administrations settings.

The market is depended upon to watch fast advancement in the coming an exceptionally prolonged stretch of time in light of the improvement and commercialization of bleeding-edge

- All submissions must be in English
- The topic must fit into systematic sessions of the conference
- Each discrete participant is allowed to submit maximum 2 abstracts
- Abstract must be succumbed online as per the given abstract template
- Abstracts must be printed in Times New Roman and font size will be 12
- Abstract must contain title, name, relationship, country, speakers biography, recent picture, image and reference

To receive the award, the awardee must submit the demonstration for which the award is given, for publication at the website, along with author permission. Failure to submit the PPT, and permission within the selected timeframe will result in forfeiture of award.

