

A New Candidate of Catalase Appears during the Germination after Priming in Naturally Aged Neem (*Azadirachta indica*) Seeds

Vimal Pandey* and Atanu Kumar Pati

School of Life Sciences, Pt. Ravishankar Shukla University, Raipur 492010, Chhattisgarh, India

Abstract

Loss of Neem (*Azadirachta indica*) seeds viability remarkably occurs with natural storages and related to decreased seed germination. We investigated that osmopriming induces hydrogen peroxide during germination that could be signaling molecule for germination improvement. A new catalase isozyme appeared during post-priming germination. The results show a high coordination between hydrogen peroxide and catalase and H₂O₂ may play a role in regulation of new catalase isozyme expression during post-priming germination of aged neem seeds.

Keywords: Seed priming; Germination; Catalase; Hydrogen peroxide; Isozyme

Introduction

Ageing accompanies diverse deleterious changes in cells that accumulate over the time that eventually leads to death. Seeds too lose viability with the progress of time. Seed viability is a complicated trait under the control of genetic, developmental, and environmental factors [1,2]. Often these factors interact with each other and produce deleterious effects during the process of seed maturation and harvest. Seed moisture and storage temperature play a significant role and confer varying length of seed viability. Concisely, we come across wide variability in seed viability, within and among seed lots and species that attributed to the interactions of above mentioned factors.

In recent years, increasing evidence have been accumulated suggesting a functional significance for reactive oxygen species in seed aging and germination [3,4]. Reactive Oxygen Species (ROS) are metabolites of molecular oxygen (O₂) that have higher reactivity than O₂. ROS can include unstable oxygen radicals, such as superoxide radical (O₂^{•-}) and hydroxyl radical (OH[•]), and non-radical molecules, like hydrogen peroxide (H₂O₂). Thus, all organisms living in an aerobic environment are exposed to ROS on a continual basis. Up to 2% of the oxygen utilized is converted into these deleterious agents and related reactive ROS, mostly, but not exclusively, within mitochondria. An enzymatic dismutation step must firstly take place to produce more stable H₂O₂ derivative from the O₂^{•-}. H₂O₂ often travel across the cell membrane and is required for a viable long-range cell-to-cell signal [5]. Additional reactive OH[•] can be formed from O₂^{•-} and H₂O₂ through Fe-catalyzed Haber-Weiss reaction that may cause lipid peroxidation [6]. The concentration of ROS is tightly controlled by ROS-scavenging pathways that metabolize ROS. However, an imbalance in generation and metabolism of ROS leads to a variety of physiological challenges by disrupting redox homeostasis of the cell, which is collectively known as "oxidative stress". Plant possesses an efficient antioxidative defense that protects the cell from oxidative damage. To minimize and/or to protect against the toxic effects of these damaging ROS, cells have evolved highly regulated enzymatic and non-enzymatic mechanisms to keep a balance between ROS production and destruction in order to maintain cellular redox homeostasis. These protective systems are composed of low molecular and enzymatic scavengers, such as superoxide dismutase (SOD), catalase (CAT) and enzymes of the Halliwell Asada pathway: ascorbate peroxidase (APX), dehydro ascorbate reductase (DHAR) and glutathione reductase (GR), connected with antioxidant compounds: ascorbate and glutathione [7]. SOD is a key enzyme in the regulation of

the amount of superoxide radicals and peroxides. The removal of H₂O₂ through CAT as well as a series of reactions is known as an ascorbate glutathione cycle in which ascorbate and glutathione participate in a cyclic transfer of reducing equivalents resulting in the reduction of H₂O₂ to H₂O using electrons derived from nicotinamide adenine dinucleotide phosphate (NADPH) [8]. The cooperative activities of these enzymes and high levels of the low molecular antioxidants may increase the resistance to oxidative injury and minimize cell damage.

Additionally, increased level of ROS, including H₂O₂ [9,10], O₂^{•-} [11,12], and hydroxyl radicals [10], as well as reactive nitrogen species have been reported to play positive role during seed germination in plant species of varied taxonomic backgrounds. Seed priming affects germination rate, seed vigor, and seedling development. It has been shown that seed priming with water or solutions of various substances positively affects traits of seeds and seedlings [13,14]. Many research groups evaluated the ROS and detoxifying enzyme activity during seed priming and germination [15,16]. Bailly et al. proposed that an oxidative window for germination restricts the occurrence of cellular germination related events to a defined range of ROS level [17].

Previously research found that Neem (*Azadirachta indica*) seeds lose their viability during natural ageing [18,19]. The seed priming improved germination to a limited extent [20]. The aims of this work are to examine H₂O₂ metabolism during ageing and its post-priming germination.

Materials and Methods

Seed collection and extraction

Mature, ripe yellow fruits of *Azadirachta indica* (Neem) were collected manually from the trees growing on the campus of Pt. Ravishankar Shukla University, Raipur, (21°14'14" N latitude and

*Corresponding author: Vimal Pandey, National Institute of Plant Genome Research, JNU Campus, P.Box No-10531, Aruna Asaf Ali Marg-110067, New Delhi, India, Tel: +91-9990566447; E-mail: vimal.bt04@gmail.com

Received August 27, 2016; Accepted September 16, 2016; Published September 20, 2016

Citation: Pandey V, Pati AK (2016) A New Candidate of Catalase Appears during the Germination after Priming in Naturally Aged Neem (*Azadirachta indica*) Seeds. J Plant Biochem Physiol 4: 171. doi:10.4172/2329-9029.1000171

Copyright: © 2016 Pandey V, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

81° 38'55" E longitude), and brought to the laboratory immediately. The initial water content of seeds was determined using randomly chosen mechanically depulped seeds. Seeds with endocarp were extracted from the fruit tissue by soaking them in 20% H₂SO₄ (Merck, India) for 30 min following depulping by rubbing over a wire mesh and washing repeatedly under tap water. Seeds with hard endocarp were dried to their initial water content under shade for storage [21]. Seeds were harvested from hard endocarp by braking gently to analyze biochemical variables after every 15 days following storage.

Sample collection and priming treatment

The seeds were collected at different percentage of germination and denoted as G_{100%}, G_{90%}, G_{70%}, G_{50%}, G_{30%} and G_{0%}. Seeds with 100% germination referred as a control. The osmopriming treatments of naturally aged seeds were carried out in 12-cm-diameter petri dish with single layers of filter paper moistened with 5 ml of -0.78 MPa PEG solutions for germination improvement as optimized for neem seed priming [20]. Ten seeds of each differentially aged (G_{90%} to G_{0%}) in five replicates were placed for treatment. After one day priming at 30°C, the seeds were washed in sterile water for twice, blotted on filter paper and immediately used for germination tests and other analysis. After each stage of ageing and treatment, following sets: dried (NA-WI), primed (Primed-NA-WI), germinated (NA-RE) and post-priming germinated (Primed-NA-RE) seeds were immediately frozen in liquid nitrogen to stop the chemical reaction, and grounded to a fine powder with a pestle and mortar in LN₂. The powder was stored at -80°C in humidity-proof vials until use.

Hydrogen peroxide estimation

Batches of five seeds at differentially aged and treated were pre-incubated for 30 min in 3 ml of K-phosphate buffer (20 mM, pH 6.0) to remove pre-formed H₂O₂ and were then incubated in 3 ml of the same buffer containing 5 μm scopoletin and 3 μg ml⁻¹ horseradish peroxidase (Boehringer Mannheim) in darkness at 25°C on a shaker. The estimation of H₂O₂ was performed according to Schopfer et al. by an increase in fluorescence (excitation: 346 nm, emission: 455 nm) in the incubation medium and was measured using reagent blanks as a reference [10]. Results corresponded to the mean ± SE of four replicates and are expressed in fluorescence, 455 nm (relative unit) per seed.

Catalase (CAT) enzyme analysis and assay

Stored LN₂ crushed differentially aged and treated seed tissue samples were weighed (Fifty mg embryonic axes), grounded again in liquid nitrogen and homogenized in cold borate buffer (0.2 M, pH 7.4) containing 4% poly-vinylpyrrolidone (PVP-40 Himedia), followed by centrifugation at 17,890 g for 20 min. The supernatants were stored at -20°C for later determination of enzyme activity spectrophotometrically.

Catalase (EC 1.11.1.6) activity was estimated measuring the decomposition of H₂O₂ at 240 nm as per the method of Aebi [22]. Enzyme activity was performed by taking, in a quartz cuvette, 2.74 ml of potassium phosphate buffer (37.5 mM, pH 6.8), to which 60 μl enzyme extract was added and the absorbance was zeroed at 240 nm. Catalase activity was triggered by adding 200 μl of H₂O₂ (60 mM). The change in absorbance was recorded for 6 min at 15 sec intervals. The blank was without enzyme extract. Catalase activity was expressed in A₂₄₀ min⁻¹mg⁻¹ protein.

CAT Isozyme

For CAT isozyme assays, stored LN₂ crushed differentially aged and

treated seed axes samples were homogenized in ice chilled Zivy's buffer (containing 30 mM Tris-base pH 8.5, 0.25 mM ascorbic acid, 1mM EDTA, 5 mM MgCl₂, 1 mM DTT and 0.1 mM PMSF) and centrifuged at 10,000 rpm for 10 min at 4°C [23].

CAT Isozymes were separated on 7% non-denaturing poly acrylamide gel electrophoresis and 30 μg protein per well was loaded. Gels were run at 20 mA constant current till the dye front reaches the bottom of the gel. Gels were stained by the method followed by Woodbury et al. [24]. After completion of gel running, gel was soaked in DDW for 15 min and incubated in 0.03% H₂O₂ for 10 min and stained with a solution of 1% ferric chloride and 1% Potassium ferricyanide. Gel turns Prussian blue except the position exhibiting CAT activity.

The protein content of the extract was determined by Bradford method using bovine serum albumin as standard [25].

Statistical analysis

Data were analyzed by one-way ANOVA in combination with Duncan's multiple-range tests at 5% level of significance (p ≤ 0.05) for post-hoc comparisons of means. Statistical tests were carried out using SPSS (version 16) for Microsoft Windows (SPSS, INC., Chicago, IL).

Results and Discussion

Seed ageing/ loss of viability addresses stochastic passive degenerative processes that may be initiated by external factors but are endogenously influenced by an uncontrollable oxidative stress. ROS accumulation induces oxidative stress when the imbalance between ROS production and antioxidant defense against these ROS takes place. The findings of the present study show that hydrogen peroxide (H₂O₂) increased with its priming (primed-WI), germination (RE) and post-priming germination (primed-RE) compared with dried/ without imbibed (WI) by about 16.68, 28.85 and 39.24% at each level of ageing (Table 1). Many earlier studies have highlighted the increase in oxidative stress during seed ageing [26-28]. The increased H₂O₂ level in seed in the current experiment observed after priming (Primed-WI), germination (RE) and post-priming germination (primed-RE) compared to dried/ without imbibed (WI), suggest that free-radical mediated processes were initiated by re-aeration of hypoxic tissues and/or higher respiration. Ageing induced a decrease in CAT activity in differentially aged seeds (Table 2). Catalase activity increased with priming, germination as well as post-priming germination in embryonic axes at each level of ageing, but such improved activity turned down with ageing. Interestingly, CAT activity was enhanced on average 35.39% after their priming (Primed-WI), 34.55% during germination (RE) and 64.82% post-priming germination (primed-RE) as compared to dried/without imbibed (WI) embryo axes during natural ageing. Gels stained for CAT activity revealed only one CAT isoform in naturally aged embryonic axes; however, new isoform appeared during post-priming germination. The band intensity gradually decreased with different level of ageing, whereas, the intensity increased after its priming as well as radicle emergence at each level of ageing (Figure 1).

Conclusion

Hydrogen peroxide enhanced during radicle emergence (RE), priming (primed-WI) and priming subsequent radicle emergence (primed-RE) compared with the dried/ without imbibed (WI) aged seeds. It is suggested that free-radical mediated processes were initiated by re-aeration of hypoxic tissues and/or higher respiration. Catalase enzyme significantly decreased with ageing CAT activity was enhanced with osmopriming and/or radicle emergence. The occurrence of extra

	Without Imbibed		Radicle Emergence	
	Aged	Primed	Aged	Primed
Natural Ageing				
G _{100%}	80.13 ± 1.23	..§	95.61 ± 2.21	..§
G _{90%}	82.50 ± 2.14	95.44 ± 2.77	113.03 ± 3.38	127.43 ± 2.69
G _{70%}	90.03 ± 3.56	110.05 ± 3.33	131.26 ± 3.31	149.90 ± 5.41
G _{50%}	102.56 ± 5.34	135.23 ± 3.32	156.33 ± 6.07	184.78 ± 4.80
G _{30%}	132.60 ± 2.73	167.20 ± 5.49	171.34 ± 5.02	211.42 ± 4.16
G _{0%}	185.02 ± 6.82	198.60 ± 3.61	-	-
F-value (1)	97.368	121.115	53.372	71.798
Df	5, 18	4, 15	4, 15	3, 12
p-value	<0.001	<0.001	<0.001	<0.001

§ Control (G_{100%}) excluded in osmopriming treatment

Table 1: Hydrogen peroxide (H₂O₂) released by aged and primed seeds at two different stages, without imbibed (WI) and radicle emergence (RE) during different levels of natural ageing. Values were represented in relative fluorescence mean at 455nm ± SE and data were expressed in per seed basis.

	Without Imbibed		Radicle Emergence	
	Aged	Primed	Aged	Primed
Axes				
G _{100%}	1.25 ± 0.03	..§	1.36 ± 0.02	..§
G _{90%}	0.72 ± 0.04	1.21 ± 0.03	1.23 ± 0.01	2.23 ± 0.05
G _{70%}	0.66 ± 0.05	1.08 ± 0.08	1.09 ± 0.01	2.17 ± 0.09
G _{50%}	0.57 ± 0.03	0.99 ± 0.02	0.96 ± 0.04	1.54 ± 0.05
G _{30%}	0.52 ± 0.02	0.85 ± 0.02	0.93 ± 0.02	1.28 ± 0.07
G _{0%}	0.37 ± 0.01	0.44 ± 0.02	-	-
F-value (1)	97.887	49.832	51.554	49.256
Df	5, 18	4, 15	4, 15	3, 12
p-value	<0.001	<0.001	<0.001	<0.001

§ Control (G_{100%}) excluded in osmopriming treatment

Table 2: Catalase (CAT) activity in axes tissue of aged and primed seeds at two different stages, without imbibed (WI) and radicle emergence (RE) during different levels of natural ageing. Values were presented in activity mean ± SE and expressed in A₂₄₀ min⁻¹ mg⁻¹ protein.




Figure 1: Native gel stain for embryonic-axes CAT of aged and primed seeds during without imbibed (WI) and radicle emergence (RE) at different level of natural ageing. Arrows indicate number of CAT isoform appeared on native gel.

CAT isoform during germination in osmoprimed seeds confirmed that it was produced by a separate CAT gene and probably expresses during germination.

References

- Betty M, Finch-Savage WE, King GJ, Campos F, Covarrubias AA (2000) Quantitative genetic analysis of seed viability and pre-emergence seedling growth traits in *Brassica oleracea*. New Phytol. 148: 277-286.
- Pukacka S, Ratajczak E (2007) Age-related biochemical changes during storage of beech (*Fagussylvatica* L.) seeds. Seed Sci Res 17: 45-53.
- Bailly C (2004) Active oxygen species and antioxidants in seed biology. Seed Sci Res 14: 93-107.
- McDonald MB (1999) Seed deterioration: physiology, repair and assessment. Seed Sci Technol 27: 177-237.
- Allan C, Fluhr R (1997) Two distinct sources of elicited reactive oxygen species in tobacco epidermal cells. Plant Cell 9:1559-1572.
- Dat J, Breusegerm VF, Vandenebee S, Vranova E, Van MM, et al. (2000) Active oxygen species and Catalase during plant stress response. Cell Mol Life Sci 57: 779-786.
- Asada K (1992) Ascorbate peroxidase- a hydrogen peroxide scavenging enzyme in plants. Physiol Plant 85: 235-41.
- Goel A, Sheoran IS (2003) Lipid peroxidation and peroxide- scavenging enzymes in cotton seed under natural ageing. Biol Plant 46: 429-434.
- Bailly C, Bogatek-Leszczynska R, Come D, Corbineau F (2002) Changes in activities of antioxidant enzymes and lipoxygenase during growth of sunflower seedlings from seeds of different vigour. Seed Sci Res 12: 47-55.
- Schopfer P, Plachy C, Frahy G (2001) Release of reactive oxygen intermediates

- (superoxide radicals, hydrogen peroxide, and hydroxyl radicals) and peroxidase in germinating radish seeds controlled by light, gibberellin, and abscisic acid. Plant Physiol 125: 1591-1602.
11. Gidrol X, Lin WS, Degousee N, Yip SF, Kush A (1994) Accumulation of reactive oxygen species and oxidation of cytokinin in germinating soybean seeds. Eur J Biochem 224: 21-28.
 12. Liu X, Xing D, Li L, Zhang L (2007) Rapid determination of seed vigor based on the level of superoxide generation during early imbibition. Photochem Photobiol Sci 6: 767-774.
 13. Bradford KJ (1986) Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. Hort Sci 21: 1105-1112.
 14. McDonald MB (2000) Seed priming. In: Seed Technology and its Biological Basis (Eds. M Black, JD Bewley), Sheffield Academic Press, Sheffield, England. 287-325.
 15. Goel A, Goel AK, Sheoran IS (2003) Changes in oxidative stress enzymes during artificial ageing in cotton (*Gossypiumhirsutum* L.) seeds. J Plant Physiol 160: 1093-1100.
 16. Hsu CC, Chen CL, Chen JJ, Sung JM (2003) Accelerated aging enhanced lipid peroxidation in bitter melon seeds and effects of priming and hot water soaking treatments. Sci Hortic 98: 201-212.
 17. Bailly C, El-Maarouf-Bouteau H, Corbineau F (2008) From intracellular signaling networks to cell death: the dual role of reactive oxygen species in seed physiology. C R Biol 331: 806-814.
 18. Sacande M, Hoekstra FA, Van Pijlen JG, Groot SPC (1998) A multifactorial study of conditions influencing neem (*Azadirachta indica*) seed storage longevity. Seed Sci Res 8: 473-478.
 19. Sacande M, Golovina EA, Van Aelst AC, Hoekstra FA (2001) Viability loss of neem (*Azadirachta indica*) seeds associated with membrane phase behaviour. J Exp Bot 92: 919-931.
 20. Pandey V (2014) Oxidative stress mediated ageing in non-orthodox seeds: involvement of ROS, scavenging enzymes, telomere and telomerase. Shodhganga.
 21. Sacandé M, BuitinkJ, Hoekstra FA (2000) A study of water relations in neem (*Azadirachta indica*) seed that is characterized by complex storage behavior. J Exp Bot 51: 635-643.
 22. Aebi HE (1983) Catalase, In: HU Bergmeyer, eds, Methods of Enzymatic Analysis. VerlageChemie, Weinheim, Germany. 123-456.
 23. Zivy M, Thiellement H, Vienne D, Hofmann JP (1983) Study on nuclear and cytoplasmic genomic expression in wheat by two-dimensional gel electrophoresis. Theo Appl Genet 66: 1-7.
 24. Woodbury W, Spencer AK, Stahmann MA (1971) An improved procedure using ferricyanide for detecting catalase isozymes. Anal Biochem 44: 301-305.
 25. Bradford MM (1997) A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein Dye Binding. Anal Biochem 72: 248-254.
 26. Bailly C, Benamar A, Corbineau F, Come D (1996) Changes in malondialdehyde content and in superoxide dismutase, catalase and glutathione reductase activities in sunflower seeds as related to deterioration during accelerated aging. Physiol Plant 97: 104-110.
 27. Kibinza S, Vinel D, Come D, Bailly C, Corbineau F (2006) Sunflower seed deterioration as related to moisture content during ageing, energy metabolism and active oxygen species scavenging. Physiol Plant 128: 496-506.
 28. Kranner I, Birtić S, Anderson KM, Pritchard HW (2006) Glutathione half-cell reduction potential: A universal stress marker and modulator of programmed cell death. Free Radic Biol Med 40: 2155-2165.