

A New Homozygous ABCB4 Mutation Identified in Two Chinese Siblings Based on Exome Sequencing

Hui Lin¹, Rong Fu², Xiumin Zhang¹, Bihui Yao³, Jing Ye¹, Jianguo Shi¹, Wen Sui⁴ and Zengshan Li^{1*}

¹Department of Pathology, Xijing Hospital and Basic Medical College, Fourth Military Medical University, Xi'an 710032, Shaanxi, P.R. China

²Department of Pediatrics, Xijing Hospital, Fourth Military Medical University, Xi'an 710032, Shaanxi, P.R. China

³Department of Surgery, Baotou Central Hospital, Baotou 014040, Inner Mongolia, P.R. China

⁴Suzhou University Children's Hospital, Suzhou 215003, Jiangsu, P.R. China

Abstract

Background: Many pediatric cholestatic liver diseases show unspecific symptoms, laboratory tests and histological features. A complete genetic background check is necessary to identify the genetic determinants as well as the diagnosis and differential diagnosis.

Method: The whole genome exome sequencing was performed in two Chinese sister siblings with unspecific cholestasis and the data was analyzed.

Results: A homozygous mutation, c.2176C-T transition (p.P726L) in exon 17 of ABCB4, was identified. Further study indicated that there are mutant alleles in the genome of their non consanguineous parents. The final diagnosis is progressive familial intrahepatic cholestasis (PFIC), type 3.

Conclusion: As a rare hereditary cholestatic liver disease, PFIC shares many similar features with other hereditary or acquired liver disease and requires a wide range of differential diagnosis.

Exome sequencing is a useful tool for mapping this kind of monogenic disease mutation and plays a very important role in genetic counseling and prenatal diagnosis.

Keywords: ABCB4; Mutation; Progressive familial intrahepatic cholestasis; Whole-exome sequencing

Abbreviations: PFIC- Progressive Familial Intrahepatic Cholestasis; MDR3 - Multidrug Resistance P-Glycoprotein 3; PC- Phosphatidyl choline; TM- Trans Membrane Segments

Introduction

Cholestasis is a common manifestation of hepatic disease in pediatric patients and a wide range of differential diagnosis, especially inherited diseases, need to be considered. Moreover, many of these entities have overlapped symptoms, histological findings as well as laboratory features. Under this kind of situations, genetic background assay is very important for a precise final diagnosis. There are different ways in which DNA sequencing can be undertaken, such as sanger sequencing, resequencing chip and denaturing high-performance liquid chromatography [1,2]. Anyway, the scope of these methods is limited sometimes, especially in detecting the unidentified genes. Whole-exome sequencing, as a new technique, is considered to be unbiased — it does not focus on specific genes or loci deemed of interest on the basis of assumptions about possible causal associations. Recent studies have shown that exome sequencing is a powerful, efficient and cost-effective strategy to identify the responsible genes for monogenic diseases such as Miller syndrome and familial combined hypolipidemia [3,4]. In this study we utilized exome sequencing in blood samples of two Chinese siblings who presented with cholestasis phenotypes to identify the underlying genetic defects responsible for the disease and confirm the clinical diagnosis. Finally a diagnosis of family progressive familial intrahepatic cholestasis was made and a new homozygous ABCB4 mutation was identified as well.

Progressive Familial Intrahepatic Cholestasis (PFIC) refers to a heterogeneous group of autosomal recessive liver disorders of childhood in which cholestasis of hepatocellular origin often presents in the neonatal period or first year of life and leads to death from liver failure

at ages usually ranging from infancy to adolescence [5]. The estimated incidence of PFIC is about one per 50000-100000 births [6]. PFIC can be classified into three types with slightly different clinical, biochemical and histological features, associated with mutations in three distinct hepatobiliary transport proteins encoded by related ATP-Binding Cassette (ABC) genes, ATP8B1 (PFIC1), ABCB11 (PFIC2) and ABCB4 (PFIC3) [7]. It's also possible that other unidentified genes involved in bile formation may be responsible for the PFIC phenotypes because mutations have not yet been identified in very rare PFIC patients [6]. Therefore molecular analysis is often recommended to confirm the diagnosis and typing of PFIC in affected patients. Histologically, PFIC exhibits extensive bile ductular proliferation, intracanalicular cholestasis, portal inflammation, and fibrosis. All these findings are nonspecific and differential diagnosis should be considered in any neonate or infant with progressive symptomatic cholestasis.

Material and Methods

Patient samples

In this study, two Chinese siblings presented with pruritus, jaundice and hepatomegaly in 9 and 15 months old separately. They

***Corresponding author:** Zengshan Li, Department of Pathology, Xijing Hospital and Basic Medical College, Fourth Military Medical University, Xi'an 710032, Shaanxi, P.R. China, Tel: 86-29-84771576; Fax: 86-29-83255697; E-mail: lizsh72@fmmu.edu.cn

Received July 28, 2014; **Accepted** September 23, 2014; **Published** September 29, 2014

Citation: Lin H, Fu R, Zhang X, Yao B, Ye J, et al. (2014) A New Homozygous ABCB4 Mutation Identified in Two Chinese Siblings Based on Exome Sequencing. J Genet Syndr Gene Ther 5: 243. doi:[10.4172/2157-7412.1000243](https://doi.org/10.4172/2157-7412.1000243)

Copyright: © 2014 Lin H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

both have similar laboratory findings, including elevated alanine aminotransferase, aspartate aminotransferase, bile salt, alkaline phosphatase and gamma-glutamyl transpeptidase. The older sister once received mother-to-daughter living donor heterotopic auxiliary part liver transplantation at the age of 2. The postoperative liver biopsy showed extensive intracanalicular cholestasis, bile ductular proliferation, portal inflammation and fibrosis. All these findings are nonspecific and the final diagnosis is unclear. Six years later her younger sibling sister was born and presented with similar symptoms in 9 months old. Their parents, non-consanguineous and healthy, are both Chinese Hui ethnic minorities (Muslim ethnic group). For further diagnosis and classification, the blood samples of the siblings were collected for exome sequencing.

DNA preparation and exome capture sequencing

Genomic DNA was extracted from venous blood with a commercially available kit, Omega Blood DNA Midi Kit (D3494-01), using the manufacturer's protocols. Qualified genomic DNA samples from each case were randomly fragmented by Covaris and enriched using Agilent's Sure Select Biotinylated RNA Library (BAITS). The array covers 1.22% of human genomic regions corresponding to the NCBI consensus coding sequence database 2008. Each captured library was then loaded on HiSeq2000 platform, and sequencing was performed for each captured library independently to ensure each sample had at least 50-fold coverage. Raw image files were processed by Illumina Pipeline v1.6 for base-calling with default parameters and the sequences of each individual were generated as 90bp paired-end reads.

Bioinformatics analysis

For the basic analysis, low quality sequences in the raw Solexa reads were filtered out as follows: base quality is more than 20, depth is between 4 and 200, estimate copy number is equal or less than 2 and the distance between two SNPs must be longer than 4. Coverage and depth of each sample were also calculated.

For the advanced analysis, the sequencing reads were aligned to the Human reference genome (NCBI 36.3) using SOA Paligner (soap2.20) and the non-synonymous variants resulting in changes of proteins were extracted as candidate variants. Based on the hypothesis that the pathogenic mutation underlying the rare familiar disease was not present in general population, these variants were then filtered against several public database, comprising dbSNP129, eight previously exome-sequenced Hap Map samples (Hap Map 8), the SNP release of the 1000 Genome Project (20100208 release) and Yan Huang (YH) 1 genome database, to remove the common variants. Then we choose the homozygous variants shared by both patients as the target variants. In addition, we used SIFT (Sorting Intolerant from Tolerant), available at <http://sift.jcvi.org/>, to predict that whether the target variants would have a phenotypic effect. Sanger sequencing using customized primers was performed to determine the presence of the target variants in the parents. Sequencing and bioinformatics analysis were performed in Beijing Genomics Institute (Beijing, China).

Because inherited syndromes of intrahepatic cholestasis commonly resulted from mutations in the genes including SERPINA1, JAG1, ATP8B1, ABCB11 and ABCB4, we simultaneously focused on these genes to investigate possible variants.

Ethics approval

Xijing Hospital Human Research Ethics Committee

Results

Exome sequencing and data analysis

For each of the two cases, we generated approximately 2.3 gigabases of effective sequence data allowing each targeted base to be read approximately 60 times. About 8000 synonymous variants were found in each case, of which approximately 7000 were missense mutations and approximately 40 were nonsense mutations (Table 1). We also detected more than 8000 insertions and deletions in each case and most of them were intergenic variants.

Given that non-synonymous variants were more likely to be the pathogenic mutations and unlikely to be present in general group, so we filtered them against several public database mentioned in methods and the number of the candidate variants was reduced to about 400. Considering that PFIC is an autosomal recessive inherited disorder and the same clinical phenotypes have been found in two siblings, so the target variants should be homozygous and shared by patients. There were only four variants meeting all the conditions. SIFT prediction showed that only one of the variants, the c.2176C-T transition (p.P726L) of ABCB4, occurred in a highly conserved position and was more likely to be functionally damaging (Table 1). The same mutation has been revealed compound heterozygous in the study of another group [8]. So we believed that this variant would be the very causation responsible for the two patients' sufferings. Otherwise, we screened those related genes mentioned previously and most of the variants were unlikely to affect protein functions. The only one non-synonymous and shared by two siblings is just the causative mutation in ABCB4 filtered out by another analysis strategy [Table 1].

Sanger sequencing

Sanger sequencing was used to screen all of the four family members. Exon 17 of ABCB4 gene, where the mutation located, was amplified using polymerase chain reaction (forward

Case	1	2
Total effective yield(Mb)	3,801.77	3,935.91
Effective sequence on target(Mb)	2,290.43	2,296.60
Average sequencing depth on target	60.71	60.87
Coverage of target region	98.3%	98.4%
covered with at least 10X	89.1%	89.2%
covered with at least 20X	79.0%	79.2%
Total number of variants*	32,432	32,349
No. Non-synonymous variants	7432	7468
No. not in dbSNP	915	951
No. not in 1000 Genomes Project	572	579
No. not in HapMap 8	472	468
No. not in YH1 genome data	468	461
No. shared by 2 cases	250	
No. homozygous	4	
SIFT prediction		
No. Tolerated	3	
No. Damaging	1	

Coverage: the average number of times each base is represented in the sequence reads; dbSNP: the Single Nucleotide Polymorphism Database (build 129) of the National Center for Biotechnology Information; HapMap 8: eight previously exome-sequenced HapMap samples; 1000 Genomes Project: the SNP release of the 1000 Genome Project (20100208 release); YH1 genome data*YanHuang 1 genome database

*Variants were filtered gradually according the criteria.

Table 1: Results of exome sequencing and details of data analysis.

primer CGAACAAACCATACTCAGCTTATG, reverse primer GAGGTTGGGAGAAGCAGCAGC)[9]. Sequence analysis showed that the mutation was homozygous in the patients and heterozygous in their parents (Figure 1). The variant has never been described in the Genbank database, so the sequence was submitted to Genbank (accession number HQ540315) and the variation was submitted to dbSNP (submitter snp number 263198237).


Discussion

In this study, we have identified a homozygous ABCB4 gene mutation using exome sequencing and finally confirmed the diagnosis as PFIC3. ABCB4 gene encodes the class III multidrug resistance P-glycoprotein 3 (MDR3) belonging to the family of ATP-binding cassette transporters. It is localized at the canalicular membrane of hepatocytes translocating Phosphatidyl Choline (PC) from the inner to the outer membrane leaflet [10]. MDR3 has 12 transmembrane segments (TM1-TM12) organized in two transmembrane domains. The mutation p.P726L is located in the TM7 which is conserved in mammals ABCB4. Mutation detected in our patients is categorized as damaging by SIFT (Score 0.01) and seems to induce a significant impairment in flippase activity. Site-directed mutagenetic analysis of P-glycoprotein has shown that mutations in transmembrane domains are important for substrate specificity [9]. Otherwise, many missense mutations may influence protein processing, causing the abnormal but potentially functional protein to be misfolded, trapped in the endoplasmic reticulum and subsequently degraded [11]. Defects of MDR3 may further induce an imbalance in the composition of primary bile, with lack of PC and a surplus of bile salts, the latter presumed to damage the biliary epithelium and cause progressive liver disease such as PFIC3 [12]. Previous study have shown that the phenotypic spectrum of PFIC3 ranges from neonatal cholestasis to cirrhosis in young adults [6]. The two siblings in our study reveal the similar clinical course characterized by progressive cholestasis and liver dysfunction.

Further sequencing identified that the mutation was inherited from the non-consanguineous parents. Notably, both parents are Hui Chinese and absent of symptoms. Lang et al. [13] have intended to establish genetic variability and haplotype structures of ABCB4 in healthy populations of different ethnic backgrounds comprising

Caucasian, African-American, Japanese, and Korean origin. Their results showed that most ABCB4 variants were population-specific, and either their occurrence or their population frequency varied among different ethnic groups. The same mutation mentioned above was detected in a French patient and compound with another heterozygous mutation. Interestingly, Degiorgio et al. [9] have ever described a genotype in an Italian patient, showing three heterozygous mutations, one of which was located on the maternal allele and similar to our finding (c.2176C-A transition, p.P726T). It seems that mutations at this site may be more important than previous perspectives. These suggest that it would make sense to investigate the frequencies of the rare variants in specific groups, however there is still short of related epidemiologic data.

Another important task is to make genetic counseling and prenatal diagnosis to some extent. PFIC transmission is autosomal recessive and mutations on both alleles are characterized in most cases. This understanding has already allowed prenatal diagnosis. Camille et al. have shown that it is possible to perform reliable molecular prenatal diagnosis for PFIC [14]. However, objects of genetic counseling and antenatal diagnosis are usually limited in individuals with a family history or from close family. The presence of mutations in ABCB4 gene had never been described in siblings of non-consanguineous parents until Raj et al. reported it in two Sicilian siblings [15]. Therefore, it's really difficult to discover individuals with genetic susceptibility using traditional methods in the situation just as we have encountered. For this reason, exome sequencing, enabling low-cost, rapid, broad sequencing, may be a potentially valuable tool in genetic counseling and prenatal diagnosis. Sequencing data reflect most of the genetic background including some possible genetic defects and has the potential to broaden our knowledge of the genetic basis of diseases. By combining with different methods or using different analysis strategies, exome sequencing is apt to be as promising a route to mapping genes for diseases comprising not only monogenic disease but also more complex diseases.[4,16,17] In our study, a total of 250 novel variants are both carried by the two siblings and some of the heterozygous variants, categorized as damaging by SIFT, may bring potential risk to next generations. Unfortunately, there are few related publications in this area.



Conclusions

Briefly, we found a novel homozygous mutation in ABCB4 gene causing deleterious results. Further functional studies of this specific gene mutation are needed to gain a clear idea of the pathogenic mechanism. We also demonstrate that exome sequencing is a useful means for mapping monogenic disease mutation. In addition, we may investigate the genetic background of certain groups and predict risk of suffering diseases by applying it to genetic counseling and prenatal diagnosis. Mutation-specific therapy may be the most promising approach in the treatment of genetic diseases such as PFIC.

Authors' Contributions

H. L. and R. F. contributed equally to this work as co-first author by providing data collection and data analysis. X. Z., B. Y., J. Y., J. S. provided valuable opinions about the manuscript. W.S. and Z.L. conceived the idea and wrote the manuscript as co-corresponding author.

Grant Sponsor

The National High Technology Research and Development Program of China (2007AA02Z470).

References

1. Liu C, Aronow BJ, Jegga AG, Wang N, Miethke A, et al. (2007) Novel resequencing chip customized to diagnose mutations in patients with inherited syndromes of intrahepatic cholestasis. *Gastroenterology* 132: 119-126.
2. Oude Elferink RP, Paulusma CC (2007) Function and pathophysiological importance of ABCB4 (MDR3 P-glycoprotein). *Pflugers Arch* 453: 601-610.
3. Krawitz PM, Schweiger MR, Rödelsperger C, Marcelis C, Kölsch U, et al. (2010) Identity-by-descent filtering of exome sequence data identifies PIGV mutations in hyperphosphatasia mental retardation syndrome. *Nat Genet* 42: 827-829.
4. Musunuru K, Pirruccello JP, Do R, Peloso GM, Guiducci C, et al. (2010) Exome sequencing, ANGPTL3 mutations, and familial combined hypolipidemia. *N Engl J Med* 363: 2220-2227.
5. Davit-Spraul A, Gonzales E, Baussan C, Jacquemin E (2009) Progressive familial intrahepatic cholestasis. *Orphanet J Rare Dis* 4: 1.
6. Hori T1, Nguyen JH, Uemoto S (2010) Progressive familial intrahepatic cholestasis. *Hepatobiliary Pancreat Dis Int* 9: 570-578.
7. van der Woerd WL, van Mil SW, Stapelbroek JM, Klomp LW, van de Graaf SF, et al. (2010) Familial cholestasis: progressive familial intrahepatic cholestasis, benign recurrent intrahepatic cholestasis and intrahepatic cholestasis of pregnancy. *Best Pract Res Clin Gastroenterol* 24: 541-553.
8. Davit-Spraul A, Gonzales E, Baussan C, Jacquemin E (2010) The spectrum of liver diseases related to ABCB4 gene mutations: pathophysiology and clinical aspects. *Semin Liver Dis* 30: 134-146.
9. Degiorgio D, Colombo C, Seia M, Porcaro L, Costantino L, et al. (2007) Molecular characterization and structural implications of 25 new ABCB4 mutations in progressive familial intrahepatic cholestasis type 3 (PFIC3). *Eur J Hum Genet* 15: 1230-1238.
10. Oude Elferink RP1, Paulusma CC (2007) Function and pathophysiological importance of ABCB4 (MDR3 P-glycoprotein). *Pflugers Arch* 453: 601-610.
11. Delaunay JL, Durand-Schneider AM, Delautier D, Rada A, Gautherot J, et al. (2009) A missense mutation in ABCB4 gene involved in progressive familial intrahepatic cholestasis type 3 leads to a folding defect that can be rescued by low temperature. *Hepatology* 49: 1218-1227.
12. Stapelbroek JM, van Erpecum KJ, Klomp LW, Houwen RH (2010) Liver disease associated with canalicular transport defects: current and future therapies. *J Hepatol* 52: 258-271.
13. Lang T, Haber M, Jung D, Drescher A, Schlagenhauer R, et al. (2006) Genetic variability, haplotype structures, and ethnic diversity of hepatic transporters MDR3 (ABCB4) and bile salt export pump (ABCB11). *Drug Metab Dispos* 34: 1582-1599.
14. Jung C, Driancourt C, Baussan C, Zater M, Hadchouel M, et al. (2007) Prenatal molecular diagnosis of inherited cholestatic diseases. *J Pediatr Gastroenterol Nutr* 44: 453-458.
15. Crupi I (2010) Progressive familial intrahepatic cholestasis (PFIC) type 3 in two Sicilian siblings of nonconsanguineous parents. *ScientificWorldJournal* 10: 1065-1066.
16. Wijers CH, de Blaauw I, Marcelis CL, Wijnen RM, Brunner H, et al. (2010) Research perspectives in the etiology of congenital anorectal malformations using data of the International Consortium on Anorectal Malformations: evidence for risk factors across different populations. *Pediatr Surg Int* 26:1093-1099.
17. Zhou H, Alexander DH, Sehl ME, Sinsheimer JS, Sobel EM, et al. (2011) Penalized regression for genome-wide association screening of sequence data. *Pac Symp Biocomput*.

Citation: Lin H, Fu R, Zhang X, Yao B, Ye J, et al. (2014) A New Homozygous ABCB4 Mutation Identified in Two Chinese Siblings Based on Exome Sequencing. *J Genet Syndr Gene Ther* 5: 243. doi:10.4172/2157-7412.1000243

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 350 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/omicsgroup/>