

A Review on the Taxonomy, Ethnobotany, Phytochemistry and Pharmacology of *Guiera senegalensis* J.F.Gmel. (Combretaceae)

Mubarak Siddig Hamad^{1*}, Hassan Elsafi¹, Ahmed Saeed¹, Eltayeb Fadul¹, Reem Hassan Ahmed²

¹Department of Medicine and Aromatic, Traditional Medicine Institute, National Center for Research, Khartoum, Sudan

²Department of Biology and Biotechnology, Industrial Science University of Bahri, Sudan

*Corresponding author: Mubarak Siddig Hamad, Department of Medicine and Aromatic, Traditional Medicine Institute, National Center for Research, Khartoum, Sudan, Tel: +249916161428; Fax: +249183786085; E-mail: mubarakiddig23@yahoo.com

Received date: June 13, 2017; Accepted date: July 03, 2017; Published date: July 09, 2017

Copyright: © 2017 Hamad MS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Guiera senegalensis (*G. senegalensis*) Family Combretaceae is shrubs or trees widely distributed in west and central Africa. In traditional medicine *G. senegalensis* used to treat malaria, dysentery and/or diarrhea, cough and/cold, abdominal pains, leprosy, hypotension and hypertension, diabetes, snakebites eczema, impotence, epilepsy, breast cancer, depressant and jaundice. Phytochemical screening investigated by scientists showed the presence of alkaloids, flavonoids, tannin, saponin, terpinoids/sterols and cardionolics mainly in leaves and roots of the plant. The biological activities showed positive results as anti-tuberculosis, anti-diarrhoeal, anti-plasmodia, analgesic and devoid of anti-inflammatory, antifungal, antioxidant anti-lipid peroxidation activity, anti-malaria and with low toxicity, antifungal, anti-acetylcholinesterase, antilipid peroxidation in rat's brain homogenate, erythrocytes hemolysis inhibitory activities and antitussive activities. The review revealed the importance of *G. senegalensis* as a natural therapy and recommend for further study and the work of the pharmaceutical formulation.

Keywords: *Guiera senegalensis*; Malaria; Diarrhea; Cough; Cold; Hypertension; Hypotension; Diabetes

Introduction

Family Combretaceae is trees or shrubs, some time climbing plants [1], comprising about 20 genera and 500 species [2,3], these species distributed around the globe and central of diversity in Africa and Asia. *G. senegalensis* belongs to this family is widely distributed in west and central Africa on leached soils, fallows; as underwood in low savanna forests, mostly in sandy soils and on very dry sites; indicative of overgrazing [4]. *G. senegalensis* has numerous traditional medicinal applications, for instance, leaves for various internal diseases, prevention of leprosy, dermatoses, as tonic, infusions as diuretic, for stomach ache, cough and so on [4]. A range of phytochemical compounds have been isolated from different plant parts. Many of the uses have been subjected to some level of pharmacological screening, and tests on its antimalarial, anti-diarrhoeal, antibacterial, anti-cough, anti-inflammatory, anti-oxidant activity have been positive. Many of these tests, however, are still at a preliminary level, and need to be followed by more detailed research. Further toxicological studies are also warranted as it has been shown that most extracts when orally ingested seem to be relatively harmless, but when injected most extracts are toxic to varying levels. *Guiera senegalensis* is also a very important species in the crop-fallow cycle in the Sahelian zone, as well for its firewood production, and its presence should be monitored in order to prevent a decline of the species.

Phytochemical screening for *G. senegalensis* showed significant secondary metabolites namely anthraquinones, terpenoids, saponins, alkaloids, coumarins, mucilages, flavonoids, tannins and cardiotonic and cyanogenic heterosides were assayed in different organs of the plant, in leaves, fruits, roots and stem bark [5].

Many studies have been carried out to *G. senegalensis* to investigate its pharmacological properties; it showed significant results in different types of illness. The present review is an attempt to highlight the Morphology, Ethnobotany, Phytochemistry and Pharmacology activities of *G. senegalensis*, the thing that reflects its importance and encourage for further study on it (Table 1).

English name	Guiera
French name	Guier du Sénégal
Arabic name	Gubeish
Scientific classification	
Kingdom	Plantae
Phylum	Magnoliophyta
Class	Magnoliopsida
Order	Myrtales
Family	Combretaceae
Genus	<i>Guiera</i>
Species	<i>G. senegalensis</i>

Table 1: Classification of *Guiera senegalensis*.

Guiera senegalensis J.F.Gmel. is an accepted name, this name is the accepted name of a species in the genus *Guiera* (family Combretaceae). The record derives from World Checklist of Selected Plant Families (WCSP) (in review) (data supplied on 2012-03-23) which reports it as an accepted name with original publication details: Syst. Nat. 675 1791. (www.theplantlist.org.).

Guiera glandulosa Sm. is a synonym of *Guiera senegalensis* J.F.Gmel. This name is a synonym of *Guiera senegalensis* J.F.Gmel.

The record derives from WCSP (in review) (data supplied on 2012-03-23) which reports it as a synonym with original publication details: *Cycl.* 17: s.p. 1811.(www.theplantlist.org.)

Morphology

Small shrub up to 3 m high, with scattered black dots; branches pubescent. Leaves grey-green, single or opposite, oblong-elliptic,

mucronate at the apex, rounded or slightly cordate at the base. About 2 to 4 cm long by 1 to 2 cm wide, softly tomentose on both surfaces, with scattered black glands beneath lateral nerves fine. Flowers globose heads, yellow-greenish, resting on short axillary peduncles. Calyx of 5 lobes densely covered by black spots, corolla of 5 petals, the stamens are 10 on two rows of 5, all inserted on the calyx [4,6,7]. The ovary has a single box containing 4 to 6 eggs [8]. Fruit narrowly cylindrical, radiating, 3.5 cm long, densely clothed with very long silky hairs, making a fluffy head (Figure 1). Flowers Sept.-Jan.; Fruits Feb.-May [6]. Source of photos; Global Biodiversity Information Facility (GBIF).



Figure 1: Close view of *Guiera senegalensis*.

Geographical distribution

Guiera senegalensis widely distributed in the savannah region of west and central Africa in Senegal, Gambia, Mali, Niger, Burkina Faso, Guinea-Bissau, Guinea, Nigeria, Chad, Mauritania and Sudan (Figure 2).

Traditional medicine uses

G. senegalensis has been reported to have numerous traditional medicinal uses, many ethnobotanical studies have been conducted by many authors confirmed that *G. senegalensis* has a good reputation as medicinal traditional plant. It is reported that it is used to treat dysentery and/or diarrhea [4,7,9-14]. Many authors reported that *G. senegalensis* used in traditional medicine to treat Malaria and /or fever [5,7,11,14-22]. It is also reported as antileprosy [4,5,7,11,17,21], Abdominal pains [4,7,9,10], Epilepsy [7], depressant [23], cough and/or cold, [4,7,15,24,25], snakebite [11,26], Eczema [7], syphilis [5], hypertension and hypotension [11,24,25], diabetes [11,17], Impotence [5,7], Breast cancer and Jaundice [27,28].

Phytochemistry review

Phytochemicals screening of different parts of *Guiera senegalensis* have been investigated by researchers. They found that leaves revealed the presence of alkaloids, flavonoids, tannins [7,24,29], saponins [24]

and terpenoids/sterols [7,29]. Also, it reported a new methoxylated naphthyl butenone, guieranone A (1) was isolated from the leaves of *Guiera senegalensis*. Its structure was elucidated as (2E)-1-(1,3,6,8-tetramethoxy-2-naphthyl) but-2-en-1-one, on the basis of spectroscopic data. Also isolated were two known naphthopyrones, 5-methyldihydroflavasperone (2) and 5-methylflavasperone [30]. The roots revealed the presence of steroids, saponins, alkaloids, terpenoids, tannins, flavonoids, cardionolic [11]. Guieranone A, a naphthyl butanone, three beta-carboline alkaloids and tetrahydroharman were purified from leaves and roots and Harmalan (dihydroharman) was isolated for the first time [31]. Four components, catechin, myricitrin, rutin and quercetin were evidenced and determined by HPLC from *G. senegalensis* [32]. Thirteen (13) components were identified for hexane-acetone (50:50) column fraction of hydroacetone extract and twenty one (21) compounds for hexane-acetone (50:50) column fraction of aqueous decoction extract [33]. Also, Methylidihydroflavasperone, a novel naphthopyran, was isolated from the chloroform extract of the leaf of *Guiera senegalensis* [34]. Mucilagens, tannins, flavonoids, alkaloids and amino acids are the so far known constituents of *Guiera senegalensis* [35,36]. from the methanolic extract of the dried leaves of this plant, flavonol aglycones as well as flavonol glycosides, some of them acylated, were isolated [37]. four flavonoids from the leaves of *G. senegalensis*, namely catechin, myricitrin, rutin and quarterin have been isolated by Ficarra et al. [32].




Figure 2: Distribution of *Guiera senegalensis* in Africa.

Pharmacological activities review

G. senegalensis has various pharmacological properties that have been reported by researchers. It showed positive effects from the methanolic roots extracts as anti-tuberculosis [38], and anti-diarrhoeal [10,11,39], also it exhibited significant result as ulcer-protective [39]. The aqueous and ethanolic leaves extract that at lower doses was not harmful to the liver and therefore can be exploited as it is served in the treatments of some illnesses [14]. Methanolic extract of leaves revealed positive result as anti-plasmodia, analgesic and devoid

of anti-inflammatory activities [5]. It is also reported that The gall *G. senegalensis* exhibited an interesting antifungal activity against all strains tested [33] while the leaves extract showed that it has no inhibition activity against all of the tested fungal strains [29], while Guieranone which isolated from the leaves of *G. senegalensis* exhibited potent activity against the fungus *Cladosporium cucumerinum* [30]. The extract of the leaves of *G. senegalensis* was found to detoxify (*in vitro*) venom from two snake species [40], Total alkaloid extract and beta carboline alkaloids presented an interesting antiplasmodial activity associated with a low cytotoxicity [31]. The extract (500 g) effectively

inhibited the formation of peroxides in sunflower oil and showed the highest antioxidative activity [41]. The chloroform extract of roots of *G. senegalensis* exhibited a pronounced ant-malarial activity while two alkaloids isolated from the active extract, Harman and tetrahydroharman, showed antimalarial activity and displayed low toxicity [18]. The gall of *G. senegalensis* by using hydroacetic extract showed the most potent antioxidant activity. And anti-lipid peroxidation activity [33]. It is also reported that gallotannins and tow condensed tannins exhibit interesting antioxidant properties [42]. The extract from galls of *G. senegalensis* possess effective anti-acetylcholinesterase, antilipid peroxidation in rat's brain homogenate and erythrocytes hemolysis inhibitory activities [43]. Leaf decoction of *G. senegalensis* was reported to possess an antitussive activity [44]. It was also reported that *G. senegalensis* the crude extracts possess *in vitro* anthelmintic activity against *Haemonchus contortus* in goat [9].

Conclusion

This review reflected that *Guiera senegalensis* is widely used traditionally as therapy for different ailments and its Phytochemical screening revealed the presence of secondary metabolites in different parts of *G. senegalensis* in particularly leaves and roots in addition to its various pharmacological properties, accordingly the Authors recommend for further study and the work of the pharmaceutical formulation of *G. senegalensis*.

References

- Andrews WF (1950) The flowering plants of the Anglo-Egyptian Sudan. Vol. I (Cycadaceae-Tiliaceae): The flowering plants of the Anglo-Egyptian Sudan. Vol. I (Cycadaceae-Tiliaceae). pp: 237.
- Heywood V, Brummitt R, Culham A, Seberg O (2007) Flowering Plant Families of the World: Royal Botanic Gardens Kew.
- Chase MW, Reveal JL (2009) A phylogenetic classification of the land plants to accompany APG III. *Botanical Journal of the Linnean Society* 2: 122-127.
- Maydell HJV, John B (1986) Trees and Shrubs of the Sahel: Their Characteristics and Uses.
- Jigam AA, Akanya HO, Dauda BE, Ogbadoyi EO (2011) Antiplasmodial, analgesic and anti-inflammatory effects of crude *Guiera senegalensis* Gmel (Combretaceae) leaf extracts in mice infected with *Plasmodium berghei*. *J Pharmacogn Phytother.* pp: 150-154.
- Amin HME (1990) Trees and Shrubs of the Sudan.
- Somoro AA, Patel K, Diallo D, Sidibe L, Chalchat J, et al. (2011) An ethnobotanical and phytochemical study of the African medicinal plant *Guiera senegalensis* JF Gmel. *J Med Plants Res* 9: 1639-1651.
- Dénou A, Togola A, Haïdara M, Sanogo R, Diallo D, et al. (2016) Review on Phytochemistry and Pharmacological Aspects of *Guiera Senegalensis* JF Gmel (Combretaceae). *International Journal of New Technology and Research* 3: 1.
- Suleiman MM, Mamman M, Sidiama A, Ighoja EJ, Tauheed M, et al. (2014) Evaluation of anthelmintic activity of Nigerian ethnoveterinary plants; *Cassia occidentalis* and *Guiera senegalensis*. *Veterinary World* 7: 3.
- Williams E, Barminas J, Akinniyi J, William A (2009) Antidiarrhoeal effects of the root extracts of *Guiera senegalensis* in male mice. *African Journal of Pure and Applied Chemistry* 8: 152-157.
- Shettima YATM, Karumi Y, Sodipo OA (2012) Phytochemical and Antidiarrhoeal Properties of Methanol root Extract of *Guiera senegalensis* J.F.Gmel. *International Research Journal of Pharmacy* 11: 3.
- Hutchinson J, Dalziel J (1965) Flora of West Tropical. Part 1. Crown Agents for Oversea Governments and Administrations, London. p: 275.
- Maxia A, Demurtas A, Kasture S, Kasture V, Fadda V, et al. (2014) Medical ethnobotany survey of the Senegalese community living in Cagliari (Sardinia, Italy). *Indian Journal of Traditional Knowledge* 2: 275-282.
- Oshobu M, Geidam M (2014) Effects of Aqueous Ethanolic Extract of the leaves of *Guiera senegalensis* JF Gmel (Combretaceae) on Liver Function in Wister Strain Albino Rats. *International Journal of Scientific Research and Management.* pp: 611-615.
- Sourabié T, Some N, Bognonou O, Ouattara Y, Ouédraogo J (2013) Ethnobotanical and ethnopharmacognostical survey on medicinal plants of Malon village and surrounding in the Cascades Region (Burkina Faso). *IOSR Journal of Pharmacy* pp: 11-15.
- Gamal EB, Awatif AB, Egami E, Waïl SA, Galal M (1997) Medicinal plants of Northern Kordofan: Medicinal plants of the Sudan.
- Gamal EB, Ghazali EAB, Ahmed K, Kariem MS (1987) Medicinal plants of the Eastern Nuba Mountaints Medicinal plants of the Sudan.
- Ancolio C, Azas N, Mahiou V, Ollivier E, Di Giorgio C, et al. (2002) Antimalarial activity of extracts and alkaloids isolated from six plants used in traditional medicine in Mali and Sao Tome. *Phytotherapy Research* 7: 646-649.
- Diarra N, van't Klooster C, Togola A, Diallo D, Willcox M, et al. (2015) Ethnobotanical study of plants used against malaria in Sélingué subdistrict, Mali. *Journal of Ethnopharmacology* pp: 352-360.
- Singh S, Singh R (2014) Herbal medicinal treatment of Malaria in Aliero Local Government Area, Kebbi, Nigeria. *Journal of Medicinal Plants* 2: 117-126.
- Oluranti AC, Michael UO, Jane U, Ayembe NA (2012) Ethno botanical studies of medicinal plants used in the management of Peptic ulcer disease in Sokoto State, North Western Nigeria. *Int Res J Pharm Pharmacol* 9: 225-230.
- Ene A, Atawodi S, Ameh D, Kwanashie H, Agomo P (2010) Locally used plants for malaria therapy amongst the Hausa, Yoruba and Ibo Communities in Maiduguri, North-Eastern Nigeria. *Indian Journal of Traditional Knowledge* 3: 486-490.
- Maria MR, Maria CD, Bucar I, Luis C (2012) Medicinal plants used to treat neurological disorders in West Africa: a case study with Guinea-Bissau flora. *American Journal of Plant Sciences.* pp: 1028-1036.
- Salihu SO, Usman AA (2015) Antimicrobial and phytochemical study of the bioactive fractions of *Guiera senegalensis* from Alasan Tambuwal, Nigeria. *Journal of Pharmacognosy and Phytochemistry* 6: 106-111.
- Dénou A, Togola A, Haïdara M, Sanogo R, Diallo D, et al. (2016) Review on Phytochemistry and Pharmacological Aspects of *Guiera Senegalensis* JF Gmel (Combretaceae). *International Journal of New Technology and Research* 3: 30-32.
- Sallau A, Njoku G, Olabisi A, Wurochekke A, Abdulkadir A, et al. (2005) Effect of *Guiera senegalensis* leaf extract on some *Echis carinatus* venom enzymes. *J Med Sci* 4: 280-283.
- Fadimu YO, Mohammed I, Zurmi S (2014) Ethnomedicinal survey of anti-typhoid plants in Ijebu Ode Local Government Area of Ogun State, Nigeria. *International Journal of Science and Nature* 2: 332-336.
- Ohemu T, Agunu A, Olotu P, Ajima U, Dafam D, et al. (2014) Ethnobotanical Survey of Medical Plants Used in the Traditional Treatment of Viral Infections in Jos, Plateau State, Nigeria. *Int J Med Arom Plants* 4: 74-81.
- Alshafei N, Elshafie A, Nour A (2016) Antitoxic, Antifungal and Phytochemical Analysis of Medicinal Compounds of *Guiera senegalensis* Leaves in Sudan. *J Plant Biochem Physiol* 4: 166.
- Silva O, Gomes ET (2003) Guieranone A, a naphthyl butenone from the leaves of *Guiera senegalensis* with antifungal activity. *Journal of Natural Products* 3: 447-449.
- Fiot J, Sanon S, Azas N, Mahiou V, Jansen O, et al. (2006) Phytochemical and pharmacological study of roots and leaves of *Guiera senegalensis* JF Gmel (Combretaceae). *Journal of Ethnopharmacology* 2: 173-178.

32. Ficarra R, Ficarra P, Tommasini S, Carulli M, Melardi S, et al. (1997) Isolation and characterization of *Guiera senegalensis* JF Gmel. active principles. *Bollettino Chimico Farmaceutico* 5: 454-459.
33. Sombié PA, Konate K, Youl E, Coulibaly AY, Kiendrébéogo M, et al. (2013) GC-MS analysis and antifungal activity from galls of *Guiera senegalensis* JF Gmel (Combretaceae). *Journal of Applied Pharmaceutical Science* 12: 6.
34. El Hadi NM, Khalid SA (1997) 5-methyldihydroflavasperone, a dihydronaphthopyran from *Guiera senegalensis*. *Phytochemistry* 4: 793-794.
35. Combier H, Becchi M, Cavé A (1977) Alcaloïdes du *Guiera senegalensis* Lam *Plantes medicinales et phytothérapie*.
36. Makkar HP, Becker K (1994) Isolation of tannins from leaves of some trees and shrubs and their properties. *Journal of Agricultural and Food Chemistry* 42: 731-734.
37. Bucar F, Schubert-Zsilavecz M, Knauder E (1996) Flavonoids of *Guiera senegalensis*. *Pharmazie* 7: 517-518.
38. Adedayo AV (2015) Pricing de Solla Price's circumvent. *Adv Res*, pp: 488-492.
39. Aniagu S, Binda L, Nwinyi F, Orisadipe A, Amos S, et al. (2005) Anti-diarrhoeal and ulcer-protective effects of the aqueous root extract of *Guierasenegalensis* in rodents. *Journal of Ethnopharmacology* 3: 549-554.
40. Abubakar M, Sule M, Pateh U, Abdurahman E, Haruna A, et al. (2000) In vitro snake venom detoxifying action of the leaf extract of *Guiera senegalensis*. *Journal of Ethnopharmacology* 3: 253-257.
41. Mariod A, Matthäus B, Hussein I (2006) Antioxidant activities of extracts from *Combretum hartmannianum* and *Guiera senegalensis* on the oxidative stability of sunflower oil Emirates. *Journal of Food and Agriculture* 2: 20-28.
42. Bouchet N, Barrier L, Fauconneau B (1998) Radical scavenging activity and antioxidant properties of tannins from *Guiera senegalensis* (Combretaceae). *Phytotherapy Research* 3: 159-162.
43. Sombie P, Hilou A, Conulibaly AY, Tibiri A, Kiendrebeogo M, et al. (2011) Brain protective and erythrocytes hemolysis inhibition potentials from galls of *Guiera senegalensis* JF Gmel (Combretaceae). *Journal of Pharmacology and Toxicology* 4: 361-370.
44. Sanogo R, De Pasquale R, Germano M (1998) The antitussive activity of *Guiera senegalensis* JF Gmel (Combretaceae). *Phytotherapy Research* 2: 132-134.