

A Short Review: Complementary and Alternative Medicine in Lung Cancer

Raj Parikh¹ and Niranjan Jeganathan^{2*}

¹Department of Internal Medicine, Rush University Medical Center, USA

²Division of Pulmonary and Critical Care Medicine, Rush University Medical Center, USA

*Corresponding author: Niranjan Jeganathan MD, Division of Pulmonary and Critical Care Medicine, Rush University Medical Center, 1750 W. Harrison Street, 297 Jelke South Center, Chicago, IL 60612, USA, Tel: 312 942-2552; Fax: 312 942-8187; E-mail: Niranjan_Jeganathan@Rush.edu

Rec date: Jun 18, 2014, Acc date: Aug 20, 2014, Pub date: Aug 22, 2014

Copyright: © 2014 Parikh R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Complementary and alternative medicine (CAM) is a growing field that has started to integrate itself into the management of many diseases and disorders, including cancer. As more patients continue to incorporate the use of CAM into their treatment plans, it is important to understand the mechanism and clinical effectiveness of CAM. This review focuses on the role of CAM in lung cancer, including the available research trials that have focused on evaluating symptom control, quality of life, and survival rates in lung cancer patients. We have also reviewed the potential risks and benefits of several CAM medications as well as the future direction of this field.

Keywords: Herbal; Complementary; Alternative; Lung Cancer; Chemotherapy

Introduction

Throughout the history of medicine, complementary and alternative medicine (CAM) has played a significant role in the treatment of cancer patients. In 1999, the National Institutes of Health (NIH) created the National Center for Complementary and Alternative Medicine (NCCAM) in recognition of this growing field. Research funding to the NCCAM has risen from \$ 2 million in 1992 to \$50 million by 2000 and is expected to be more than \$110 million by the mid-2000's [1]. A survey conducted in 2005 indicated that approximately 35.9% of cancer patients attempted use of CAM during their treatment course [2]. In fact, that percentage may be even higher as studies have shown that many patients do not disclose the use of CAM therapy unless asked specifically [3]. Physicians poorly communicate with their patients regarding the potential benefits or harms of CAM because they have minimal understanding of CAM therapy and its physiological role in managing diseases [4,5]. Patients are often dissatisfied with their physicians' lack of knowledge regarding CAM therapy [6]. Patients who have a history of incorporating CAM into their management plan are generally driven by personal beliefs and principles regarding life and health [7]. An increased use of CAM has been seen in patients with increased psychosocial stress or poor prognosis at initial diagnosis [8,9]. Additionally, females and younger patients have a higher predilection for using CAM therapy [10]. There are numerous modalities of CAM such as spiritual healing or prayer, herbal medicine, and chiropractics [11]. This review focuses on medicinal therapies used as CAM for patients suffering from malignancy, specifically lung cancer.

Methods

A detailed search was performed on PubMed using the following terms: "lung cancer" and "herbal therapy", "complementary and alternative medicine", "CAM" AND "malignancy", "CAM" AND "lung cancer", and "CAM" AND "chemotherapy". Relevant clinical

trials were included. The bibliographies of these and other review articles were reviewed in order to identify additional relevant material.

CAM in Lung Cancer

CAM specific to lung cancer has focused on therapeutic options that may have less systemic toxicity compared to current chemotherapy options such as Gemcitabine, Paclitaxel, and Etoposide. Herbal plants that have been used in lung cancer therapy include *Platycodon grandiflorum* (Campanulaceae), *Morus alba* (Moraceae), *Prunus armeniaca* (Rosaceae), *Rhus verniciflua* (Anacardiaceae), *Perilla frutescens* (Labiatae), *Stemona japonica* (Stemonaceae), *Tusilago farfara* (Compositae), and *Draba nemorosa* (Brassicaceae) [12]. Although these are used as alternative modes of therapy, it is more common for herbal remedies to be used as adjuvant modalities with chemotherapy [13]. The role of these herbal therapies is multifactorial. They can be used to reduce chemotherapy-induced toxicity, mitigate cancer-related symptoms, or directly increase anti-cancer effects [14]. A randomized control trial (RCT) of 61 patients evaluated the use of herbal medications as adjuvant therapy and showed an enhancement in median survival in the group that was treated with Sheng-mai and Gu- jin Granule along with Navelbine and Cisplatin [15]. This group had a 48.5% increase in survival compared to 32.2% in the control arm that only received the two chemotherapy drugs. Another study enrolled 232 non-small cell lung cancer patients and showed that Shenqi-fuzheng, an herbal injection similar to Sheng-mai, served as a useful adjuvant to chemotherapy by improving quality of life [16]. Similarly another RCT of 121 patients with advanced non-small cell lung cancer showed improved quality of life and a trend towards increase in survival time in patients assigned to Kangliuzhengxiao and Feiyanning in addition to chemotherapy [17]. Fiji Recipe therapy has been shown to alleviate chemotherapy side effects and enhance therapeutic efficacy [18]. Similarly, other studies have demonstrated the benefits of CAM therapy in mitigating the adverse effects of chemotherapy, including myalgia and diarrhea [19,20]. A case report showed regression of small cell lung cancer with the use of Chinese herbal medicines [21]. Lastly, another case report

showed that herbal remedies decrease tumor marker levels in lung cancer [22].

Discussion

In the last few years there has been more research devoted to CAM. However, there still remain many obstacles for CAM to be integrated into cancer treatment, especially lung cancer. The lack of resources (plants) to produce these herbal remedies could result in inadequate supply for the potentially high demand. The technology required for plant extraction and mixture preparation is another hurdle in establishing an adequate supply. Another obstacle is the lack of

understanding of the amount of herbal medication needed for monotherapy or adjuvant therapy. What dose is safe? What dose could interact with other chemotherapy drugs? In 2008, a study showed adverse drug interactions when a mixture of herbal medications including Ginseng and Fomesfomentarius were used with Gefitinib [23]. While CAM provides potential treatment options, there is much to be understood. It will be important to incorporate CAM into larger trials to measure its effect on survival, safety, and quality of life. In a medical world that continues to strive towards understanding how therapies work it is important to study the mechanisms and signaling pathways regulated by CAM in bench research and preclinical animal studies.

Study	Medication	Study Design	Sample Size	Control (Chemo therapy alone)	Results
Chen YZ, Li ZD, Gao F, Zhang Y, Sun H, et al. [15]	Sheng-mai, Gu-jin Granule	Randomized control study	n=61	Navelbine plus cisplatin	48.5% survival rate compared to 32.2% survival in control subjects
Lin H, Li DR [16]	Shenqi-Fuzheng	Randomized control study	n=232	Navelbine plus cisplatin or paclitaxel plus cisplatin	Improved quality of life* in experimental arm
Xu ZY, Jin CJ, Zhou CC, Wang ZQ, Zhou WD, et al. [17]	Kangliuzeng xiao, Feiyanning	Randomized control study	n=121	Navelbine plus cisplatin	Improved quality of life** and a trend toward prolonged survival time in experimental arm
Tian JH, Liu LS, Shi ZM, Zhou ZY, Wang L [18]	Fiji Recipe	Randomized control study	n=70	Vinorelbine plus cisplatin or gemcitabine plus cisplatin	Improved quality of life* in regards to side effects of chemotherapy in experimental arm
Yamamoto K, Hoshiai H, Noda K [19]	Shakuyaku-kanzo-to	Randomized control study	n=10	Paclitaxel plus carboplatin	Decreased adverse effects of chemotherapy- induced myalgia
Mori K, Kondo T, Kamiyama Y, Kano Y, Tominaga K [20]	Hangeshash in-to	Randomized control study	n=44	Cisplatin plus irinotecan	Decreased adverse effects of chemotherapy- induced diarrhea
*As assessed by quality of life scale of European Organization for Research on treatment of Cancer					
**As assessed by the Karnofsky score					

Table 1: List of CAM Therapies Evaluated in Clinical Trials as adjuvant to chemotherapy.

References

- Institute of Medicine (2003) Exploring complementary and alternative medicine. In: The Richard and Hinda Rosenthal Lectures 2001. The National Academies Press 26.
- Molassiotis A, Fernandez-Ortega P, Pud D, Ozden G, Scott JA, et al. (2005) Use of complementary and alternative medicine in cancer patients: a European survey. *Ann Oncol* 16: 655-663.
- Metz J, Jones H, Devine P, Hahn S, Glatstein E (2001) Cancer patients use unconventional medical therapies far more frequently than standard history and physical examination suggest. *Cancer J* 7: 149-154.
- Schofield PE, Juraskova I, Butow PN (2003) How oncologists discuss complementary therapy use with their patients: an audio-tape audit. *Support Care Cancer* 11: 348-355.
- Newell S, Sanson-Fisher RW (2000) Australian oncologists' self-reported knowledge and attitudes about non-traditional therapies used by cancer patients. *Med J Aust* 172: 110-113.
- Gray RE, Greenberg M, Fitch M, Parry N, Douglas M, et al. (1997) Perspectives of cancer survivors interested in unconventional therapies. *J Psychosoc Oncology* 15: 149-171.
- Astin JA (1998) Why patients use alternative medicine: results of a national study. *JAMA* 279: 1548-1553.
- Burstein HJ, Gelber S, Guadagnoli E, Weeks JC (1999) Use of alternative medicine by women with early-stage breast cancer. *N Engl J Med* 340: 1733-1739.
- Risberg T, Kaasa S, Wist E, Melsom H (1997) Why are cancer patients using non-proven complementary therapies? A cross-sectional multicentre study in Norway. *Eur J Cancer* 33: 575-580.
- Gansler T, Kaw C, Crammer C, Smith T (2008) A population-based study of prevalence of complementary methods use by cancer survivors: a report from the American Cancer Society's studies of cancer survivors. *Cancer* 113: 1048-1057.
- Ni H, Simile C, Hardy AM (2002) Utilization of complementary and alternative medicine by United States adults: results from the 1999 national health interview survey. *Med Care* 40: 353-358.
- Liang HLM, Xue CCL, Zhou DH, Li CG (2011) Chinese herbal medicine for lung cancer: a critical literature review. *CJIM* 9: 157-160.
- Richardson MA, White JD (2000) Complementary/alternative medicine and cancer research. A national initiative. *Cancer Pract* 8: 45-48.
- Gerber B, Scholz C, Reimer T, Briese V, Janni W (2006) Complementary and alternative therapeutic approaches in patients with early breast cancer: a systematic review. *Breast Cancer Res Treat* 95: 199-209.

15. Chen YZ, Li ZD, Gao F, Zhang Y, Sun H, et al. (2009) Effects of combined Chinese drugs and chemotherapy in treating advanced non-small cell lung cancer. *Chin J Integr Med* 15: 415-419.
16. Lin HS, Li DR (2007) Multi-center randomized clinical study on Shenqi-fuzheng injection combined with chemotherapy in the treatment for lung cancer. *Zhonghua Zhong Liu Za Zhi* 29: 931-934.
17. Xu ZY, Jin CJ, Zhou CC, Wang ZQ, Zhou WD, et al. (2011) Treatment of advanced non-small-cell lung cancer with Chinese herbal medicine by stages combined with chemotherapy. *J Cancer Res Clin Oncol* 137: 1117-1122.
18. Tian JH, Liu LS, Shi ZM, Zhou ZY, Wang L (2010) A randomized controlled pilot trial of "Feiji Recipe" on quality of life of non-small cell lung cancer patients. *Am J Chin Med* 38: 15-25.
19. Yamamoto K, Hoshiai H, Noda K (2001) Effects of shakuyaku-kanzo-to on muscle pain from combination chemotherapy with paclitaxel and carboplatin. *Gynecol Oncol* 81: 333-334.
20. Mori K, Kondo T, Kamiyama Y, Kano Y, Tominaga K (2003) Preventive effect of Kampo medicine (Hangeshashin-to) against irinotecan-induced diarrhea in advanced non-small-cell lung cancer. *Cancer Chemother Pharmacol* 51: 403-406.
21. Liang HL, Xue CC, Li CG (2004) Regression of squamous cell carcinoma of the lung by Chinese herbal medicine: a case with an 8-year follow-up. *Lung Cancer* 43: 355-360.
22. Kamei T, Kumano H, Iwata K, Nariai Y, Matsumoto T (2000) The effect of a traditional Chinese prescription for a case of lung carcinoma. *J Altern Complement Med* 6: 557-559.
23. Hwang SW, Han HS, Lim KY, Han JY (2008) Drug interaction between complementary herbal medicines and gefitinib. *J Thorac Oncol* 3: 942-943.