

A Short Review on Chiral Alcohols Verses Bio-Catalysis

Sreelakshmi Chittamuru^{*1,2}, BVS Reddy² and AB Rao²

¹Palmer Centre for Chiropractic Research, 734 Brady Street, Davenport Iowa, 52803, USA

²Indian Institute of Chemical Technology, Hyderabad, Telangana-500007, India

*Corresponding author: Sreelakshmi Chittamuru, Palmer Centre for Chiropractic Research, 734 Brady Street, Davenport Iowa, 52803, USA, Tel: 5635282192; E-mail: chittamurus@gmail.com

Received date: February 26, 2016; Accepted date: April 27, 2016; Published date: May 28, 2016

Copyright: © 2016 Sreelakshmi Chittamuru, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Catalyst is a chemical molecule or metal substance which enhances the rate of reaction is called catalyst, the catalyst present in the living organisms to carryout biochemical reactions or metabolic pathways. Exames oxidoreductases, lyases, ligases, proteases, hydrolases, esterases, pectinases etc. are Different methodologies available in synthesis of chiral molecules chemical/biocatalysis. The enormous potential of biocatalysts (microorganisms and enzymes) in synthesis of chiral molecules in mild conditions (pH and temperate) with high chemo-, regio-, enantio and functional selectivity with decrease formation of by-products, with short reaction steps (reactions which are not easily conducted classical organic reactions) (Long chemical processes with tedious blocking and de-blocking steps). Thus the use of biocatalyst has attracted a great attention from the green chemistry perspective (expensive chiral reagents/ environmentally hazardous heavy metals).

Biocatalysts are well known for their enzymatic activity towards numerous reactions ranging from in-vivo living cells (biochemical pathways) to in-vitro chemical reactions (reduction and transesterification) with enantiomeric purity and specificity. In this review the discussion is about synthesis of chiral using different kinds of biocatalysts. *Daucus Carota*, *Pisum Sativa*, *Novozyme P-435*, as biocatalysts for chiral alcohols.

Keywords: Biocatalysis; Biocatalysts; Sterio selectivity; Chiral alcohols; Anti-fungal; Antimicrobial; Active pharma ingredients; Pharma intermediates; Agrochemicals; Flavours; Lactonization; Bioreduction; Trans-esterification; Lipase; *Novozyme P-435*

Overview of the Chiral Alcohols Verses Bio-catalysis

Life and chirality are strictly connected in carrying out different metabolic functions of the living organism, thus chirality plays an important role in the life of plants and animals. Louis Pasteur (1853) coined the name dissymmetric. Later the term was changed to asymmetry and later it was replaced by the term chirality, a familiar analogy of the mirror image relationship like left and right [1-14]. The chemical complexity is the fundamental from of life and thus the chirality occupies the main features of the living world. Chirality is a phenomenon of geometrical or spatial arrangement of an object which is a non-super imposable mirror image. Chiral molecules that consist of a tetrahedral carbon atom attached to four different groups are called stereo-isomers, and possess identical chemical properties [12-25].

Chiral molecules are widely distributed in nature, and these enantioselective compounds are known to have selective/specific potent biological activities with reduced side effects compared to their racemates [26-29]. Hence, the synthesis of chiral molecules has drawn immense attention in the pharmaceutical industry as well as by drug regulatory authorities (FDA). A majority of naturally obtained compounds are chiral molecules that are active single isomers. For e.g., between the two enantiomers of Ketamine, S(+) Ketamine is the active anesthetic whereas the other isomer, R(-)Ketamine show undesirable side effects [30-38].

Reduction of different classes of prochiral ketones ranging from aromatic, heterocyclic and aliphatic ketones to their corresponding chiral alcohols. These chiral secondary alcohols are being used as drug intermediates in pharmaceuticals and also as toxophores in agrochemicals. These chiral synthons can be synthesized by economically viable and eco-friendly methods by using biocatalysts [39-48]. The use of biocatalysts from plant sources as enzymes have been well recognized for the synthesis of chiral products. Among the plant enzymes, sprouted seeds (e.g., *Pisum sativum*) have been considered as suitable biocatalytic systems for the reduction of aliphatic and aromatic prochiral ketones. an enzymatic approach for synthesis of (S)-1-phenethyl alcohol in 72% yield with 98% ee, by selective reduction of acetophenones, using sprouted *Pisum sativum* in an aqueous medium. Similarly substituted acetophenone derivatives, aliphatic and hetero cyclic ketones, tetrahydro pyran-4-ones and β -keto esters were taken up for selective bio reduction by sprouted *Pisum sativum* to obtain chiral secondary alcohols with a high degree purity with 91-98% enantioselectivity (ee) showing (S) configuration, following Prelog's rule. Enzymatic approach for the synthesis of chiral tetrahydropyranols by enantioselective reduction of 2-substituted tetrahydropyran-4-ones using *Daucus carota*. Reduction of (\pm)-2-phenyl-tetrahydro-2H-pyran-4-ones with *Daucus carota* afforded (2S, 4S)-2-phenyl-tetrahydropyranol and (2R,4S)-2-phenyl-tetrahydropyranol in a 1:1 ratio. The enantiomeric excesses 92% was observed in both the cis- & trans-tetrahydropyranols as determined by HPLC using chiral columns. The absolute stereochemistry of (2R, 4S)- & (2S, 4S)-2-aryl- or 2-alkyl-tetrahydropyranols was established by comparison with authentic samples. This methodology has wide substrate selectivity like alkyl and aryl pro chiral ketones to give corresponding chiral alcohols with optical purity and enantio & stereo selectivity. Acetyl-pyridines which are known as aromatic components

of perfumes, smoking suppressants and flavours used in foods [5,6,49-54]. Heterocyclic aromatic compounds contain nitrogen or oxygen or sulfur in the heterocyclic ring is important core groups found in natural and synthetic products of biological interest. Chiral hetero aryl alcohols have numerous applications as important intermediates in the synthesis of biologically active molecules and also act as chiral ligands/auxiliaries in a number of asymmetric addition reactions. This study highlights the selective asymmetric reduction of the heteroaryl prochiral ketones to the corresponding chiral alcohols using *Daucus carota*. The results confirms that the production of high yields of chiral alcohols (60% - 95%) with high enantioselectivity (76% - 99%) for (S)-configuration following the Prelog's rule. Only a single isomer was obtained, as confirmed through HPLC using a chiral column (no second isomer was present in the reaction medium). The un-reacted starting compound was recovered [7-13, 55-67].

Total Synthesis of Some Chiral γ -Lactones

Current discussion highlights the total synthesis of γ -Lactones like the fungicide Paecilocin A, (-)-3-butyl-7-hydroxyphthalide and the insect pheromones (4R)-Dodecanolide and (4R)-Octanolide molecules. Total synthesis of polyketide Paecilocin A which was isolated from the fungus Paecilomyces variotii. This molecule is known to exhibit inhibitory activity against pathogenic bacteria including methicillin-resistant bacteria. The molecule 3-butyl-7-hydroxyphthalide was isolated from a culture broth of Penicillium vulpinum. Furthermore, the lactones are known to function such as hormones, pheromones, and antibiotics [14,68-72]. Therefore the synthesis of these natural products taken up in an elegant way using lipase mediated kinetic resolution of propargyl alcohol (to create the stereo center of the target molecule) followed by Alder-Rickert reaction (to construct the functionalized aromatic precursor) as the key steps. The synthesis of target molecules, Paecilocin A and (-)-3-butyl-7-hydroxyphthalide were accomplished by the transesterification of racemic compounds respectively, which were easily synthesized by alkyne esters and anisole derivative of diene via Alder-Rickert reaction [73]. The optically pure esters were formed from secondary alcohols via the Enzymatic kinetic trans-esterification of racemic propargylic alcohols using Novozym-435 in the presence of vinyl acetate in tert-butyl methyl ether afforded enantioselective (S)-acetates. Propargylic alcohols were synthesized from the commercially available aldehydes respectively [74]. In summary, we have successfully demonstrated the enantioselective total synthesis of Paecilocin A & (-)-3-butyl-7-hydroxyphthalide using lipase mediated kinetic resolution of racemic propargyl alcohol and Alder-Rickert reaction. The synthesized chiral molecules possess a potent anti-fungal activity.

Synthesis of (R)-Dodecan-4-olide & (R)-Octan-4-olide Molecules

This Section describes the total synthesis of γ -lactones; optically active 5- and 6-alkyl-substituted (or) γ - and δ -lactones which are important scaffolds in medicinal chemistry and also attractive building blocks in the synthesis of natural products. Lactones contain structural moieties that are frequently present in insect pheromones, cardenolides, lignans and flavor components [16-18]. Among various γ -lactones, 4(R)-Dodecanolide is a unique and naturally occurring butanolide; it is isolated from fruits and butterfat. It is a defensive secretion of the pygidial glands of rove beetles, Bledius mandibularis and Bledius spectabilis and is also produced during the bioconversion of soya bean fatty acids by Penicillium roqueforti spores in the presence

of an exogenous lipase. These lactones were used as a flavouring agent. The γ -lactones, (4R)-Octanolide is found in strawberry, peach and apricot, it exhibits significant physiological activities [19,75]. The synthesis of target molecules, γ -lactones were accomplished from alkyne esters respectively, which were easily prepared from secondary alcohols via the enzymatic kinetic resolution of racemic propargylic alcohols using Novozym-435 in the presence of vinyl acetate in tert-butyl methyl ether afforded (R)-alcohols. Propargylic alcohols were synthesized from the commercially available aldehydes respectively.

Summary

The current review successfully demonstrated chemo enzymatic total synthesis of wide variety of chiral alcohols such as aromatic and aliphatic chiral alcohols via bio-reduction and synthesis of γ -lactones such as (4R)-Dodecanolide and (4R)-Octanolide using lipase mediated kinetic resolution followed by lactonization as the key steps.

Acknowledgment

Dr. Sreelakshmi Chittamuru says thanks to Professor Ram Gudavalli and Professor William Reed at Palmer Center for Chiropractic Research Davenport USA, for giving the opportunity as a visiting researcher at Biomechanics core department and CSIR, New Delhi, India for financial support in the form of a fellowship from 2007-2013 (financial support as a part of XII five year plan program under title ORIGIN (CSC-0108). And also thanks for Dr. A.B.R. and Dr. B.V.S.R IICT Hyderabad India for their continuous support.

References

1. Pasteur LCR (1853) Transformation of Tartaric Acid into Racemic Acid. Academie des Sciences 37: 162.
2. Mislow K, Siegel J (1984) Stereoisomerism and local chirality. J Am Chem Soc 106: 3319-3328.
3. Faul MM, Huff BE (2000) Strategy and Methodology Development for the Total Synthesis of Polyether Ionophore Antibiotics. Chem Rev 100: 2407-2474.
4. Babine RE, Bender SL (1997) Molecular Recognition of Protein-Ligand Complexes: Applications to Drug Design. Chem Rev 97: 1359-1472.
5. Garg R, Gupta S, Gao H, Babu MS, Debnath A, et al. (1999) Comparative Quantitative Structure-Activity Relationship Studies on Anti-HIV Drugs. Chem Rev 99: 3525-3602.
6. Andrew SW, Gregory LF, Peter CM, John WW (1986) The use of enzymes in the manufacture of active pharmaceutical ingredients & drug quality Wells. A S Organic Process Research & Development 16: 1986-1993.
7. Rakeshwar B, Apurba B, Louis D, Peter D, Kenneth F, et al. (2012) Green Chemistry articles of interest to the pharmaceutical industry. Organic Process Research & development 16: 1887-1896.
8. Routhi AM (2002) Chem Eng pp43.
9. Wanda MK, Mironowicz A (2004) Tetrahedron: Asymmetry 13: pp1965.
10. Challenger CA (2001) Chiral Drugs John Willey & Sons New York.
11. Gotor V, Alfonso I, Urdiales GE (2008) Asymmetric Organic Synthesis with Enzymes. Wiley Weinheim.
12. Yale HY (1961) Pyridine and Its Derivatives: part 2 In: Klingsberg E Interscience Publisher INC New York.
13. Hayoz P, Von ZA, Evans SH (1993) Stereoselective synthesis of octahedral complexes with predetermined helical chirality. Journal of American Chem Society 115: 5111-5114.
14. Kawano S, Horikawa M, Yasohara Y, Hasegawa J (2003) Bioscience Biotechnology and Bio-chemistry 67: pp 809.
15. Cordell GA, Lemos TLG, Monte FJQ, De Mattos MCJ (2007) Nat Prod 70: pp478.

16. Blanchard N, Weghe PVD (2006) *Daucus carota* L. mediated bioreduction of prochiral ketones. *Org Biomol Chem* 4: 2348-2353.
17. Akakabe Y, Takahashi M, Kamezawa M, Kikuchi K, Tachibana H (1995) Preparation of chiral enantiopure 2-(hydroxyalkyl)pyridine derivatives. Use of the chiral pool. *Chem Soc Perkin Trans 1*: 1983-1990.
18. Baskar B, Ganesh S, Lokeswari TS, Chadka A (2004) Highly stereoselective reduction of 4-Aryl-2-oxo but-3-enoic carboxylic esters by plant cell culture of *Daucus carota*. *J Mol Catal B: Enzym* 27: 13-17.
19. Baldassarre F, Bertoni G, Chiappe C, Marioni F (2000) Preparative synthesis of chiral alcohols by enantioselective reduction with *Daucus carota* root as biocatalyst. *J Mol Catal B: Enzym* 11: 55-58.
20. Yadav JS, Reddy PT, Hashim SR (2000) *Synlett* pp1049.
21. Yadav JS, Reddy PT, Hashim SR (2000) *Synlett* pp473.
22. Maczka WK, Mironowicz A (2002) Enantioselective hydrolysis of 1-aryl ethyl acetates and reduction of aryl methyl ketones using carrot, celeriac and horseradish enzyme systems. *Tetrahedron: Asymmetry* 13: 2299-2302.
23. Stampfer W, Kosjek B, Faber K, Kroutil W (2003) Biocatalytic Asymmetric Hydrogen Transfer Employing *Rhodococcus ruber* DSM 44541. *J Org Chem* 68: 402-406.
24. Gröger H, Hummel W, Rollmann C, Chamouveau F, Hüsken H, et al. (2004) Preparative asymmetric reduction of ketones in a biphasic medium with an (S)-alcohol dehydrogenase under in situ-cofactor-recycling with a formate dehydrogenase. *Tetrahedron* 60: 633-640.
25. Edegger K, Stampfer W, Seisser B, Faber K, Mayer SF (2006) Regio- and Stereoselective Reduction of Diketones and Oxidation of Diols by Biocatalytic Hydrogen Transfer. *J Org Chem* 8: 1904-1908.
26. Yang ZH, Zeng R, Yang G, Wang Y, Li LZ (2008) Asymmetric reduction of prochiral ketones to chiral alcohols catalyzed by plants tissue. *J Ind Microbiol Biotechnol* 35: 1047-1051.
27. Caron D, Coughlan AP, Simard M, Bernier J, Piche Y, et al. (2005) Stereoselective reduction of ketones by *Daucus carota* hairy root cultures. *Biotech Lett* 27: 713-716.
28. Scarpi D, Occhiato EG, Guarna A (2005) Selectivity of *Daucus carota* roots and baker's yeast in the enantioselective reduction of γ -nitroketones. *Tetrahedron: Asymmetry* 16: 1479-1483.
29. Maczka WK, Mironowicz A (1965) Enantioselective reduction of bromo- and methoxy-acetophenone derivatives using carrot and celeriac enzymatic system. *Tetrahedron: Asymmetry* 15: 1965-1967.
30. Comasseto JV, Omori AT, Porto ALM, Andrade LH (2004) Preparation of chiral organochalcogeno- α -methylbenzyl alcohols via biocatalysis. The role of *Daucus carota* root *Tetrahedron Lett* 45: 473-476.
31. Kumaraswamy G, Ramesh S (2003) Soaked *Phaseolus aureus* L: an efficient biocatalyst for asymmetric reduction of prochiral aromatic ketones. *Green Chem* 5: 306-308.
32. Danielle Skropeta (2008) Deep-sea natural products. *Natural Product Reports* 25: 1131-1166.
33. Shaw JT (2009) Naturally diverse: highlights in versatile synthetic methods enabling target- and diversity-oriented synthesis. *Nat Prod Rep* 26: 11-26.
34. Blunt JW, Copp BR, Hu WP, Munro MHG, Northcote PT, et al. (2009) Marine natural products. *Nat Prod Rep* 26: 144-222.
35. Jin Z (2009) Muscarine, imidazole, oxazole and thiazole alkaloids. *Nat Prod Rep* 26: 382-445.
36. Cragg GM, Grothaus PG, Newman DJ (2009) Impact of natural products on developing new anti-cancer agents. *Chem Rev* 109: 3012-3043.
37. Midland MMJ (1975) Preparation of monolithium acetylde in tetrahydrofuran. Reaction with aldehydes and ketones. *Org Chem* 40: 2250-2252.
38. Duffault JM, Einhorn J, Alexakis A (1995) Synthesis of regioselectively deuterated hydroxy conjugated dienes. *Tetrahedron Lett* 36: 2765-2768.
39. Rao YS (1976) Recent advances in the chemistry of unsaturated lactones *Chem Rev* 1 76: 625-694.
40. Fukusaki E, Senda S, Nakazono Y, Omata T (1991) Lipase-catalyzed kinetic resolution of methyl 4-hydroxy-5-tetradecynoate and its application to a facile synthesis of Japanese beetle pheromone. *Tetrahedron* 47: 6223-6230.
41. Mori K (1989) Synthesis of optically active pheromones. *Tetrahedron* 45: 3233-3298.
42. Wheeler JW, Happ GM, Araujo J, Pasteels JM (1972) γ -Dodecalactone from rove beetles. *Tetrahedron Lett* 13: 4635.
43. Bloch R, Gilbert L (1987) Synthesis of both enantiomers of γ -substituted α,β -unsaturated γ -lactones. *J Org Chem* 52: 4603-4605.
44. Gunasekera SP, Mc Carthy PJ, Kelly-Borges M, Lobkovsky E, Clardy J (1996) Dysidiolide: A Novel Protein Phosphatase Inhibitor from the Caribbean Sponge *Dysidea etheria* de Laubenfels. *J Am Chem Soc* 118: 8759-8760.
45. Avcibasi H, Anil H (1987) Four terpenoids from *Cedrus libanotica*. *Phytochemistry* 26: 2852-2854.
46. Miles DH, Chittawong V, Lho DS, Payne AM, de La Cruz AA, et al. (1991) Toxicants from Mangrove Plants, VII. Vallapin and Vallapianin, Novel Sesquiterpene Lactones from the Mangrove Plant *Heritiera littoralis*. *J L J Nat Prod* 54: 286-289.
47. Hoffmann HMR, Kover A, *Angew J* (1985) *Chem Int Ed Engl* 24: 94.
48. Boukoulas J, McCann LC (2011) Concise, regiocontrolled synthesis of yanginhualine A. *Tetrahedron Lett* 52: 1202-1204.
49. Hentzer M, Eberl L, Nielsen J, Givskov M (2003) Quorum Sensing: A Novel Target for the Treatment of Biofilm Infections. *Biodrugs* 17: 241-250.
50. Hjelmggaard T, Persson T, Rasmussen TB, Givskov M, Nielsen J (2003) *Bioorg Med Chem* 11: 3261-3271.
51. Fusetani N (2004) Biofouling and antifouling. *Nat Prod Rep* 21: 94-104.
52. Cateni F, Zillic J, Zacchigna M, Bonivento P, Frausin F et al. (2006) Synthesis and biological properties of new α -methylene- γ -butyrolactones and α,β -unsaturated δ -lactones. *Eur J Med Chem* 41: 192-200.
53. Marcos IS, Escola MA, Moro RF, Basabe P, Diez D, et al. (2007) Synthesis of novel antitumoural analogues of dysidiolide from ent-halimic acid. *Bioorg Med Chem* 15: 5719-5737.
54. Takahashi M, Dodo K, Sugimoto Y, Aoyagi Y, Yamada Y, et al. (2000) Synthesis and biological properties of novel analogues of dysidiolide and their structure-activity relationship. *Bioorg Med Chem Lett* 10: 2571-2574.
55. Brohm D, Philippe N, Metzger S, Bhargava A, Muller O (2002) Solid-Phase Synthesis of Dysidiolide-Derived Protein Phosphatase Inhibitors *J Am Chem Soc* 124: 13171-13178.
56. Li DH, Zhu TJ, Liu HB, Fang CY, Gu QQ (2006) Four butenolides are novel cytotoxic compounds isolated from the marine-derived bacterium, *Streptovorticillium luteovorticillatum* 11014. *Arch Pharm Res* 29: 624-626.
57. Ma S, Shi Z, Yu Z (1999) Synthesis of β -halobutenolides and their Pd(0)-catalyzed cross-coupling reactions with terminal alkynes and organozinc reagents. A general route to β -substituted butenolides and formal synthesis of cis-whisky lactone. *Tetrahedron* 55: 12137-12148.
58. Chalier P, Crouzet J (1998) Enantiodifferentiation of four γ -lactones produced by *Penicillium Roqueforti*. *Chirality* 10: 786-790.
59. Flath RA, Black DR, Guadagni DG, McFadden WH, Schultz TH (1967) *J. Agric. Food Chem* 15: 29.
60. Shimotori Y, Miyakoshi T (2010) Combination of Novozym 435-catalyzed hydrolysis and Mitsunobu reaction for production of (R)- γ -lactones. *Synth Commun* 40: 1607.
61. Habel A, Boland W (2008) Efficient and flexible synthesis of chiral γ - and δ -lactones. *Org. Biomol. Chem* 6: 1601-1604.
62. Gansauer A, Fan CA, Keller F, Keil JJ (2007) Titanocene-Catalyzed Regiodivergent Epoxide Openings. *Am Chem Soc* 129: 3484-3485.
63. Pollock JA, Clark KM, Martynowicz BJ, Pridgeon MG, Rycenga MJ, et al. (2007) A mild biosynthesis of lactones via enantioselective hydrolysis of hydroxynitriles. *Tetrahedron: Asymmetry* 18: 1888-1892.

64. Sabitha G, Reddy EV, Yadagiri K, Yadav JS (2006) Total synthesis of insect pheromones (R)-4-dodecanolide and (S)-5-hexadecanolide. *Synthesis* 19: 3270.
65. Taylor S, Arnold C, Gerds A, Ide N, Law KM, et al. (2004) Lactone synthesis via biotransformations of γ -hydroxyamides. *Tetrahedron: Asymmetry* 15: 3819-3821.
66. Benincori T, Rizzo S, Pilati T, Ponti A, Sada M, et al. (2004) Process-scale preparation of enantiomerically pure γ -lactones by asymmetric hydrogenation of γ -ketoesters and comparative tests of the sensory properties of some antipodes. *Tetrahedron: Asymmetry* 15: 2289-2297.
67. Corbett M, Liu X, Sanyal A, Synder J (2003) Cycloadditions of chiral anthracenes: effect of the trifluoromethyl group. *Tetrahedron Lett* 44: 931.
68. Movassaghi M, Jacobsen E (2002) *J Am Chem Soc* 124: 2468.
69. Krief A, Ronvaux A, Tuch A (1998) Syntheses of rac- and optically active 2-alkyl- γ -butyrolactones and 2-alkyl-cyclobutanones. *Tetrahedron* 54: 6903-6908.
70. Harcken C, Bruckner R (1997) *Angew Synthesis of Optically Active Butenolides and γ -Lactones by the Sharpless Asymmetric Dihydroxylation of β,γ -Unsaturated Carboxylic Esters.* *Int Ed Engl* 36: 2750.
71. Masuya K, Tanino K, Kuwajina I (1994) 3-(Alkylthio)-1,2-bis(siloxy)-3-butenes as efficient chirality transferred building blocks *Tetrahedron Lett* 35: 7965-7968.
72. Federici C, Chiummiento L, Funicello M (1994) Ring opening of 2,3-epoxy 1-tosylates to halohydrins and subsequent elaboration to asymmetrical alcohols. *Tetrahedron Lett* 35: 797-800
73. Ohkuma T, Kitamura M, Moyori R (1990) Enantioselective synthesis of 4-substituted γ -lactones. *Tetrahedron Lett* 31: 5509-5512.
74. Nemoto H, Ishibashi H, Mori M, Fujita S, Fukumoto K (1990) *J Chem Soc Perkin Trans pp*: 2835.
75. Utaka M, Watabu H, Takeda A (1987) Asymmetric reduction of the prochiral carbonyl group of aliphatic γ - and δ -keto acids by use of fermenting bakers' yeast. *J Org Chem* 52: 4363-4368.