

Abundance Detection and Molecular Characterization of *Toxoplasma gondii* by SAG1 Gene in Rodents and Cattle of Golestan Province, Northeast of Iran

Javid Sadraie and Ehsan Shariat Bahadory*

Medical University of Tarbiat Modares, Tehran, Iran

Abstract

Background: *Toxoplasma* parasite is from Toxoplasmatidea family that initially was seen in *ctinodactylus gondii* rodent. *Toxoplasma* parasites that extracted from different rodents are same in immunologic and morphologic characteristics but have differences in pathogenicity and genotypes in mice. The rodents and cattle are most reservoir hosts in environment that by attention of human environment vicinity to cattle and rodent's environment causes *Toxoplasma* dispersion in that area. The aim of this study was Molecular detection of *Toxoplasma gondii* in rodents and cattle of Golestan province, northeast of Iran.

Materials and methods: In this study we collected 285 mice and 185 cattle tissues from Golestan forest and extracted brains of rodents and hearts tissues of cattle to obtain DNA from these tissues. We divided these rodents to 4 groups and then detected the positive samples by PCR method.

Results: In this study we found 68 samples of these rodents were positive for SAG1 gene. 38 samples were *Ratus ratus*, 10 samples were *Ratus norvegicus*, 10 samples were *Mus musculus* and 10 samples were *Rombomys opimus*. Also 81 samples of cattle hearts were positive for SAG1 gene.

Conclusion and discussion: in this study we found that the different types of rodents were responsible to spread of toxoplasmosis, also SAG1 gene was very useful marker to detect toxoplasmosis in rodents of northeast area of Iran. Also in this area the numbers of toxoplasmosis cattle were in very high range.

Keywords: Toxoplasmosis; SAG1 gene; PCR; Golestan forest; Rodents; Cattle

Introduction

Toxoplasma gondii is an intracellular parasite that infected many hosts in Iran, including human, rodents, cats and domestic animals. Because domestic and feral cats are the natural definitive host, they play an important role in disseminating of toxoplasmosis [1-5]. Rodents are very important reservoir in dissemination of toxoplasmosis in Iran. The brain of rodents and the heart of cattle are the most important tissues that can infect by toxoplasmosis. Toxoplasmosis infection causes tissue cysts in these organs of rodents and cattle. The major gene that extracted from these organs to detect toxoplasmosis is SAG1 gene [5-10].

Toxoplasmosis infection pathway is eating of infected tissues from rodents by cats or uncooked meats from domestic animals to human, then laying Oocysts from cats in environments to dissemination of infection to human or domestic animals. The major disease of this parasite in human is encephalitis or brain disorders [11-13]. Primary routes of acute human *T. gondii* infection include ingestion of tissue cysts in undercooked, contaminated meat, congenital infection through the placenta, and ingestion of Oocysts from soil, water, or cat litter. Oocysts are shed in cat feces and can remain viable in soil and water samples for months to years. The infected rodents are the main food for cats, and this cycle is the important cause of spreading toxoplasmosis infection. My purpose was the abundance detection of toxoplasmosis in rodents and cattle of Golestan province using SAG1 gene in brain and heart tissues [14-18].

Materials and Methods

Different regions of Golestan province have different climate and are notably heterogeneous. Northern parts are located in the arid and semi-arid climate, southern parts show a mountainous climate, and

central and southern west parts have a moderate Mediterranean climate (Weather Centre Hashem). 185 cattle were collected from 20 villages of 5 towns locating in different climates of Golestan province. Also rodents were collected from Golestan forest. This study was done on March to September 2016. Because of this study was based on molecular detection; we did not use other examinations for example histopathology examinations.

Preparation of samples

We collected 285 mice from Golestan forest and 185 cattle hearts from 20 villages. Then brains and hearts of mice and cattle were removed. These organs were fixed in 95% Ethanol, and preserved in 4°C until DNA extraction. These rodents divided in 4 groups: (*Rattus rattus*, *Rattus norvegicus*, *Mus musculus* and *Rombomys opimus*) (Table 1).

DNA extraction

To extraction genomic DNA, we cut approximately 3 g of brain and heart tissues by DNG/proteinase K method from Sinacolon company and eluted into 50 µl DDH₂O according to the manufacturer's

*Corresponding author: Ehsan Shariat Bahadory, PhD of Medical Parasitology, Medical University of Tarbiat Modares, Tehran, Iran, Tel: 0098912418960; E-mail: ehsanshariat63@gmail.com

Received April 22, 2017; Accepted June 12, 2017; Published June 14, 2017

Citation: Sadraie J, Bahadory ES (2017) Abundance Detection and Molecular Characterization of *Toxoplasma gondii* by SAG1 Gene in Rodents and Cattle of Golestan Province, Northeast of Iran. J Vet Sci Technol 8: 446. doi: [10.4172/2157-7579.1000446](https://doi.org/10.4172/2157-7579.1000446)

Copyright: © 2017 Sadraie J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Rodents type	Numbers
<i>Rattus rattus</i>	130
<i>Rattus norvegicus</i>	45
<i>Mus musculus</i>	60
<i>Rombomys opimus</i>	50

Table 1: The total samples of rodents divided in 4 groups [14-18].

recommendations. A PCR targeting the *T. gondii* SAG1 gene was performed to detect possible infection with *T. gondii*. DNA samples giving positive SAG1 amplification were then used for genetic characterization.

We cut 3 g of tissue into small places and placed the samples into a 1.5 ml sterile tube. Then added 180 µl (microliter) lyse buffer (pepsin) to homogenization and 400 microliter (DNG/proteinase K) and then homogenized the samples. If the sample size was larger than 3 g we should increase the amount of lyse buffer proportionally.

Added 20 µl proteinase K to the samples. Then mixed immediately by shaking for 20 seconds. Then incubated at 60°C for 1 hour to lyse samples. If tissue was difficult to lyse, we increased the incubation time to 2-3 hours. Then inverted the samples every 10-15 minutes. Then we added 300 microliter isopropanol to DNA precipitation. After 5-10 minutes we washed the tubes by 70% ethanol and finally we eluted the DNA by DDH2O. The eluted DNA preserved in -20°C freezer until using PCR method to detection of infection. The purification of DNA was done with Nanodrop instrument.

PCR analysis for *T. gondii* SAG1 gene

To genetically identify the presence of KI-1 Tachyzoites in visceral organs, PCR analysis was performed to detect *Toxoplasma* SAG1 gene as a diagnostic gene. DNA extraction was performed using the DNeasy[®]Tissue kit (DNG Sinacolon). The primers were designed by BLAST method from NCBI site and produced by Pishgam Company.

Primer sequences were:

5'- GCTGTAACATTGAGCTCCTTGASTTCCTG-3' for forward and

5'- CCGGAACAGTACTGATTGTTGTCTTGAG-3' for reverse

Amplification of the SAG1 gene was completed in the following conditions: 1 cycle of 5 min at 95°C for initial denaturation followed by 30 cycles of 1 min at 95°C, 1 min at 62°C, and 3 min at 74°C. The best annealing temperature was 62°C. Amplification was performed using a DNA thermal cyler (Eppendorf instrument). PCR amplification products were examined in 1.5% agarose gels and confirmed by staining with Safe stain and visualized under Gel Doc using UV. The statistical surveys were done with SPSS 18 software (Table 2).

Results

In this study we found 68 samples of these rodents were positive for SAG1 gene. 38 samples were *Rattus rattus*, 10 samples were *Rattus norvegicus*, 10 samples were *Mus musculus* and 10 samples were *Rombomys opimus*. The samples were positive in 1180 bp location. Also 81 heart samples of cattle were positive for SAG1 gene. *Toxoplasma gondii* SAG1 gene after NCBI blast, complete cds (Table 3; Figures 1 and 2).

Origin:

1 acaatgtgca cctgtaggaa gctgtagtca ctgctgattc tcactgttct cggaagggc

61 cgacgaccgg agtacagttt ttgtggcag agccgttggt cagctttccg ttgttctcgg

Materials	Volume	Final concentration
Master Mix	7.5 µl	2X
DNA Sample	2 µl	10-100 ng/µl
Work Primer	1 µl	20 pmol
Distilled water	4.5 µl	-
Total	15.0 µl	-

Table 2: PCR substrates [14-18].

Rodents type	Numbers	Numbers of positive
<i>Rattus rattus</i>	130	38
<i>Rattus norvegicus</i>	45	10
<i>Mus musculus</i>	60	10
<i>Rombomys opimus</i>	50	10

Table 3: Total samples and positive samples for SAG1 gene [14-18].




Figure 1: PCR conclusion for SAG1 gene. P: Positive control. M: Size marker. S: Sample. N: Negative control [14-18].




Figure 2: PCR conclusion for SAG1 gene. P: Positive control. M: Size marker. S: Sample. N: Negative control [14-18].

121 ttgtgtcaca tgtgtcattg tcgtgtaaac acacggttgt atgtcggttt cgctgcacca
 181 cttcattatt tcttctggtt tttgacgag tatgtttccg aaggcagtga gacgcgccgt
 241 cacggcaggg gtgtttgccg cgcccact gatgtcgttc ttgcgatgtg gcttatggc
 301 atcggatccc cctcttggc ccaatcaagt tgcactctgc ccagataaaa aatcgacagc
 361 cgcggtcatt ctacaccga cggagaacca cttactctc aagtgccta
 aaacagcgt
 421 cacagagcct cccactcttg cgtactacc caacaggcaa atctgccag
 cgggtactac
 481 aagtactgtg acatcaaagg ctgtaacatt gagtccttg atctctgaag cagaagatag
 541 ctggtggacg ggggattctg ctagtctga cacggcaggc atcaaactca
 cagttcaat
 601 cgagaagttc cccgtgaca cgcagacgtt tgtggtcgtg tgcataagg
 gagacgacgc

661 acagattgt atggtcacag tgacagtaca agccagagcc tcatcggctc
tcaataatgt

721 cgcaaggtgc tctcaggtg cagacagcac tcttggctct gtcaagttgt
ctgcggaagg

781 acccactaca atgaccctcg tgtgcgggaa agatggagtc aaagttctc
aagacaaca

841 tcagttactgt tccgggacga cgctgactgg ttgcaacgag aaatcgttca aagatattt

901 gccaaaatta actgagaacc cgtggcaggg taacgcttcg agtgataagg
gtgccacgct

961 aacgatcaag aaggaagcat ttccagccga gtcaaaaagc gtcattattg
gatgcacagc

1021 gggatgcct gagaagcatc actgtaccgt gaaactggag ttgcccgggg
ctgcagggtc

1081 agcaaaatcg gctgcgggaa cagccagtcg cgtttccatt cttgcatgg
tgatcggact

1141 tattggctct atcgagcgtt gttgcgctg agtgattacc gttgtgc

These results showed that positive samples in 1180 bp location by SAG1 gene. These results obtained from GelDock instrument. Primer designation was done by BLAST software from NCBI site. In 130 samples from *Rattus rattus* 38 samples were positive, in 45 samples from *Rattus norvegicus* 10 samples were positives, in 60 samples from *Mus musculus* 10 samples were positive and finally in 50 samples from *Rombomys opimus* 10 samples were positives. Also 81 samples of heart tissues of cattle were positive for toxoplasmosis.

Figure 3 showed that 1=*Rattus rattus*, 2=*Rattus norvegicus*, 3=*Mus musculus* and 4=*Rombomys opimus*. Series 1 (blue) were total sample and series 2 (red) were positive samples. Positive samples bands detected on 1180 bp region [14-18]. These results obtained from GelDock instrument (Figure 4). Figure 5 showed that in the 185 heart tissues of cattle, 85 samples were positive for *Toxoplasma* SAG1 gene.

Discussion

Regarding to free living of rodents, cattle and also presence of them in large number in rural areas, obtaining data about *T. gondii* dynamic in rodents and cattle population of rural area is critical for the establishment of monitoring programs [19-21]. My purpose of this study was abundance detection of toxoplasmosis in rodents and cattle of Golestan province using SAG1 gene. Toxoplasmosis is a zoonosis infection between rodents, cats, human and domestic animals. Rodents are main reservoir host for toxoplasmosis to infect feral cats and uncooked meat of domestic animals such as cattle or sheep can transfer toxoplasmosis to human. In both ways the human can plague with toxoplasmosis. In Golestan province, the population of rodents is higher than cattle but both animals have same role in spreading of toxoplasmosis. Climate characteristic of Golestan province is optimum for dissemination of toxoplasmosis. Prevalence of 40% *T. gondii* antibodies in stray cats in Sari, Northern Iran, has been reported by Sharif et al. They also survey anti *T. gondii* antibodies with latex agglutination test (LAT) on 100 serum samples collected from stray cats in five urban areas of Sari. Sari is located near Golestan province in North Iran and has humid climate which has been introduced suitable for *T. gondii* growth [22-25]. A study in Tabriz by Jamali clarified 36/2% *T. gondii* infection of cats by using dye test that differed with our methods. In this study most positive sampled has been belonged to Azadshahr villages and also Gorgan villages. Compare to Bandar-Turkmen,






Figure 4: Positive samples of cattle heart tissues. M: Size marker. S: Samples. N: Negative control. P: Positive control.



Gonbad and Aghghala that have semi-arid and arid climates, these two regions have humid climate. So detection of higher positive sample in Azadshahr villages and also Gorgan villages can be due to more suitable condition for growth of *T. gondii* oocyst in these areas. Also Mostafavi et al. reported highest prevalence of human toxoplasmosis, 70%, in humid regions of North Iran. Prevalence of *Toxoplasma gondii* antibodies in cats from Urmia, Northwest of Iran. In Sari by contrast, differences in *T. gondii* infection were detected between male stray cats and female stray cats. In 2013, Cong detected house sparrows toxoplasmosis in China, Khademvatan detected birds toxoplasmosis in southwest of Iran and Ortega found pigs toxoplasmosis in Mexico [22-26]. Most of the

studies didn't reported significant different in *T. gondii* infection of two sexes and the role of sexuality in *T. gondii* exposure is not clear. In recent years had not any studies in rodents toxoplasmosis in Golestan area. In 2011, Hong did very important study in genotyping of cat's toxoplasmosis. In 2011, Dubey done a study in genotyping of zoonosis toxoplasmosis in USA. In 2012, Selseleh did a study in genotyping of Tehran rodent's toxoplasmosis by SAG1 gene. In 2012, Habibi did very important study in detection of sheep toxoplasmosis by SAG1 gene. In 2013, Ling jang did immense study in detection of rodent toxoplasmosis by SAG1 gene in china. In 2013, Cabral did a study in detection of rodent toxoplasmosis in Brazil. In 2014, Barros et al. had found genetic characteristics of *Toxoplasma gondii* from doves in Brazil. In 2014, Yan et al. found genetic characteristics of *Toxoplasma gondii* from rodents in china. In 2014, Gjerede et al. had found genetic characteristics of *Toxoplasma gondii* from muscles of Lutra in Norway. In 2014, Chen et al. detected *Toxoplasma gondii* from HIV positive people in China [26-30]. In recent years have not perform any studies to detect rodents and cattle toxoplasmosis at the same time in Golestan area. This study showed important role of rodents and cattle in dissemination of toxoplasmosis in humid area. In this study we showed that the SAG1 gene was very important marker to detection of abundance of zoonotic toxoplasmosis and brain or heart tissue was main tissues to follow SAG1 gene from Tachyzoites of *Toxoplasma* parasite, also the rodents and cattle were very important reservoirs in dissemination of toxoplasmosis to human in Golestan area, northeast of Iran [26-30].

Conclusion

In this study we concluded: The northeast of Iran was very important region for following toxoplasmosis infection, specifically in animals such as rodents and cattle. In other study that had done four month after this surveillance, we found that the dominant genotype of *Toxoplasma gondii* in that area was genotype using PCR-RFLP by HaeII restriction enzyme.

Acknowledgements

The current research was funded by Tarbiat Modares University.

References

- Evers F, Garcia JL, Navarro IT, Zulpo DL, Nino BS, et al. (2013) Diagnosis and isolation of *Toxoplasma gondii* in horses from Brazilian slaughterhouses. Rev Bras Parasitol Vet 22: 58-63.
- Quan JH, Zhou W, Cha GH, Choi IW, Shin DW, et al. (2013) Kinetics of IL-23 and IL-12 secretion in response to *Toxoplasma gondii* antigens from THP-1 monocytic cells. Korean J Parasitol 51: 85-92.
- Dubey JP, Hill D, Zarlenga D, Choudhary S, Ferreira LR, et al. (2013) Isolation and characterization of new genetic types of *Toxoplasma gondii* and prevalence of *Trichinella murrelli* from black bear (*Ursus americanus*). Vet Parasitol 196: 24-30.
- Assmar M, Yassaie F, Terhovanesian A, Esmaeili AR, Nahrevanian H, et al. (2004) Prenatal Diagnosis of Congenital Toxoplasmosis Validity of PCR Using Amniotic Fluid against Indirect Fluorescent antibody assay in mothers. Iranian J Public Health 33: 1-4.
- Pena HF, Vitaliano SN, Beltrame MA, Pereira FE, Gennari SM, et al. (2013) PCR-RFLP genotyping of *Toxoplasma gondii* from chickens from Espírito Santo state, Southeast region, Brazil: new genotypes and a new SAG3 marker allele. Vet Parasitol 192: 111-117.
- Quan JH, Kim TY, Choi UI, Lee YH (2008) Genotyping of a Korean isolate of *Toxoplasma gondii* by multilocus PCR-RFLP and microsatellite analysis. Korean J Parasitol 46: 105-108.
- Hong SH, Jeong YI, Kim JY, Cho SH, Lee WJ, et al. (2013) Prevalence of *Toxoplasma gondii* infection in household cats in Korea and risk factors. Korean J Parasitol 51: 357-361.
- Dubey JP, Randall AR, Choudhary S, Ferreira LR, Verma SK, et al. (2013) Occurrence, isolation, and genetic characterization of *Toxoplasma gondii* from white-tailed deer (*Odocoileus virginianus*) in New Jersey. J Parasitol 99: 763-769.
- Selseleh M, Modarressi MH, Mohebbali M, Shojaee S, Eshragian MR, et al. (2012) Real-time RT-PCR on SAG1 and BAG1 gene expression during stage conversion in immunosuppressed mice infected with *Toxoplasma gondii* Tehran strain. Korean J Parasitol 50: 199-205.
- Habibi G, Imani A, Gholami M, Hablolvarid M, Behroozkikah A, et al. (2012) Detection and Identification of *Toxoplasma gondii* Type One Infection in Sheep Aborted Fetuses in Qazvin Province of Iran. Iran J Parasitol 7: 64-72.
- Selseleh MM, Keshavarz H, Mohebbali M, Shojaee S, Modarressi M, et al. (2012) Production and Evaluation of *Toxoplasma gondii* Recombinant Surface Antigen 1 (SAG1) for Serodiagnosis of Acute and Chronic *Toxoplasma* Infection in Human Sera. Iran J Parasitol 7: 1-9.
- Lim H, Lee SE, Jung BK, Kim MK, Lee MY, et al. (2012) Serologic survey of toxoplasmosis in Seoul and Jeju-do, and a brief review of its seroprevalence in Korea. Korean J Parasitol 50: 287-293.
- Lee SE, Hong SH, Lee SH, Jeong YI, Lim SJ, et al. (2012) Detection of ocular *Toxoplasma gondii* infection in chronic irregular recurrent uveitis by PCR. Korean J Parasitol 50: 229-231.
- El Behairy AM, Choudhary S, Ferreira LR, Kwok OC, Hilali M, et al. (2013) Genetic characterization of viable *Toxoplasma gondii* isolates from stray dogs from Giza, Egypt. Vet Parasitol 193: 25-29.
- Edwards JF, Dubey JP (2013) *Toxoplasma gondii* abortion storm in sheep on a Texas farm and isolation of mouse virulent atypical genotype *T. gondii* from an aborted lamb from a chronically infected ewe. Vet Parasitol 192: 129-136.
- Dubey JP, Choudhary S, Kwok OC, Ferreira LR, Oliveira S, et al. (2013) Isolation and genetic characterization of *Toxoplasma gondii* from mute swan (*Cygnus olor*) from the USA. Vet Parasitol 195: 42-46.
- Herrmann DC, Wibbelt G, Götz M, Conraths FJ, Schares G (2013) Genetic characterisation of *Toxoplasma gondii* isolates from European beavers (*Castor fiber*) and European wildcats (*Felis silvestris silvestris*). Vet Parasitol 191: 108-111.
- Lilly EL, Wortham CD (2013) High prevalence of *Toxoplasma gondii* oocyst shedding in stray and pet cats (*Felis catus*) in Virginia, United States. Parasit Vectors 6: 266.
- Zhang XX, Lou ZZ, Huang SY, Zhou DH, Jia WZ, et al. (2013) Genetic characterization of *Toxoplasma gondii* from Qinghai vole, Plateau pika and Tibetan ground-tit on the Qinghai-Tibet Plateau, China. Parasit Vectors 6: 291.
- Cabral AD, Gama AR, Sodré MM, Savani ES, Galvão-Dias MA, et al. (2013) First isolation and genotyping of *Toxoplasma gondii* from bats (Mammalia: Chiroptera). Vet Parasitol 193: 100-104.
- Mancianti F, Nardoni S, D'Ascenzi C, Pedonese F, Mugnaini L, et al. (2013) Seroprevalence, detection of DNA in blood and milk, and genotyping of *Toxoplasma gondii* in a goat population in Italy. Biomed Res Int 2013: 905326.
- Ortega-Pacheco A, Acosta Viana KY, Guzmán-Marín E, Segura-Correa JC, Alvarez-Fleites M, et al. (2013) Prevalence and risk factors of *Toxoplasma gondii* in fattening pigs farm from Yucatan, Mexico. Biomed Res Int 2013: 231497.
- Galván-Ramírez ML, Sánchez-Orozco LV, Rodríguez LR, Rodríguez S, Roig-Melo E, et al. (2013) Seroepidemiology of *Toxoplasma gondii* infection in drivers involved in road traffic accidents in the metropolitan area of Guadalajara, Jalisco, Mexico. Parasit Vectors 6: 294.
- Shichao X, Chen Z, Lei H, Tongyao W, Liusong N, et al. (2013) DNA Detection of *Toxoplasma gondii* with a Magnetic Molecular Beacon Probe via CdTe@Ni Quantum Dots as Energy Donor. J Nanomater 2013: 473703.
- Cong W, Huang SY, Zhou DH, Zhang XX, Zhang NZ, et al. (2013) Prevalence and genetic characterization of *Toxoplasma gondii* in house sparrows (*Passer domesticus*) in Lanzhou, China. Korean J Parasitol 51: 363-367.
- Khademvatan S, Saki J, Yousefi E, Abdzadeh R (2013) Detection and genotyping of *Toxoplasma gondii* strains isolated from birds in the southwest of Iran. Br Poult Sci 54: 76-80.
- Barros LD, Taroda A, Zulpo DL, Cunha IA, Sammi AS, et al. (2014) Genetic characterization of *Toxoplasma gondii* isolates from eared doves (*Zenaidura macroura*) in Brazil. Rev Bras Parasitol Vet 23: 443-448.
- Yan C, Liang LJ, Zhang BB, Lou ZL, Zhang HF, et al. (2014) Prevalence and genotyping of *Toxoplasma gondii* in naturally-infected synanthropic rats (*Rattus norvegicus*) and mice (*Mus musculus*) in eastern China. Parasit Vectors 7: 591.

29. Gjerde B, Josefsen TD (2014) Molecular characterisation of *Sarcocystis lutrae* n. sp. and *Toxoplasma gondii* from the musculature of two Eurasian otters (*Lutra lutra*) in Norway. Parasitol Res 114: 873-886.
30. Chen LJ, Jia YX, Leng L, Luo M, Gao J, et al. (2014) Comparison of *Toxoplasma gondii* separated from HIV-positive people and RH strain GRA6 gene. Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi 26: 434-436.

Citation: Sadraie J, Bahadory ES (2017) Abundance Detection and Molecular Characterization of *Toxoplasma gondii* by SAG1 Gene in Rodents and Cattle of Golestan Province, Northeast of Iran. J Vet Sci Technol 8: 446. doi: [10.4172/2157-7579.1000446](https://doi.org/10.4172/2157-7579.1000446)

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission/