

Alcoholism: Common and Oxidative Damage Biomarkers

Saé Muñiz-Hernández¹, Jesús B Velázquez-Fernández², José Díaz-Chávez¹, Rosa C López-Sánchez³, José A Hernández³ and Adela Rendón-Ramírez^{4*}

¹Subdirección de Investigación Básica, Instituto Nacional de Cancerología, Secretaría de Salud Mexico (SS), USA

²Laboratorio de Bioquímica Toxicológica, Posgrado en Ciencias Biológico Agropecuarias y Pesqueras, Universidad Autónoma de Nayarit, USA

³Tecnológico de Monterrey, Escuela de Medicina y Ciencias de la Salud, USA

⁴Unidad de Biofísica (Centro Mixto CSIC, UPV/EHU), Universidad del País Vasco, Spain

*Corresponding author: Adela Rendón Ramírez, Unidad de Biofísica (CSIC-UPV/EHU) - Fundación Biofísica Bizkaia, Aptdo 644, 48080 Bilbao, España, Spain, Tel: +34647 496 436; E-mail: adela_rendon1@ehu.es

Received date: Mar 03, 2014; Accepted date: Apr 28; 2014, Published date: Apr 30, 2014

Copyright: © 2014 Saé Muñiz-Hernández, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

For many years it has been assumed that moderate alcohol intake might provide health benefits to humans, while alcohol abuse generally produces some systemic complications, even when physical and mental dependence on alcohol has not developed. Many studies have focused on distinct molecules that might serve as useful early or late biomarkers in individuals that consume important amounts of alcohol. Indeed, the diversity of molecules postulated as biomarkers is quite wide and this is the direct reflection of the broad range of mechanisms involved in the impact that alcohol has on human health, either as a result of acute or chronic intake, the latter being a cause death. In this review, we have considered most of the molecules that have been followed-up in individuals who drink large amounts of alcohol, as well as some of those that have been less intensely studied, and we present a pair of oxidative biomarkers that could be used to evaluate early stage alcoholism.

Keywords: Alcoholism; Oxidative damage; Biomarkers

Introduction

Alcoholism represents one of the most serious socioeconomic and health problems worldwide. The American Medical Association defines alcoholism as “an illness characterized by significant impairment that is directly associated with persistent and excessive intake of alcohol”, which also results in the deterioration of social behavior and/or a physical illness, with the concomitant development of other adverse effects. Children and adolescents are more vulnerable to the harmful effects of alcohol than adults, although it is considered that this may reflect the fact that they have no context or reference point to regulate their drinking and they are building their own range of tolerance [1].

The important difference between alcohol abuse and alcoholism has often been highlighted. Accordingly, alcoholism indicates a physical and mental dependence on alcohol, which leads to regular or periodic, heavy and uncontrolled alcohol consumption [2]. By contrast, the individual who is considered to abuse alcohol has experienced severe sequels of their habit in the past year, and they may develop alcoholic liver disease and its systemic complications without developing the physical and mental dependence on alcohol [3]. Lamentably, in alcohol-dependent individuals most pathological processes precede the onset of clinical symptoms and unfortunately, many individuals with a history of chronic alcohol abuse die before reaching the clinical stage. However, if the problems provoked by alcohol are recognized at an early stage, a physician may be able to prevent their further development and progression. Therefore, it is essential to characterize the early events leading to alcoholic diseases and to define markers that may aid in the implementation of early preventive measures [4]. In addition, it is important to first identify the markers of long-term

alcoholism (either early or late) that can be detected before medical symptoms appear.

In this review, we aim to consider the molecules more commonly associated with these alcohol induced changes, as well as some of those less commonly considered. In addition, we focus on a pair of oxidative molecules that could be used as markers of the early stage of alcoholism, in general gaining a global idea of the potential early and late biomarkers of alcohol consumption.

Alcohol metabolism

Alcohol is generally accepted to be a compound toxic to cells or tissues that is readily metabolized to acetaldehyde by alcohol dehydrogenase (ADH). A variety of ADH isoforms have been identified, present largely in the liver, and they require a vitamin-related co-factor, nicotinamide adenine dinucleotide (NAD⁺), to accept reducing equivalents (hydrogen atoms and electrons) from the alcohol. Circulating levels of acetaldehyde are low under normal conditions and much of the acetaldehyde produced by the oxidation of alcohol is in turn oxidized in the liver to acetate. This leaves the liver and circulates to peripheral tissues where it is converted to a key Acetyl CoA.

In the liver, ethanol may also be metabolized through two additional pathways, that involving cytochrome P450 2E1 (CYP2E1) or catalase [5]. CYP2E1 plays an important role in ethanol oxidation in the brain (Figure 1).




Figure 1: Alcohol Metabolism. The enzymes alcohol dehydrogenase (ADH), cytochrome P450 (CYP450) and catalase, all contribute to the oxidative metabolism of ethanol. ADH in the cell converts alcohol to acetaldehyde, which is then metabolized mainly by aldehyde dehydrogenase (ALDH) in the mitochondria to form acetate and $\text{NADH} + \text{H}^+$.

However, this system that yields additional reactive oxygen radicals during alcohol metabolism [6,7] that can damage mitochondria, cause DNA modifications and lipid peroxidation, and elevate cytokine production [8].

In addition, chronic alcohol consumption produces fatty acid accumulation in hepatocytes, thereby decreasing the functional capacity of the liver. Chronic alcohol ingestion also alters various metabolic pathways within the liver, which ultimately leads to the production of free radicals [9]. Indeed, the lipid peroxidation mediated by free radicals is considered to be the most important mechanism responsible for cell membrane destruction and cell damage [10]. Accordingly, the uncontrolled production of free radicals is considered as an important factor in the tissue damage induced by several pathologies. Moreover, the body's defense mechanisms (enzymatic and non-enzymatic) play an important role in the formation of antioxidants that serve to minimize such damage, adapting to the stressful situations that arise.

Alcohol elimination rate

The Alcohol Elimination Rate (AER) is a measure that is conditioned by several factors, some of which are associated with the characteristics of the individual while others are more specifically modulated by the environment. As indicated above, ADH is the most important enzyme in alcohol metabolism and it directly influences the AER. With respect to the age of individuals, studies realized on animal models have indicated that the sub-maximal expression of ADH in young animals leads to a low AER [11]. Whereas the AER is thought to be higher in individuals that express the beta3 class I ADH isoform than in those who express the beta1 isoform. Moreover, alcohol metabolism is higher in the fed nutritional state than in the fasted state, presumably due to the higher ADH levels and the elevated capacity of substrate shuttle mechanisms to transport reducing equivalents into the mitochondria. Indeed, food intake may increase

liver blood flow and it is known that the sugar fructose enhances alcohol metabolism by providing substrates that help to convert NADH to NAD^+ , as well as by enhancing mitochondrial oxygen uptake. The increased AER associated with food was similar for meals with different compositions, and there appears to be no difference between carbohydrate, fat and protein on the rate of alcohol metabolism [11].

However, given that first pass metabolism occurs in the stomach, it is possible that intake of a similar dose of alcohol may produce a higher blood ethanol concentration in females than males. Conversely, men and women exhibit a similar AER when this rate is calculated in terms of gr/hr or gr/dm^3 liver volume. Moreover, liver size may at least partially explain the ethnic and gender differences in AERs [12].

Biochemical parameters and biomarkers

Despite efforts to study the molecules that could serve as early markers in individuals with an important consumption of alcohol, none have yet been identified. Nevertheless, much information has been generated regarding the biochemical and physiological parameters that are modified at these stages. For such, molecules to be considered and used as biomarkers, ideally they should display high sensitivity and specificity, discriminating between safe and hazardous drinking. Importantly, such markers should not be elevated by non-alcohol-induced organ damage and they should be non-invasive [13].

Blood, urine or breath alcohol analyses provide information as to whether the patient has alcohol in their system, yet such approaches provide no information as to the severity of alcohol abuse [14]. Moreover, the detection time is limited to hours until the alcohol consumed is metabolized to acetylaldehyde. At present, no single value can differentiate between chronic and acute abuse [15].

Below, we shall consider what are thought to be the most common potential biomarkers, as well as other non-conventional and oxidative biomarkers that could be used to determine chronic and sub-chronic exposure to ethanol (Table 1).

Biomarkers of Alcohol abuse			
Biomarkers type	Conventional	Non-conventional	Oxidative
Biochemical parameters	Blood or air ethanol levels	Phosphatidyl ethanol	Adducts formation
	Gama glutamyl transferase	5-Hydroxytryptophol	GST
	Transaminases	Sera beta-hexosaminidase	GSH, GSSG
	Carbohydrate deficient transferrin	Fatty acid ethyl esters	Catalase
	Mean corpuscular volume		Lipid peroxidation
	HDL-lipid profile		DNA damage
	Ethyl glucuronide		

Table 1: Biomarkers of alcohol Abuse. Metabolites and adducts generated during ethanol metabolism.

Conventional Biomarkers

Biomarkers have mainly been used to screen patients for possible problems of alcohol use and the conventional biomarkers considered to date will be discussed here (Figure 2).




Figure 2: Conventional Biomarkers of Alcohol: Ethylglycol (EtG), carbohydrate deficient transferrine (CDT), mean erythrocyte corpuscular volume (MCV), aspartate transferase (AST), alanine transferase (ALT) and Gamma-glutamyl transferase (GGT).

Blood ethanol

Blood ethanol is a widely accepted marker for recent ethanol consumption. Levels exceeding 1.5% with no evidence of intoxication, or of 3% at any time, are indicative of the ethanol tolerance typically found in alcohol abusers and alcohol-dependent patients. This parameter is useful for emergency clinics, although the rapid elimination of ethanol from the blood nearly always makes it impossible to assess anything other than recent ethanol ingestion in this way: within 6–8 h [16].

γ -Glutamyl transferase

Gamma-glutamyl transferase (GGT) is a glycoprotein found at the membrane of cells in several tissues. This enzyme catalyzes the first step in the degradation of GSH, and it plays important roles in glutathione homeostasis and mercapturic acid metabolism. In individuals that consume excessive alcohol, the release of GGT from the cell membrane may be enhanced [17,18] and indeed, serum-GGT has been widely used as an indicator of liver dysfunction and a marker of alcohol intake [19]. Thus, GGT represents a sensitive and inexpensive, age-dependent marker, although its specificity is diminished by obesity, diabetes, non-alcoholic liver disease, pancreatitis, hyperlipidemia, cardiac insufficiency, severe trauma, medication, nephrotic syndrome and renal rejection [17].

Mean corpuscular volume

Red blood cell size (MCV, mean corpuscular volume) is often used in screening procedures to detect alcohol abuse [18]. An increased MCV follows chronic heavy drinking and it is correlated with both the amount and frequency of alcohol ingestion. However, it may take up

to a month of drinking more than 60 g of alcohol daily to raise the MCV above the reference range [14,16]. The principal mechanism by which alcohol causes an increase in the MCV may be through a direct toxic effect of alcohol on erythrocytes and through folic acid deficiency rather than hepatic damage [20].

Formation of Carbohydrate-Deficient transferrin (CDT)

Transferrin is a liver protein that participates in iron transport and that is synthesized and secreted by the liver, with a half-life of 7–10 days. This marker is a desialylated transferrin variant that appears in the serum [21]. Alcohol abuse hampers the glycosylation of this protein such that they have significantly higher amounts of carbohydrate deficient forms of transferrin. A rise in CDT occurs after a daily consumption of 60-80 g of alcohol for at least one week and this can be normalized slowly during abstinence [22]. Serum CDT levels may be influenced by other elements unrelated to alcohol use like anorexia nervosa and pregnancy, and indeed, false positives have been associated with genetic factors, iron levels, age and liver disease [23].

Transaminases

Aspartate aminotransferase (AST) and Alanine aminotransferase (ALT) are enzymes that transform α -keto acids into amino acids. ALT is mainly present in hepatic tissue, while AST (also known as serum glutamic oxaloacetic transaminase) is found predominantly in the liver but it is also found in considerable amounts in other tissues like heart and muscle. The ratio of AST to ALT in serum may aid the diagnosis of some liver diseases and while in most patients with acute liver injury the ratio is 1 or less, in alcoholic hepatitis it is generally about 2. Predictably, the sensitivity of both these enzymes in the context of alcohol abuse is low and varies greatly [24], such that these enzymes are apparently not good biomarkers.

HDL

Alcohol intake may raise plasma high density lipoprotein (HDL) levels, either by altering the synthesis of HDL or by affecting the enzymes and proteins that influence HDL metabolism. In addition, alcohol may also alter HDL levels by increasing the plasma concentrations of apolipoproteins AI and AII, the main components of HDL [25]. Since HDL may increase as a result of drinking relatively low amounts of alcohol (≤ 5 drinks per day), it has been used to detect early phase drinking problems in individuals, as well as in the follow-up of patients with no significant liver injury. In lipid profiles, alcohol abusers also frequently show increased concentrations of serum triglycerides and free fatty acid ethyl ester following recent bouts of heavy drinking [26].

Acetaldehyde

Acetaldehyde is a product of oxidative ethanol metabolism and since it is a reactive molecule, forming Schiff bases with amines, it readily binds to proteins, leading to an irreversible reaction that give rise to an acetaldehyde-protein adduct. The concentration of acetaldehyde after alcohol intake is highly variable, with a lifetime of approximately 3 h. Two approaches have been adopted to detect acetaldehyde as a marker of alcohol intake. The first involves acetaldehyde detection in its free form, or reversibly bound to plasma proteins or blood cells. This acetaldehyde is liberated from the blood and it can be measured by gas or liquid chromatography [27]. The second approach involves the use of an immunoassay to detect

epitopes derived from acetaldehyde on proteins in the plasma. Another option is to directly measure the levels of anti-adduct IgAs as biomarkers of alcohol intake. This approach may reflect the close relationship between marker levels and the actual amounts of recent ethanol ingestion. Indeed, the specificity 88-94% and sensitivity of 65-73% confirm the potential of this approach [28].

Ethyl glucuronide (EtG)

Ethyl glucuronide (EtG) is a non-volatile, water-soluble, direct metabolite of ethanol. EtG forms in the liver when alcohol reacts with glucuronic acid, a substance that participates in the detoxification of drugs by turning them into water-soluble compounds that can be easily removed from the body [29]. EtG is only detected if alcohol is consumed and it remains present in urine well after the disappearance of ethanol itself. EtG appears to be a highly sensitive, specific and reliable marker of recent alcohol intake (up to 5 days). Moreover, EtG has also been shown to be present in other tissues and in hair, which may have additional diagnostic implications [18]. EtG has a half-life of approximately 2-3 hours [30], and it can be measured by GC/MS, LM/MS, LC/MS-MS and ELISA. However, all these techniques are currently rather laborious and technically demanding for routine laboratories.

Non-Conventional Biomarkers

Non-conventional biomarkers enable alcohol consumption to be assessed in a chronic and sub-chronic setting. Some characteristics of these non-conventional biomarkers can be seen schematically in Figure 3.




Figure 3: Non-Conventional Biomarkers of Alcohol: Hexosaminidase, phosphatidylethanol (PEth), 5-Hydroxytryptophol (5HTOL), Fatty acid ethyl ester (FAEE).

Phosphatidylethanol (PEth)

Phosphatidylethanol PEth is also a specific metabolite of ethanol that is synthesized by phospholipase D in the presence of ethanol. Therefore, the diagnostic specificity of PEth as a biomarker of alcohol is theoretically 100%. However, this marker only reacts to a single dose greater than 60 g of alcohol and thus, single doses of 32-47 g alcohol

will not be detected by this marker [31]. The half-life of PEth is about 4 days and PEth may remain in circulation for more than 2 weeks, possibly being detected for up to 3 weeks in heavy consumers. This marker is sensitive to storage and to avoid *in vitro* formation of PEth in blood, samples must be frozen at -80 °C [32].

5-Hydroxytryptophol

The serotonin metabolite 5-Hydroxytryptophol (5-HTOL) is a normal constituent of urine and after alcohol consumption, there is a marked increase in 5-HTOL probably due to inhibition of ALDH or the increase in the NAD/NADH ratio. Indeed, the concentration of the normal serotonin metabolite 5-hydroxytryptophol-3-acetic acid (5-HIAA) decreases correspondingly [24]. This marker seems to have a high diagnostic accuracy to detect recent alcohol consumption and for monitoring relapses. Accordingly, the 5-HTOL:5-HIAA ratio was found to have 100% sensitivity 4 h after consumption of a moderate dose of ethanol, although the reliability of this marker decreases fairly rapidly after 7 h [33,34]. Moreover, the HPLC-based methods currently used to determine the 5-HTO:5-HIAA ratio are difficult to translate into routine clinical practice.

Serum β -hexosaminidase

β -Hexosaminidase (HEX) is a lysosomal enzyme that is involved in the metabolism of carbohydrates and gangliosides in the liver. In both serum and urine, β -HEX has long been known to be a very sensitive biomarker for chronic alcohol use. Its activity in serum reflects excessive alcohol consumption (more than 60 g of alcohol daily for more than 6.5 days [35]). Moreover, one of the major potential strengths of β -HEX is that it can be measured using standard and inexpensive laboratory techniques (spectrophotometry and fluorimetry). However, elevated serum β -HEX also occurs in patients with hypertension, diabetes, cirrhosis, myocardial infarction, in pregnancy and after oral contraceptive use. Nevertheless, β -HEX may be a useful marker to discriminate between alcoholics and abstemious individuals or social drinkers [36].

Fatty acid ethyl esters (FAEEs)

Fatty acid ethyl ester (FAEE) synthases catalyze the reaction between ethanol and a fatty acid to produce a fatty acyl ethyl ester. FAEEs may be toxic, inhibiting DNA and protein synthesis, and high concentrations can be detected in organs commonly damaged by alcohol abuse, in particular the pancreas and liver. In serum, FAEE levels remain elevated for up to 12-24 hours after alcohol consumption [37], although the number of false positive cases represents one of the disadvantages of FAEE as a biomarker. For example, such false negatives may be produced in hair specimens as a result of the indistinctive or daily use of alcohol based lotions or sprays. The principal methods to analyze FAEEs are laborious and time-consuming, including gas chromatography-mass spectrometry, which may prevent its routine use in clinical laboratories [38].

Oxidative Biomarkers

Alcohol consumption is associated with a number of changes in cell function and in the oxidant-antioxidant system. The liver is known to be the major site of ethanol oxidation and in the first oxidation step ethanol is metabolized to acetaldehyde mainly through oxidation catalyzed by ADH. Subsequently, this acetaldehyde is rapidly oxidized to acetate by liver acetaldehyde dehydrogenase [39].

Alcohol is also metabolized in non-liver tissues, such as by CYP2E1 and catalase enzyme in the brain. This causes oxidative damage by augmenting the levels of reactive oxygen species (ROS) [40], and through reduced glutathione and superoxide dismutase activity [41].

In addition, the metabolism of alcohol increases the levels of NADH, which provides electrons for the mitochondrial electron transport chain (ETC), leading to increased one-electron reduction of oxygen to superoxide [42]. Furthermore, formation of acetaldehyde by alcohol has been shown to cause mitochondrial damage, which may also lead to increased formation of superoxide within the ETC [43]. Finally, ethanol metabolism is not only involved directly in the production of ROS but also, it has been associated with establishing a setting favorable to oxidative stress (e.g., hypoxia and endotoxaemia: Figure 4) [44].




Figure 4: Oxidative Biomarkers of Alcohol. MDA, HNE, GSH/GSSG and acetaldehyde could represent relevant biomarkers, which are produced by oxidative ethanol metabolism.

Cytochrome P450

The cytochrome P450 (CYP) enzymes, especially the CYP2E1, CYP1A2 and CYP3A4 isozymes, are involved in alcohol oxidation in the liver [45]. Specifically, CYP2E1 is induced by chronic alcohol consumption and it assumes an important role in metabolizing ethanol to acetaldehyde in the presence of elevated ethanol concentrations ($K_m = 8$ to 10 mM, as opposed to 0.2 to 2.0 mM for hepatic ADH). Since the combined activity of CYP1A2 and CYP3A4 is comparable to that of CYP2E1, these CYP can contribute significantly to microsomal ethanol oxidation and therefore, they may also be involved in any related pathophysiological effects [46]. In the metabolism of ethanol to acetaldehyde via CYP2E1, and through a typical monooxygenase mechanism, ethanol can be oxidized by liver microsomes through hydroxyl radicals. These may include those originating from iron catalyzed Haber-Weiss and Fenton reactions degradation of H_2O_2 [47]. In addition, acetaldehyde's toxicity is due, in part, to its capacity to bind to microsomal proteins, including CYP2E1 [48]. The formation of protein adducts results in antibody

production, enzyme inactivation and decreased DNA repair, as well as a score of other toxic manifestations that include a decrease in GSH [49].

Glutathione S-Transferase (GST)

A superfamily of enzymes exists that catalyze the transfer of a glutathione group to electrophilic xenobiotics or substrates, and that also exhibit peroxidase activity. Both, these transferase and peroxidase activities have been proved to be altered by alcohol consumption [50]. Initial studies found that Glutathione S-Transferase (GST) from the livers of ethanol fed rats is modified by a product of ethanol metabolism, the malondialdehyde-acetaldehyde adduct [51]. These results suggest that GST enzymes play an important role in the detoxification of reactive aldehydes, including those that participate in alcohol-induced liver injury [52]. The Alpha, Mu, Theta and Pi families of GST are involved in antioxidant effects and as such, the GSTs represent a group of polymorphic enzymes that are important in the protection against oxidative stress [53].

Reduced glutathione (GSH)

The reduced glutathione (GSH) tripeptide is involved in many cellular functions and it plays a central role in antioxidant defense, especially in the liver. Chronic ethanol consumption selectively depletes GSH in the mitochondria due to the defective activity of this carrier, probably mediated by decreased fluidity of the mitochondrial inner membrane [54]. The endogenous glutathione-glutathione peroxidase system and catalase are important antioxidants, and they form part of the cytoprotective machinery in the hepatocyte mitochondria that is activated on exposure to ethanol [55]. Glutathione reductase and catalase activities increase on exposure to ethanol, and chronic alcoholism induces alterations in the GSH/GSSG ratio that might enhance the oxidative modification of liver mitochondrial proteins [56].

Catalase

Catalase is found in the cytosol and mitochondria, although it is mainly active in peroxisomes. This enzyme is very efficient in promoting the conversion of hydrogen peroxide to water and molecular oxygen. Catalase is capable of oxidizing ethanol in vitro in the presence of a hydrogen peroxide (H_2O_2)-generating system, such as the NADPH oxidase enzyme complex or the enzyme xanthine. This is considered a minor pathway of alcohol oxidation, except in the fasting conditions [57]. Chronic alcohol consumption by rats results in increased H_2O_2 production in pericentral regions of the liver lobule and increased catalase activity [58]. Moreover, several studies have suggested that catalase may be involved in the metabolism of alcohol to acetaldehyde in the brain [46], which could in turn mediate some of the behavioral effects of alcohol.

Lipid peroxidation

Lipid peroxidation is a complex process whereby polyunsaturated fatty acids in the phospholipids of cellular membranes undergo a reaction with oxygen to yield lipid hydroperoxides (LOOH) [59]. This process results in the formation of compounds known as malondialdehyde (MDA) and 4-hydroxy-2-nonenal (HNE), both of which can form adducts with proteins [60]. MDA concentrations in alcoholics with or without cirrhosis are higher than in matched healthy controls [61] and therefore, MDA is considered as an excellent

biomarker to reveal and monitor oxidative stress triggered by alcohol consumption [62]. As mentioned earlier, ethanol metabolism by CYP2E1 and NADH oxidation by the ETC generate ROS, which drive lipid peroxidation. Lipid peroxidation products can be generated from the attack of ROS and iron on lipids or by CYP2E1 activity [63]. An increase in lipid peroxidation products has been detected in individuals with a 4-5 year history of alcohol drinking [10].

DNA damage

Chronic ethanol consumption leads to an increase in ROS and acetaldehyde, which have both been shown to induce DNA damage, including oxidative modifications, adducts and cross-linking [64]. 7,8-dihydro-8-oxo-2'-deoxyguanosine as the most representative product of modified DNA, while the primary acetaldehyde-derived DNA adduct is N-2-ethylidene-deoxyguanosine, which may be converted to N 2-ethyldeoxyguanosine (N 2-ethyl-dG)in vivo, a potential biomarker of acetaldehyde induced DNA damage in human alcoholics [65].

Endogenous DNA damage is constantly being produced and repaired in normal conditions, reflecting a low-steady state level of damage that is compatible with normal cellular function. Indeed, many studies have quantified the five basic types of DNA repair [66]. However, in conditions when DNA repair is deficient, endogenous damage is not repaired and accumulates over time, resulting in a variety of symptoms or diseases, such as liver cirrhosis in chronic stages or alcohol related neuronal loss in the brain. In the case of alcohol drinkers, it was assumed that oxidative process could only be observed with long-term and high doses of alcohol intake, however, excess alcohol consumption produces deleterious effects on many tissues in the body, even if this excess occurs in a short time window, such as a week. In this sense, we recently published a study where we observed biomarkers of oxidative damage in a group of young people who have been drinking alcohol for a relatively short period of time (4-5 years) [10].

Conclusions

In this review, we have focused on most of molecules that have been used as markers to identify individuals that consume an important amount of alcohol, each of which is are involved in some manner in the development of diverse signs and pathological processes that have an impact on the health of these individuals. However, the majority of work carried out both in vitro and in a clinical setting has concentrated on high consumption over long periods. Fewer studies have assessed biomarkers of early detection that could be present in individuals considered to be "social drinkers" or those that consume a moderate quantity of alcohol every so often. We believe that important changes exist in the cells of drinkers even within a short period after the onset of alcohol consumption, ergo, before any symptoms of organ or tissue damage can be perceived. Indeed, we reported that biomarkers that reflect oxidative damage and that were validated long ago for the determination of chronic stages of alcoholism could also be suitable to be used at earlier stages of consumption. Nevertheless, more studies should be conducted to identify and validate the use of these or other biomarkers in the early stages of alcoholism, which will have important benefits in clinical practice and with the goal of preventing a major deterioration of health. Finally, it is important to note that most studies analyzing molecules or biomarkers have been carried out on adults. However, we think that is important to realize such analysis in young people, due to the period of consumption and

the future establishment of behaviors. More research is necessary to explore novel biomarkers or combinations of biomarkers known to reflect drinking patterns with high sensitivity and specificity.

References

1. Anderson P (2009) Is it time to ban alcohol advertising? *Clin Med* 9: 121-124.
2. Okulicz-Kozaryn K, Foxcroft DR (2012) Effectiveness of the Strengthening Families Programme 10-14 in Poland for the prevention of alcohol and drug misuse: protocol for a randomized controlled trial. *BMC Public Health* 12: 319.
3. Wakefield JC, Schmitz MF (2014) How Many People have Alcohol Use Disorders? Using the Harmful Dysfunction Analysis to Reconcile Prevalence Estimates in Two Community Surveys. *Front Psychiatry* 5: 10.
4. Hartwig S, Auwärter V, Pragst F (2003) Fatty Acid ethyl esters in scalp, pubic, axillary, beard and body hair as markers for alcohol misuse. *Alcohol Alcohol* 38: 163-167.
5. Crabb DW, Liangpunsakul S (2007) Acetaldehyde generating enzyme systems: roles of alcohol dehydrogenase, CYP2E1 and catalase, and speculations on the role of other enzymes and processes. *Novartis Found Symp* 285: 4-16.
6. Zakhari S (2013) Alcohol metabolism and epigenetics changes. *Alcohol Res* 35: 6-16.
7. Dupont I, Bodénez P, Berthou F, Simon B, Bardou LG, et al. (2000) Cytochrome P-450 2E1 activity and oxidative stress in alcoholic patients. *Alcohol Alcohol* 35: 98-103.
8. Caro AA, Cederbaum AI (2004) Oxidative stress, toxicology, and pharmacology of CYP2E1. *Annu Rev Pharmacol Toxicol* 44: 27-42.
9. Lieber CS (1988) Biochemical and molecular basis of alcohol-induced injury to liver and other tissues. *N Engl J Med* 319: 1639-1650.
10. Rendón-Ramírez A, Cortés-Couto M, Martínez-Rizo AB, Muñiz-Hernández S, Velázquez-Fernández JB (2013) Oxidative damage in young alcohol drinkers: A preliminary study. *Alcohol* 47: 501-504.
11. Cederbaum AI (2012) Alcohol metabolism. *Clin Liver Dis* 16: 667-685.
12. Cole-Harding S, Wilson JR (1987) Ethanol metabolism in men and women. *J Stud Alcohol* 48: 380-387.
13. Das SK1, Nayak P, Vasudevan DM (2003) Biochemical markers for alcohol consumption. *Indian J Clin Biochem* 18: 111-118.
14. Neumann T, Spies C (2003) Use of biomarkers for alcohol use disorders in clinical practice. *Addiction* 98 Suppl 2: 81-91.
15. Helander A, Eriksson G, Stibler H, Jeppsson JO (2001) Interference of transferrin isoform types with carbohydrate-deficient transferrin quantification in the identification of alcohol abuse. *Clin Chem* 47: 1225-1233.
16. Aston ER, Liguori A (2013) Self-estimation of blood alcohol concentration: a review. *Addict Behav* 38: 1944-1951.
17. Bataller-Sifré R, Guiral-Olivan V, Bataller-Alberola L (2011) New clinical and toxicological scenario of gammaglutamyltranspeptidase. *Rev Esp Enferm Dig* 103: 586-590.
18. Tavakoli HR1, Hull M, Michael Okasinski L (2011) Review of current clinical biomarkers for the detection of alcohol dependence. *Innov Clin Neurosci* 8: 26-33.
19. Zhang H, Forman HJ (2009) Redox regulation of gamma-glutamyl transpeptidase. *Am J Respir Cell Mol Biol* 41: 509-515.
20. Sharpe PC (2001) Biochemical detection and monitoring of alcohol abuse and abstinence. *Ann Clin Biochem* 38: 652-664.
21. Kwoh-Gain I, Fletcher LM, Price J, Powell LW, Halliday JW (1990) Desialylated transferrin and mitochondrial aspartate aminotransferase compared as laboratory markers of excessive alcohol consumption. *Clin Chem* 36: 841-845.
22. Koch H, Meerkerk GJ, Zaat JO, Ham MF, Scholten RJ, et al. (2004) Accuracy of carbohydrate-deficient transferrin in the detection of

- excessive alcohol consumption: a systematic review. *Alcohol Alcohol* 39: 75-85.
23. Fleming MF, Anton RF, Spies CD (2004) A review of genetic, biological, pharmacological, and clinical factors that affect carbohydrate-deficient transferrin levels. *Alcohol Clin Exp Res* 28: 1347-1355.
 24. Torrente MP, Freeman WM, Vrana KE (2012) Protein biomarkers of alcohol abuse. *Expert Rev Proteomics* 9: 425-436.
 25. Collins MA, Neafsey EJ, Mukamal KJ, Gray MO, Parks DA, et al. (2009) Alcohol in moderation, cardioprotection, and neuroprotection: epidemiological considerations and mechanistic studies. *Alcohol Clin Exp Res* 33: 206-219.
 26. Laposata M (1997) Fatty acid ethyl esters: short-term and long-term serum markers of ethanol intake. *Clin Chem* 43: 1527-1534.
 27. Musshoff F (2002) Chromatographic methods for the determination of markers of chronic and acute alcohol consumption. *J Chromatogr B Analyt Technol Biomed Life Sci* 781: 457-480.
 28. Hietala J, Koivisto H, Latvala J, Anttila P, Niemelä O (2006) IgAs against acetaldehyde-modified red cell protein as a marker of ethanol consumption in male alcoholic subjects, moderate drinkers, and abstainers. *Alcohol Clin Exp Res* 30: 1693-1698.
 29. Skipper GE, Weinmann W, Thierauf A, Schaefer P, Wiesbeck G, et al. (2004) Ethyl glucuronide: a biomarker to identify alcohol use by health professionals recovering from substance use disorders. *Alcohol Alcohol* 39: 445-449.
 30. Schmitt G, Droenner P, Skopp G, Aderjan R (1997) Ethyl glucuronide concentration in serum of human volunteers, teetotalers, and suspected drinking drivers. *J Forensic Sci* 42: 1099-1102.
 31. Hansson PI, Caron M, Johnson G, Gustavsson L, Alling C (1997) Blood phosphatidylethanol as a marker of alcohol abuse: levels in alcoholic males during withdrawal. *Alcohol Clin Exp Res* 21: 108-110.
 32. Niemelä O (2007) Biomarkers in alcoholism. *Clin Chim Acta* 377: 39-49.
 33. Borucki K, Schreiner R, Dierkes J, Jachau K, Krause D, et al. (2005) Detection of recent ethanol intake with new markers: comparison of fatty acid ethyl esters in serum and of ethyl glucuronide and the ratio of 5-hydroxytryptophol to 5-hydroxyindole acetic acid in urine. *Alcohol Clin Exp Res* 29: 781-787.
 34. Helander A, Some M (2000) Dietary serotonin and alcohol combined may provoke adverse physiological symptoms due to 5-hydroxytryptophol. *Life Sci* 67: 799-806.
 35. Hannuksela ML, Liisanantti MK, Nissinen AE, Savolainen MJ (2007) Biochemical markers of alcoholism. *Clin Chem Lab Med* 45: 953-961.
 36. Kärkkäinen P, Poikolainen K, Salaspuro M (1990) Serum beta-hexosaminidase as a marker of heavy drinking. *Alcohol Clin Exp Res* 14: 187-190.
 37. Bisaga A, Laposata M, Xie S, Evans SM (2005) Comparison of serum fatty acid ethyl esters and urinary 5-hydroxytryptophol as biochemical markers of recent ethanol consumption. *Alcohol Alcohol* 40: 214-218.
 38. Auwärter V, Sporkert F, Hartwig S, Pragst F, Vater H, et al. (2001) Fatty acid ethyl esters in hair as markers of alcohol consumption. Segmental hair analysis of alcoholics, social drinkers, and teetotalers. *Clin Chem* 47: 2114-2123.
 39. Zhu H, Jia Z, Misra H, Li YR (2012) Oxidative stress and redox signaling mechanisms of alcoholic liver disease: updated experimental and clinical evidence. *J Dig Dis* 13: 133-142.
 40. Lu Y, Gong P, Cederbaum AI (2008) Pyrazole induced oxidative liver injury independent of CYP2E1/2A5 induction due to Nrf2 deficiency. *Toxicology* 252: 9-16.
 41. Kessova IG, Cederbaum AI (2007) Mitochondrial alterations in livers of Sod1^{-/-} mice fed alcohol. *Free Radic Biol Med* 42: 1470-1480.
 42. Das SK, Vasudevan DM (2007) Alcohol-induced oxidative stress. *Life Sci* 81: 177-187.
 43. Boveris A, Fraga CG, Varsavsky AI, Koch OR (1983) Increased chemiluminescence and superoxide production in the liver of chronically ethanol-treated rats. *Arch Biochem Biophys* 227: 534-541.
 44. Sergent O, Griffon B, Cillard P, Cillard J (2001) [Alcohol and oxidative stress]. *Pathol Biol (Paris)* 49: 689-695.
 45. Salmela KS, Kessova IG, Tsyrllov IB, Lieber CS (1998) Respective roles of human cytochrome P-4502E1, 1A2, and 3A4 in the hepatic microsomal ethanol oxidizing system. *Alcohol Clin Exp Res* 22: 2125-2132.
 46. Lieber CS (1999) Microsomal ethanol-oxidizing system (MEOS): the first 30 years (1968-1998)--a review. *Alcohol Clin Exp Res* 23: 991-1007.
 47. Cederbaum AI (1989) Oxygen radical generation by microsomes: role of iron and implications for alcohol metabolism and toxicity. *Free Radic Biol Med* 7: 559-567.
 48. Behrens UJ, Hoerner M, Lasker JM, Lieber CS (1988) Formation of acetaldehyde adducts with ethanol-inducible P450IIE1 in vivo. *Biochem Biophys Res Commun* 154: 584-590.
 49. Lieber CS (2005) Metabolism of alcohol. *Clin Liver Dis* 9: 1-35.
 50. Sherrat PJ, Hayes JD (2001) Enzyme Systems that Metabolise Drugs and Other Xenobiotics: Glutathione S-Transferases. (1stedn), Costas Ioannides.
 51. Xu D, Thiele GM, Kearley ML, Haugen MD, Klassen LW, et al. (1997) Epitope characterization of malondialdehyde-acetaldehyde adducts using an enzyme-linked immunosorbent assay. *Chem Res Toxicol* 10: 978-986.
 52. Khan AJ, Choudhuri G, Husain Q, Parmar D (2009) Polymorphism in glutathione-S-transferases: a risk factor in alcoholic liver cirrhosis. *Drug Alcohol Depend* 101: 183-190.
 53. Brind AM, Hurlstone A, Edrington D, Gilmore I, Fisher N, et al. (2004) The role of polymorphisms of glutathione S-transferases GSTM1, M3, P1, T1 and A1 in susceptibility to alcoholic liver disease. *Alcohol Alcohol* 39: 478-483.
 54. Fernández-Checa JC, García-Ruiz C, Colell A, Morales A, Mari M, et al. (1998) Oxidative stress: role of mitochondria and protection by glutathione. *Biofactors* 8: 7-11.
 55. Kurose I, Higuchi H, Kato S, Miura S, Ishii H (1996) Ethanol-induced oxidative stress in the liver. *Alcohol Clin Exp Res* 20: 77A-85A.
 56. Bailey SM, Patel VB, Young TA, Asayama K, Cunningham CC (2001) Chronic ethanol consumption alters the glutathione/glutathione peroxidase-1 system and protein oxidation status in rat liver. *Alcoholism, Clinical and Experimental Research* 25: 726-733.
 57. Handler JA, Thurman RG (1990) Redox interactions between catalase and alcohol dehydrogenase pathways of ethanol metabolism in the perfused rat liver. *J Biol Chem* 265: 1510-1515.
 58. Misra UK, Bradford BU, Handler JA, Thurman RG (1992) Chronic ethanol treatment induces H₂O₂ production selectively in pericentral regions of the liver lobule. *Alcohol Clin Exp Res* 16: 839-842.
 59. Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M (2006) Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chem Biol Interact* 160: 1-40.
 60. Worrall S, Thiele GM (2001) Protein modification in ethanol toxicity. *Adverse Drug React Toxicol Rev* 20: 133-159.
 61. Huang MC, Chen CC, Peng FC, Tang SH, Chen CH (2009) The correlation between early alcohol withdrawal severity and oxidative stress in patients with alcohol dependence. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 66-69.
 62. Chen CH, Pan CH, Chen CC, Huang MC (2011) Increased oxidative DNA damage in patients with alcohol dependence and its correlation with alcohol withdrawal severity. *Alcohol Clin Exp Res* 35: 338-344.
 63. Setshedi M, Wands JR, Monte SM (2010) Acetaldehyde adducts in alcoholic liver disease. *Oxid Med Cell Longev* 3: 178-185.
 64. Brooks PJ, Theruvathu JA (2005) DNA adducts from acetaldehyde: implications for alcohol-related carcinogenesis. *Alcohol* 35: 187-193.
 65. Fang JL, Vaca CE (1995) Development of a 32P-postlabelling method for the analysis of adducts arising through the reaction of acetaldehyde with 2'-deoxyguanosine-3'-monophosphate and DNA. *Carcinogenesis* 16: 2177-2185.

-
66. Kruman II, Henderson GI, Bergeson SE (2012) DNA damage and neurotoxicity of chronic alcohol abuse. *Exp Biol Med (Maywood)* 237: 740-747.

This article was originally published in a special issue, entitled: "[Drug and Alcohol abuse](#)", Edited by Asok K Dasmahapatra, University of Mississippi, USA