

An update on Dendritic Cell-Based Cancer Immunotherapy

Shimodaira S^{1*}, Hirabayashi K¹, Koya T¹, Higuchi Y¹, Yanagisawa R¹, Okamoto M² and Koido S³¹Center for Advanced Cell Therapy, Shinshu University Hospital, Matsumoto, Nagano 390-8621, Japan²Department of Advanced Immunotherapeutics, Kitasato University School of Pharmacy, Minato-ku 108-8641, Tokyo, Japan³Department of Gastroenterology and Hepatology, the Jikei University School of Medicine, Kashiwa, Chiba 277-8567, Japan

Abstract

Although treating advanced cancers that affect organs with distant metastasis remains challenging, the pace of recent advances has accelerated; these advances have particularly focused on the inhibitors of key immune potentiators. Research on therapeutic vaccination involving active dendritic cell (DC)-based immunotherapy is also being performed for the induction of an efficient immune response against cancer-associated antigens by the acquired immune system. Cancer vaccines prepared with autologous monocyte-derived mature DCs have been generated using granulocyte-macrophage colony-stimulating factor and interleukin-4, which are principally attributed to the presence of tumor-associated antigens. Wilms' tumor 1 (WT1) is an attractive target antigen that is widely detected in many cancers. DC-based immunotherapy targeting WT1 may elicit a strong therapeutic response to cancers. DC vaccines primed with HLA class I/II-restricted WT1 peptides (WT1-DC) are a feasible option for patients with advanced cancers. Immune response monitoring using tetramer analysis and/or enzyme-linked immunosorbent spot assay has been applied to determine the efficacy of WT1-DC. The inhibition of immune suppressors and acceleration of anti-cancer immunity with WT1-DC may comprise a promising future therapeutic strategy for treating advanced cancers.

Keywords: Dendritic cells; Cancer vaccination; Wilms' tumor 1; Tetramer analysis; Enzyme-linked immunosorbent spot assay

Abbreviations: APC: Antigen-Presenting Cell; DCs: Dendritic Cells; RECIST: Response Evaluation Criteria in Solid Tumors; HLA: Human Leukocyte Antigen; MDSCs: Myeloid-Derived Suppressor Cells; CTLs: Cytotoxic T Cells; GM-CSF: Granulocyte-Macrophage Colony-Stimulating Factor; TGF: Transforming Growth Factor; IL: Interleukin; VEGF: Vascular Endothelial Growth Factor; PGE2: Prostaglandin E2; PD-L1: Programmed Death-Ligand 1; FDA: Food and Drug Administration; WT1: Wilms' Tumor 1; MUC1: Mucin 1 Cell Surface Associated; HER2: Human Epidermal Growth Factor Receptor 2; CEA: Carcinoembryonic Antigen; PSA: Prostate-Specific Antigen; WT1-DC: DC Vaccines Primed with HLA class I/II-Restricted WT1 Peptides; ELISPOT: Enzyme-Linked Immunosorbent Spot; WT1-CTLs: WT1 Antigen-Specific Cytotoxic T Lymphocytes; IFN: Interferon; PBMCs: Peripheral Blood Mononuclear Cells; HSCT: Hematopoietic Stem Cell Transplantation; G-CSF: Granulocyte Colony-Stimulating Factor

Introduction

Immune checkpoint inhibitors have resulted in significant advances in cancer therapeutics however; the treatment of advanced cancers affecting organs and involving distant metastasis remains extremely difficult [1-6]. Clinical benefit of an anti-programmed death 1 (PD1) immune checkpoint inhibitor was shown in patients with progressive metastatic colorectal cancer with mismatch repair-deficiency [7].

Antigen-presenting cell (APC)-based immunotherapy using active dendritic cells (DCs) is under investigation for developing therapeutic vaccination against cancer [8]. Autologous DC-based immunotherapy appears to trigger few adverse reactions with limited clinical effectiveness if conventional evaluation procedures such as response evaluation criteria in solid tumors (RECIST) are employed [9,10]. Long-term acquired immunity following DC vaccination produces a delayed separation of survival curves with an advantage pertaining to prolonging the overall survival [11,12].

Human leukocyte antigen (HLA) molecules containing cancer antigen peptides on DCs bind with T cell receptors on the CD8+


killer and CD4⁺ helper T cells, producing an immune response against cancers. In contrast, immune suppressor cells such as regulatory T cells, and myeloid-derived suppressor cells (MDSCs) suppress the auto reactive and anti-cancer immunity [13-15]. Moreover, tolerogenic DCs with immunosuppressive cytokines induce antigen-specific anergy and regulatory properties in memory CD4⁺ T cells [16-18]. Conversely, inflammatory DCs are driven by infection or inflammation release inflammatory cytokines, including tumor necrosis factor and the inducible nitric oxide synthase [19]. Moreover, inflammatory DCs drive Th1- and Th17-mediated immunity and facilitate the induction of regulatory T cells [20,21] (Figure 1). Immune suppressive factors are also produced by cancer cells; these factors include transforming growth factor (TGF)- β , interleukin (IL)-10, vascular endothelial growth factor (VEGF), prostaglandin E2 (PGE2), and programmed death-ligand 1 (PD-L1) [22] (Figure 1). Chemotherapeutic drugs, targeted anti-cancer agents and various other biological and physicochemical therapies such as radiation therapy have been identified as employing immunogenic cell death. DCs are primed with cancer cells succumbing to these immunogenic cell death inducers [23]. Radiation therapy aside from the induction of cancer cell death is useful to shift an immunosuppressive tumor microenvironment to a more beneficial immune stimulation. Radiation therapy can also enhance the expression of HLA class I on the surface of cancer cells, boosting the recognition and killing of irradiated cancer cells through T cells and NK cells [24]. DCs in combination with chemoradiotherapy may accelerate the development of acquired

***Corresponding author:** Shigetaka Shimodaira, Center for Advanced Cell Therapy, Shinshu University Hospital, Matsumoto, Nagano 390-8621, Japan, Tel: +81-263-37-3580; Fax: +81-263-37-3027; E-mail: shimodai@shinshu-u.ac.jp

Received January 06, 2016; **Accepted** March 09, 2016; **Published** March 15, 2016

Citation: Shimodaira S, Hirabayashi K, Koya T, Higuchi Y, Yanagisawa R, et al. (2016) An update on Dendritic Cell-Based Cancer Immunotherapy. Immunome Res 12: 106. doi: [10.4172/17457580.1000106](https://doi.org/10.4172/17457580.1000106)

Copyright: © 2016 Shimodaira S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



cancer immunity to induce antigen-specific cytotoxic T lymphocytes (CTLs) [25,26]. Chemoradiotherapy induce immunogenic cell death, which could trigger T-cell immunity mediated by high-mobility group box 1 protein in patients with oesophageal squamous cell cancer [27]. Radiotherapy and chemotherapeutic drugs with off-target effects may allow the in situ modulation of regulatory T cells and other suppressors within the tumor microenvironment (Figure 1). DCs and regulatory T cells are more resistant to radiation than other lymphocytes in mouse model [28,29]. Human monocyte-derived DCs as well as macrophages are more radio resistant than monocytes [30]. Tumor necrosis factor α and interferon (IFN)- γ secreted by activated T cells react with cancer cells, which causes the formation of complex of PD-L1 on cancer cells with PD1 on activated T cells. PD1-PDL1 complex suppresses the interactions between HLA molecule and T cell receptor, resulting in regression of anti-cancer immunity [31].

DC Vaccine and Vaccination

A manufacturing technique is being developed to promote the strong induction of T cells against tumor antigens. Oil adjuvants for peptide vaccines act by locally accelerating the activation of lymphocytes [32]. However, DCs have potential bioactivity that can be used as a suitable adjuvant [33-34]. DCs expressing tumor-specific antigens used in active cancer immunotherapies [35,36] have been conventionally generated using peripheral monocytes with granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin-4 (IL-4). Antigenic peptides, protein, tumor lysate, and RNA have been used to deliver cancer-associated antigens into DCs [37-42]. Sipuleucel-T (Provenge[®], Dendreon Corporation, Seattle, WA), which has been approved by the US Food and Drug Administration (FDA), is an autologous, DC-based immunotherapy for patients with metastatic hormone-refractory

prostate cancer. Sipuleucel-T is manufactured using patient's blood cells exposed to a recombinant fusion protein comprising a prostatic acid phosphatase with GM-CSF, which enhances its activity against prostate cancer. The autologous DC product is administered as per a 3-dose schedule with approximately 2-week intervals between each dose. This regimen yields a survival benefit of 4.1 months when given to patients with hormone-resistant prostate cancer [43].

Among potential cancer antigens such as Wilms' tumor 1 (WT1); mucin 1, cell surface associated (MUC1); human epidermal growth factor receptor 2 (HER2); carcinoembryonic antigen (CEA); survivin; and prostate-specific antigen (PSA), WT1 was identified to be the most potent cancer-associated antigen fulfilling immunological and clinical effectiveness criteria with respect to therapeutic functions, immunogenicity, specificity, and oncogenicity [44]. WT1 has recently been shown to regulate the expression of VEGF, a major mediator of angiogenesis, suggesting it to be another target within the cancer microenvironment [45,46]. Furthermore, the WT1 peptide was restricted to HLA-A*24:02 and modified WT1235-243 peptide (CYTWNQMNL) and the second amino acid (methionine: M) was replaced with tyrosine (Y), which can more effectively induce cytotoxic T cells than wild-type peptides [47-50]. The WT1-332-347-class II peptide is compatible with HLA-DRB1*04:05, HLA-DRB1*08:03, DRB1*15:01, DRB1*15:02, DPB1*05:01, or DPB1*09:01. Phase I clinical trials of this peptide have been conducted for various types of cancers and hematologic malignancies [51-53].

DC vaccines primed with HLA class I/II-restricted WT1 peptides (WT1-DC) have been shown to be safe and feasible with few adverse reactions reported for patients with advanced cancers, including lung, breast, stomach, biliary tract, pancreatic, colorectal, ovarian,

and high-grade gliomas [54-62]. Clinical studies have indicated the efficacy of DC vaccination as an add-on to chemotherapy [54-59] and chemoradiotherapy [62,63] and even suggested a survival benefit in some patients. Combinations of adjuvant chemotherapy and/or radiotherapy as well as the periods required for adaptation to these therapies have been investigated. The development of combination therapies that potentially include immune checkpoint inhibitors may help improve the outcomes of personalized therapy for cancer patients.

Immune Monitoring

Immunological monitoring of DC vaccination using tetramer analysis and/or enzyme-linked immunosorbent spot (ELISPOT) assays in clinical studies and trials requires both reproducibility and validation. The presence of WT1 antigen-specific cytotoxic T cells (WT1-CTLs) is defined according to the following criteria: presence of greater than 0.02% WT1-positive CD8⁺ T cells among the 50,000–10,000 lymphocytes analyzed with no evidence of false positive cells and WT1-positive cell population being clustered but not diffused with slight modification, as previously described [64]. ELISPOT assays were performed to measure WT1-specific IFN- γ production by peripheral blood mononuclear cells (PBMCs). The presence of WT-CTLs was defined according to the following criteria: presence of at least 15 WT1-specific spots per 1 \times 10⁶ PBMCs and at least 50% more WT1-specific spots than negative peptide (HIV peptide) spots [64].

We determined WT1-CTLs by both WT1-peptide/HLA-A*24:02 tetramer analysis and ELISPOT assay after one course of DC vaccination during maintenance chemotherapy in a patient with gastric cancer. After one course of DC vaccination, the immune monitoring assay demonstrated that WT1-CTLs comprised 1.10% of the CD8⁺ T cell population (Figure 2A), with over 100 WT1-specific spots observed in ELISPOT assays (Figure 2B, upper panel). ELISPOT assays performed using WT1-332 (HLA-class II peptide) also demonstrated a specific number of IFN- γ -spots (Figure 2B, lower panel). Specific WT-CTLs were persisted for more than 1 year after DC vaccination.


DC Vaccination Technology

Allogeneic vaccines induce T cell infiltration and aggregate formation in pancreatic cancer, resulting in the induction of

immunosuppressive regulatory mechanisms [65]. However, allogeneic DC vaccination targeting WT1 also represents a potential strategy for treating patients with relapsed leukemia following hematopoietic stem cell transplantation (HSCT). Subsequent evidence of immune monitoring would provide proof of the validity of this concept. This strategy may be safe, tolerable, and even feasible for pediatric donors and patients with relapsed leukemia following HSCT, as previously described [66]. In this report, a 15-year-old girl with acute lymphoblastic leukemia received allogeneic DC vaccination pulsed with WT1 peptide after her third HSCT. The vaccines were generated from the third HSCT donor, the patient's younger 12-year-old sister, who was a HLA-A*24:02 match. The patient received 14 vaccine doses with no occurrence of graft-versus-host disease or any systemic adverse effects, except for a grade 2 local skin reaction at the injection site. WT1-specific immune responses were detected post-vaccination by both WT1-tetramer analysis and ELISPOT assay. The patient experienced 44 months of remission after the third HSCT with DC vaccination, whereas she had been in remission for less than 14 months between her second and third HSCT. This suggests that WT1-specific DC vaccination contributed to extended remission after the patient's third HSCT. These findings form the basis for developing individualized therapy for future prospective HSCT clinical trials.

One potential way to overcome the phenomenon of tumor cells escaping immune detection is the generation of IFN-DCs from monocytes using GM-CSF and IFN- α previously described in a clinical study on metastasized medullary thyroid carcinoma [67] under Good Gene, Cell and Tissue Manufacturing Practice conditions. Mature IFN-DCs would induce CTLs together with their adaptive antitumor effects as well adoptive immunotherapy with natural killer cell activity independent of HLA antigen expression [67,68]. Another approach is the administration of granulocyte colony-stimulating factor (G-CSF), resulting in the upregulation of adhesion molecules, to evaluate the hypothesis of increased acquired immunity using G-CSF-primed DC vaccines.

Immune checkpoint inhibitors are rapidly being developed as cancer therapeutic agents [69]. In combination with DC vaccination and immune checkpoint inhibition, further studies are required to evaluate whether the number of effector memory T cell present



prior to vaccination and the exhausted marker of PD1-positive CTLs after DC vaccination influence the efficacy of DC vaccination and/or immune check point inhibitors. Such future clinical trials could reveal the effectiveness of DC vaccine in combination with immune checkpoint inhibitors in treating cancers, sarcomas, and hematopoietic malignancies in the near future. Biomarkers that predict the potential efficacy of DC vaccination targeting WT1 are highly relevant to current standards of personalized cancer therapy.

Conflict of Interest

All authors declare no conflict of interest that could be perceived as prejudicing the impartiality of the reported research.

Acknowledgments

This study was supported by the grants-in-aid for scientific research from the Japan Society for the Promotion of Science (JSPS). The authors would like to thank Enago (www.enago.jp) for the English language review.

References

- Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, et al. (2010) Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 363: 711-723.
- Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, et al. (2012) Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 366: 2443-2454.
- Hamid O, Robert C, Daud A, Hodi FS, Hwu WJ, et al. (2013) Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *N Engl J Med* 369: 134-144.
- Brahmer JR, Tykodi SS, Chow LQ, Hwu WJ, Topalian SL, et al. (2012) Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med* 366: 2455-2465.
- Wolchok JD, Kluger H, Callahan MK, Postow MA, Rizvi NA, et al. (2013) Nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med* 369: 122-133.
- Weber JS, Kudchadkar RR, Yu B, Gallenstein D, Horak CE, et al. (2013) Safety, efficacy, and biomarkers of nivolumab with vaccine in ipilimumab-refractory or -naive melanoma. *J Clin Oncol* 31: 4311-4318.
- Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, et al. (2015) PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med* 372: 2509-2520.
- Steinman RM (2012) Decisions about dendritic cells: past, present, and future. *Annu Rev Immunol* 30: 1-22.
- Rosenberg SA, Yang JC, Restifo NP (2004) Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 10: 909-915.
- Ridgway D (2003) The first 1000 dendritic cell vaccinees. *Cancer Invest* 21: 873-886.
- Hoos A (2012) Evolution of end points for cancer immunotherapy trials. *Ann Oncol* 23 Suppl 8: viii47-52.
- Schlom J, Arlen PM, Gulley JL (2007) Cancer vaccines: moving beyond current paradigms. *Clin Cancer Res* 13: 3776-3782.
- Nishikawa H, Sakaguchi S (2010) Regulatory T cells in tumor immunity. *Int J Cancer* 127: 759-767.
- Whiteside TL (2014) Induced regulatory T cells in inhibitory microenvironments created by cancer. *Expert Opin Biol Ther* 14: 1411-1425.
- Nagaraj S, Gabrilovich DI (2008) Tumor escape mechanism governed by myeloid-derived suppressor cells. *Cancer Res* 68: 2561-2563.
- Torres-Aguilar H, Aguilar-Ruiz SR, González-Pérez G, Munguía R, Bajaña S, et al. (2010) Tolerogenic dendritic cells generated with different immunosuppressive cytokines induce antigen-specific anergy and regulatory properties in memory CD4+ T cells. *J Immunol* 184:1765-1775.
- Ezzelarab M, Thomson AW (2011) Tolerogenic dendritic cells and their role in transplantation. *Semin Immunol* 23: 252-263.
- Svajcer U, Rozman P (2014) Tolerogenic dendritic cells: molecular and cellular mechanisms in transplantation. *J Leukoc Biol* 95: 53-69.
- Schlitzer A, Ginhoux F (2014) Organization of the mouse and human DC network. *Curr Opin Immunol* 26: 90-99.
- Segura E, Touzot M, Bohineust A, Cappuccio A, Chiochia G, et al. (2013) Human inflammatory dendritic cells induce Th17 cell differentiation. *Immunity* 38:336-348.
- Osorio F, Fuentes C, López MN, Salazar-Onfray F, González FE (2015) Role of Dendritic Cells in the Induction of Lymphocyte Tolerance. *Front Immunol* 6: 535.
- Guo C, Manjili MH, Subjeck JR, Sarkar D, Fisher PB, et al. (2013) Therapeutic cancer vaccines: past, present, and future. *Adv Cancer Res* 119: 421-475.
- Garg AD, Galluzzi L, Apetoh L, Baert T, Birge RB, et al. (2015) Molecular and Translational Classifications of DAMPs in Immunogenic Cell Death. *Front Immunol* 6: 588.
- Derer A, Deloch L, Rubner Y, Fietkau R, Frey B, et al. (2015) Radio-Immunotherapy-Induced Immunogenic Cancer Cells as Basis for Induction of Systemic Anti-Tumor Immune Responses - Pre-Clinical Evidence and Ongoing Clinical Applications. *Front Immunol* 6: 505.
- Duffy AG, Gretten TF (2014) Immunological off-target effects of standard treatments in gastrointestinal cancers. *Ann Oncol* 25: 24-32.
- Shimodaira S, Kobayashi T, Hirabayashi K, Horiuchi K, Koya T, et al. (2015) Induction of antigen-specific cytotoxic T lymphocytes by chemoradiotherapy in patients receiving Wilms' tumor 1-targeted dendritic cell vaccinations for pancreatic cancer. *OMICS J Radiol* 4: 4.
- Suzuki Y, Mimura K, Yoshimoto Y, Watanabe M, Ohkubo Y, et al. (2012) Immunogenic tumor cell death induced by chemoradiotherapy in patients with esophageal squamous cell carcinoma. *Cancer Res* 72: 3967-3976.
- Bogdándi EN, Balogh A, Felgyinszki N, Szatmári T, Persa E, et al. (2010) Effects of low-dose radiation on the immune system of mice after total-body irradiation. *Radiat Res* 174: 480-489.
- Kachikwu EL, Iwamoto KS, Liao YP, DeMarco JJ, Agazaryan N, et al. (2011) Radiation enhances regulatory T cell representation. *Int J Radiat Oncol Biol Phys* 81: 1128-1135.
- Bauer M, Goldstein M, Christmann M, Becker H, Heylmann D, et al. (2011) Human monocytes are severely impaired in base and DNA double-strand break repair that renders them vulnerable to oxidative stress. *Proc Natl Acad Sci U S A* 108: 21105-21110.
- Ribas A (2012) Tumor immunotherapy directed at PD-1. *N Engl J Med* 366: 2517-2519.
- Hailemichael Y, Dai Z, Jaffarzar N, Ye Y, Medina MA, et al. (2013) Persistent antigen at vaccination sites induces tumor-specific CD8 α^+ T cell sequestration, dysfunction and deletion. *Nat Med* 19: 465-472.
- Banchereau J, Steinman RM (1998) Dendritic cells and the control of immunity. *Nature* 392: 245-252.
- Steinman RM, Banchereau J (2007) Taking dendritic cells into medicine. *Nature* 449: 419-426.
- Figdor CG, de Vries IJ, Lesterhuis WJ, Melief CJ (2004) Dendritic cell immunotherapy: mapping the way. *Nat Med* 10: 475-480.
- O'Neill DW, Adams S, Bhardwaj N (2004) Manipulating dendritic cell biology for the active immunotherapy of cancer. *Blood* 104: 2235-2246.
- Nestle FO, Farkas A, Conrad C (2005) Dendritic-cell-based therapeutic vaccination against cancer. *Curr Opin Immunol* 17: 163-169.
- Copier J, Ward S, Dalgleish A (2007) Cell based cancer vaccines: regulatory and commercial development. *Vaccine* 25 Suppl 2: B35-46.
- Ramakrishnan R, Antonia S, Gabrilovich DI (2008) Combined modality immunotherapy and chemotherapy: a new perspective. *Cancer Immunol Immunother* 57: 1523-1529.
- Baxeveanis CN, Perez SA, Papamichail M (2009) Combinatorial treatments including vaccines, chemotherapy and monoclonal antibodies for cancer therapy. *Cancer Immunol Immunother* 58: 317-324.
- Koido S, Homma S, Takahara A, Namiki Y, Tsukinaga S, et al. (2011) Current immunotherapeutic approaches in pancreatic cancer. *Clin Dev Immunol* 2011: 267539.
- Koido S, Ohkusa T, Homma S, Namiki Y, Takakura K, et al. (2013) Immunotherapy for colorectal cancer. *World J Gastroenterol* 19: 8531-8542.

43. Kantoff PW, Higano CS, Shore ND, Berger ER, Small EJ, et al. (2010) Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* 363: 411-422.
44. Cheever MA, Allison JP, Ferris AS, Finn OJ, Hastings BM, et al. (2009) The prioritization of cancer antigens: a national cancer institute pilot project for the acceleration of translational research. *Clin Cancer Res* 15: 5323-5337.
45. McCarty G, Awad O, Loeb DM (2011) WT1 protein directly regulates expression of vascular endothelial growth factor and is a mediator of tumor response to hypoxia. *J Biol Chem* 286: 43634-43643.
46. Katuri V, Gerber S, Qiu X, McCarty G, Goldstein SD, et al. (2014) WT1 regulates angiogenesis in Ewing Sarcoma. *Oncotarget* 5: 2436-2449.
47. Oka Y, Elisseeva OA, Tsuboi A, Ogawa H, Tamaki H, et al. (2000) Human cytotoxic T-lymphocyte responses specific for peptides of the wild-type Wilms' tumor gene (WT1) product. *Immunogenetics* 51: 99-107.
48. Ohnishi H, Yasukawa M, Fujita S (2000) HLA class I-restricted lysis of leukemia cells by a CD8(+) cytotoxic T-lymphocyte clone specific for WT1 peptide. *Blood* 95: 286-293.
49. Tsuboi A, Oka Y, Uda K, Murakami M, Masuda T, et al. (2002) Enhanced induction of human WT1-specific cytotoxic T lymphocytes with a 9-mer WT1 peptide modified at HLA-A*2402-binding residues. *Cancer Immunol Immunother* 51: 614-620.
50. Fujiki F, Oka Y, Tsuboi A, Kawakami M, Kawakatsu M, et al. (2007) Identification and characterization of a WT1 (Wilms Tumor Gene) protein-derived HLA-DRB1*0405-restricted 16-mer helper peptide that promotes the induction and activation of WT1-specific cytotoxic T lymphocytes. *J Immunother* 30: 282-293.
51. Oka Y, Tsuboi A, Taguchi T, Osaki T, Kyo T, et al. (2004) Induction of WT1 (Wilms' tumor gene)-specific cytotoxic T lymphocytes by WT1 peptide vaccine and the resultant cancer regression. *Proc Natl Acad Sci U S A* 101: 13885-13890.
52. Oka Y, Tsuboi A, Murakami M, Hirai M, Tominaga N, et al. (2003) Wilms tumor gene peptide-based immunotherapy for patients with overt leukemia from myelodysplastic syndrome (MDS) or MDS with myelofibrosis. *Int J Hematol* 78: 56-61.
53. Nishida S, Koide S, Takeda Y, Homma S, Komita H, et al. (2014) Wilms tumor gene (WT1) peptide-based cancer vaccine combined with gemcitabine for patients with advanced pancreatic cancer. *J Immunother* 37: 105-114.
54. Kimura Y, Tsukada J, Tomoda T, Takahashi H, Imai K, et al. (2012) Clinical and immunologic evaluation of dendritic cell-based immunotherapy in combination with gemcitabine and/or S-1 in patients with advanced pancreatic carcinoma. *Pancreas* 41: 195-205.
55. Takahashi H, Okamoto M, Shimodaira S, Tsujitani S, Nagaya M, et al. (2013) Impact of dendritic cell vaccines pulsed with Wilms' tumour-1 peptide antigen on the survival of patients with advanced non-small cell lung cancers. *Eur J Cancer* 49: 852-859.
56. Kobayashi M, Sakabe T, Abe H, Tanii M, Takahashi H, et al. (2013) Dendritic cell-based immunotherapy targeting synthesized peptides for advanced biliary tract cancer. *J Gastrointest Surg* 17: 1609-1617.
57. Kobayashi M, Shimodaira S, Nagai K, Ogasawara M, Takahashi H, et al. (2014) Prognostic factors related to add-on dendritic cell vaccines on patients with inoperable pancreatic cancer receiving chemotherapy: a multicenter analysis. *Cancer Immunol Immunother* 63: 797-806.
58. Koide S, Homma S, Okamoto M, Takakura K, Mori M, et al. (2014) Treatment with chemotherapy and dendritic cells pulsed with multiple Wilms' tumor 1 (WT1)-specific MHC class I/II-restricted epitopes for pancreatic cancer. *Clin Cancer Res* 20: 4228-4239.
59. Kobayashi M, Chiba A, Izawa H, Yanagida E, Okamoto M, et al. (2014) The feasibility and clinical effects of dendritic cell-based immunotherapy targeting synthesized peptides for recurrent ovarian cancer. *J Ovarian Res* 7: 48.
60. Kobayashi M, Sakabe T, Chiba A, Nakajima A, Okamoto M, et al. (2014) Therapeutic effect of intratumoral injections of dendritic cells for locally recurrent gastric cancer: a case report. *World J Surg Oncol* 12: 390.
61. Sakai K, Shimodaira S, Maejima S, Udagawa N, et al. (2015) Dendritic cell-based immunotherapy targeting Wilms' tumor 1 in patients with recurrent malignant glioma. *J Neurosurg* 123: 989-997.
62. Shimodaira S, Sano K, Hirabayashi K, Koya T, Higuchi Y, et al. (2015) Dendritic cell-based adjuvant vaccination targeting Wilms' tumor1 in patients with advanced colorectal cancer. *Vaccines* 3: 1004-1018.
63. Shibamoto Y, Okamoto M, Kobayashi M, Ayakawa S, Iwata H, et al. (2013) Immune-maximizing (IMAX) therapy for cancer: Combination of dendritic cell vaccine and intensity-modulated radiation. *Mol Clin Oncol* 1: 649-654.
64. Higuchi Y, Koya T, Yuzawa M, Yamaoka N, Mizuno Y, et al. (2015) Enzyme-linked immunosorbent spot assay for the detection of Wilms' tumor1-specific T cells induced by dendritic cell vaccination. *Biomedicines* 3: 304-315.
65. Lutz ER, Wu AA, Bigelow E, Sharma R, Mo G, et al. (2014) Immunotherapy converts nonimmunogenic pancreatic tumors into immunogenic foci of immune regulation. *Cancer Immunol Res* 2: 616-631.
66. Saito S, Yanagisawa R, Yoshikawa K, Higuchi Y, Koya T, et al. (2015) Safety and tolerability of allogeneic dendritic cell vaccination with induction of Wilms tumor 1-specific T cells in a pediatric donor and pediatric patient with relapsed leukemia: a case report and review of the literature. *Cytotherapy* 17: 330-335.
67. Papewalis C, Jacobs B, Wuttke M, Ullrich E, Baehring T, et al. (2008) IFN-alpha skews monocytes into CD56+ -expressing dendritic cells with potent functional activities in vitro and in vivo. *J Immunol* 180: 1462-1470.
68. Chan CW, Crafton E, Fan HN, Flook J, Yoshimura K, et al. (2006) Interferon-producing killer dendritic cells provide a link between innate and adaptive immunity. *Nat Med* 12: 207-213.
69. Shih K, Arkenau HT, Infante JR (2014) Clinical impact of checkpoint inhibitors as novel cancer therapies. *Drugs* 74: 1993-2013.

Citation: Shimodaira S, Hirabayashi K, Koya T, Higuchi Y, Yanagisawa R, et al. (2016) An update on Dendritic Cell-Based Cancer Immunotherapy. *Immunome Res* 12: 106. doi: [10.4172/17457580.1000106](https://doi.org/10.4172/17457580.1000106)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>