



---

**Research Article**

---

**ANTHELMINTIC ACTIVITY OF METHANOLIC EXTRACT OF RHIZOMES OF *PICRORRHIZA KURROA ROYAL EX. BENTH.***

A. Rajani<sup>1</sup>, K.Hemamalini<sup>2\*</sup>, D.Satyavati<sup>3</sup>, S.K.Arifa Begum<sup>1</sup>, NDVR.Saradhi<sup>1</sup>

<sup>1</sup>Associate Professors, <sup>3</sup>Principal/Professor, Sree Dattha Institute of Pharmacy, Sheriguda, Ibrahimpatnam, Greater Hyderabad. <sup>2</sup>Associate Professor & HOD, Teegala Ram Reddy College of Pharmacy, Meerpet, Hyderabad.

**\*Corresponding Author: Email**

**(Received: August 11, 2013; Accepted: September 19, 2013)**

**ABSTRACT**

The objective of this study is to evaluate and compare the Anthelmintic activity of methanolic extract of *Picrorrhiza kurroa Royle ex. Benth* (Scrophulariaceae). *Picrorrhiza kurroa* a small perennial herb growing in the hilly parts of the North-western Himalayan region in India and Nepal. Earth worms were used for Anthelmintic activity. Piperazine citrate was used as standard drug. Time required for paralysis and death of the earth worms were noted for each sample.

**Keywords:** Pheretima posthuma, Piperazine citrate, *Picrorrhiza kurroa*, Anthelmintic activity.

**INTRODUCTION**

*Picrorrhiza kurroa Royle ex. Benth.* Belonging to the family Scrophulariaceae is a small perennial herb that is widely distributed in the North-west India on the slopes of Himalayas between 3000 and 5000 meters<sup>1,2</sup>. *Picrorrhiza kurroa* is valued as hepatoprotective, Anti-periodic, Cholagogue, Stomachic, Anti-amoebic, Anti-oxidant, Anthelmintic, Anti-inflammatory, Cardiotonic, Laxative, Carminative, Expectorant etc<sup>3, 4</sup>. On account of its use as Anthelmintic as well as bitter, this study was undertaken to evaluate the Anthelmintic potential. Anthelmintics are the drugs that expel parasitic worms from the body either by paralyzing or killing them<sup>5</sup>. Helminth infections are now being recognized as the cause of many acute as well as chronic ill healths among the various human beings as well as cattle's. More than half of the population of the world suffers from

infection of one or the other and majority of cattle population suffer from worm infections<sup>6</sup>. The objective of the present study is to evaluate Anthelmintic activity of methanolic extract of rhizomes of *Picrorrhiza kurroa* plant.

**MATERIALS AND METHODS**

***Authentication of the plant material:***

The plant specimen was collected from S.V University; Tirupati and has been identified as *Picrorrhiza kurroa Royle ex. Benth.* Belonging to the family Scrophulariaceae, Voucher No: SDIP, Ref No: 002 dated 26/10/2012 and authenticated by Dr.K.Madhavachetty, Botanist, Tirupati. The rhizomes of the plant were dried in vacuum oven at 40° C.

***Preparation of the plant extract:***

Rhizomes of *Picrorrhiza kurroa* plant are coarsely powdered and are successively extracted by continuous hot percolation

method using Soxhlet apparatus employing chloroform and methanol. Methanolic extraction yielded sufficiently good quantity of the product. From the dried methanolic extract, accurately 20mg/ml, 40mg/ml, 60mg/ml, 80mg/ml, 100mg/ml, 120mg/ml solutions were prepared using distilled water.

**Standard used for the activity:**

Piperazine citrate is used as the standard to compare the test results. The standard drug was prepared in the concentrations of 10mg/ml using distilled water as solvent.

**Animals:**

Adult earth worms (*Pheretima posthuma*) of about 6-8cms long were chosen for Anthelmintic activity.

pin, which stimulated or induced movements if the worm was alive.

**RESULTS AND DISCUSSION**

Results of preliminary phytochemical screening showed the presence of carbohydrates, glycosides, saponins, steroids like phytochemical constituents. Cucurbitacins, Phenolic, Iridoid glycosides are some of the principle constituents responsible for various pharmacological activities. Iridoid glycosides like kutkin, Picroliv, Picrisides I, II, III & IV, Kutkosides are the chemical moieties that may be responsible for Anthelmintic activity<sup>8</sup>. The Table no: 01 reveals that the methanolic extract obtained from the rhizomes of *Picrorrhiza kurroa* are active against earth worm. Methanolic extract of *Picrorrhiza*

**Table: 1 Anthelmintic activity of methanolic extract of leaves *Picrorrhiza kurroa***

Treatment	Concentration used (mg/ml)	Time taken for paralysis (min) X ± S.D	Time taken for death (min) X ± S.D
Control	-	-	-
Standard (Piperazine citrate)	10mg/ml	17 ± 1.571*	39 ± 1.932*
Methanolic extract of <i>Picrorrhiza kurroa</i>	20mg/ml	76 ± 3.303*	99.33 ± 0.402*
	41mg/ml	65.33 ± 2.883**	90.67 ± 3.921**
	60mg/ml	56 ± 2.017**	85.00 ± 5.310**
	80mg/ml	48.33 ± 1.498**	69.83 ± 2.496**
	100mg/ml	48.33 ± 1.498**	51.67 ± 2.108**
	120mg/ml	27.33 ± 2.060**	40.33 ± 0.9888***

All values are mean ± SEM and analyzed by one way ANOVA followed by Dunnett's test n= 5, \*p<0.05, \*\*p<0.01, \*\*\*p<0.001

**Method:**

The Anthelmintic assay was carried as per the method of Ajaiyeoba et.al. with minor modifications<sup>7</sup>. The Anthelmintic activity was evaluated on Indian adult worms (*Pheretima posthuma*) available at a nearby vermin culture plant. Six different concentrations i.e. 20mg/ml, 40mg/ml, 60mg/ml, 80mg/ml, 100mg/ml, 120mg/ml of test drug and 10mg/ml of standard were prepared in different petri dishes. Six worms were placed in each petri dish. Observations were made for the time taken (in minutes) for paralysis/immobility and death of individual worms in all the 12 petri dishes. Death of the worm was ascertained when the worm showed no further movements upon transferring the worm into a beaker containing hot water at 50°C or pricking it by a ball

*kurroa* rhizomes showed good Anthelmintic activity in a dose dependent manner. The Anatomy and Physiology of *Pheretima posthuma* is similar to that Helminthes<sup>9</sup>. Therefore earth worms were used in this study. It has been demonstrated that all Anthelmintics are toxic to earth worms, and a substance toxic to earth worm is worthy for investigation as an Anthelmintic<sup>10</sup>. Further studies are needed to establish the mode of activity. The Anthelmintic activity can be ascertained by testing the drug on other species of helminthes which is our future part of research work.

**ACKNOWLEDGEMENTS**

Authors are thankful to manager of the vermin culture plant for providing us the earth worms. Authors also express

sincere gratitude to the management and Directors of Sree Dattha Institutions, Sheriguda, Ibrahimpatnam, Greater Hyderabad for providing us the facilities to carry out the research activity and their timely support.

#### REFERENCES

1. Hooker J.D, Flora of British India, 246-290, 1885.
2. Chopra R.N and Ghosh S, Some common indigenous remedies, Indian journal of Medical Research 22; 263-264, 1934.
3. Pacpute A.P., Deshmukh T.A., "Antioxidant and Hepatoprotective Activity of an Ethanol Extract of Piper Cubeba Fruits", Int. J. Res. Dev. Pharm. L. Sci., 2012, 2 (2), pp. 321-329.
4. Kapahi B.K, Srinivastava T.N and Sarin Y.K, Description of Picrorrhiza kurroa, A source of the Ayurvedic drug Kutaki, International Journal of Pharmacognosy 31: 217-222, 1993.
5. Singh G.B, Sarang B, Singh S, Khajuria A, Sharma M.L, Gupta B.d and Banerjee S.K, Anti inflammatory activity of the Iridoids kutkin, picroside-I and Kutkoside from picrorrhiza kurroa, Phytotherapy Research 7: 402-407, 2006.
6. Encyclopaedia Britannica, 2009.
7. Anonymous, The Ayurvedic Pharmacopoeia of India, First edition, Part-I, Vol-II, Control publication, Delhi, 2000.
8. Ajaiyeoba E.O, Onoeha P.A, Olarenwaju O.T, In-vitro Anthelmintic properties of Buchholzia coriacea and Gynandropsis gynandra extract, Pharm Biol., 39, 217-220, 2001.
9. Parimala S., Kuamr V.H., Kumar S.M.S., Suthakaran R., Selvan A.T., "Analgesic and CNS depressant Activity of Methanolic Extracts of Cissus Vitegenia And Cissus Pallida", Int. J. Res. Dev. Pharm. L. Sci., 2012, 2 (2), pp. 330-332.
10. Ghisalberti E.L, Biological and Pharmacological activity of naturally occurring Iridoids and Secoiridoids, Phytomedicine 5, 147-163, 1998.
11. Edwards CA, Testing the effects of chemicals on earth worms. The advantages and limitations of field tests. Ecotoxicology of earth worms 1, 75-84, 1992.
12. Sollmann T, Anthelmintics: Their efficiency as tested on tested on earth worms. J. Pharm Exp Ther 12: 129, 1918.

#### How to cite this article:

Rajani A., Hemamalini K., Satyavati D., Begum S. K. A., Saradhi NDVR., "Anthelmintic activity of methanolic extract of rhizomes of *picrorrhiza kurroa royal ex. benth.*", Int. J. Res. Dev. Pharm. L. Sci., 2013, 2(6), pp. 705-707.