

## Antibiogram of *Klebsiella* Sp. Isolated from Ascitic Fluid of Canine Infected with Pyometra

Durgalakshmi R\*, Sheeba A, Rajagopalan V, Masilamoni Ronald BS, Ananda Chitra M, Ponnusamy P and Manickam R

Department of Veterinary Microbiology, Veterinary College and Research Institute, Orathanadu, Tamil Nadu, India

### Abstract

Pyometra, a hormonally mediated diestrual disorder is described by thickening of the uterine lining and accumulation of uterine secretions (cystic endometrial hyperplasia). These secretions leaks out to the abdomen and forms ascites, which provides environment for bacterial growth. *E. coli* is the common organism that migrates from the vagina into the uterus and causes ascites. As a first report, a bacterium was isolated from such an ascitic fluid of pyoderma infected Rottweiler canine breed and further the organism was identified as *klebsiella sp.* based on biochemical tests. Antibiogram of *klebsiella sp.* revealed that the organism was sensitive to ciprofloxacin, erythromycin, tetracycline, ampicillin, amikacin and gentamycin and resistant to methicillin.

**Keywords:** Ascitic fluid; Pyometra; Rottweiler

### Introduction

Pyometra is an important deadly disease, which affects the fertility of female dogs. "Pyo" meaning pus and "metra" meaning uterus. Thus pyometra factually means pus filled uterus. This condition is similar to appendicitis in humans. The major cause of Pyometra is the hormonal changes in reproductive tract. Following estrus cycle ("heat"), progesterone levels gets elevated for eight to ten weeks and this thickens the lining of the uterus in preparation for pregnancy. If pregnancy does not happen for several subsequent oestrus cycles, the thickened lining forms cyst (Cystic endometrial hyperplasia) [1] and secretes fluid in which bacteria can infect and grow. The path of the bacteria is the opening of the cervix during estrus cycle. Once the bacteria enter, the cervix gets tightly closed, and forms a barrier between the uterus and the outside. Later the uterus ruptures and leaks out the pus into abdomen, causing distended and enlarged abdomen, a condition known as Ascites [2]. Since the bacteria cannot expel, bacterial toxins may enter the bloodstream (Toxemia). As a result, dogs with "closed" pyometra and ascites can quickly become ill developing breathing difficulty, cough, weakness and pale gums and finally the dog expires. *E.coli* is the common bacterium isolated from ascitic fluid, and this is the first report of isolating the *Klebsiella sp.* Very few reports explains the penetration of antibiotics onto ascitic fluid [3]. In this study, Antibiogram test for *klebsiella sp.* was conducted and found that it was sensitive to ciprofloxacin, erythromycin, tetracyclin, ampicillin, amikacin and gentamycin and resistant to methicillin.

### Materials and Methods

A Rottweiler female canine aged seven years was brought to veterinary dispensary, Pudukkottai district, with a history of vomiting, loss of appetite, depression and increased drinking and urination. On clinical examination, the abdomen looks bloated and on tapping the abdomen produced a dull, flat noise indicating the case of ascites due to pyometra. Exudate fluid from ascites was collected by advancing a needle into the most pendulous area of the abdomen while keeping a slight negative pressure in the syringe [2]. The ascitic fluid was subjected to microbiological analysis. Fluid sample was streaked on 5% sheep blood agar and MacConkey agar plates and kept for aerobic incubation at 37°C for 24 hrs. Colonies from the plates were subsequently isolated and identified on the basis of morphology (Grams staining and capsule

staining) and standard biochemical analysis (Oxidase test, Indole, Methyl red, Voges proskauer, citrate, Triple sugar iron, Motility test and catalase tests) as described by Quinn et al. [4]. The organism was subjected to Antibiotic susceptibility test on Muller Hinton Agar (MHA) as described by Bauer et al. [5]. The efficiency of antibiotics was determined by measuring the diameter of zone of inhibition around the discs.

### Results and Discussion

Pure growth of mucoid colonies in blood agar and pink colonies in MacConkey agar were observed after 24 hrs of incubation. Gram's staining revealed that the organism was gram negative and rod shaped. Capsule staining confirmed the presence of capsule which is also the characteristic of *Klebsiella sp.* Biochemical analysis described that the organism was non motile, oxidase negative, catalase positive and further IMViC test showed that the organism was indole negative, Methyl red negative, voges proskauer positive, citrate positive and also in TSI, organism exhibited acidic slant and acidic butt without H<sub>2</sub>S production. Thus based on these interpretations, the organism was identified as the opportunistic pathogen *Klebsiella sp.* Antibiogram results demonstrated that the organism was sensitive to ciprofloxacin, Erythromycin, tetracycline, ampicillin, amikacin and gentamycin and resistant to methicillin (Figure 1).

Although antimicrobial treatment is not the complete treatment for pyometra, this can reduce the severity of the infection that spreading from abdomen. So far *E.coli* has been isolated from canine pyometra exudate fluid and this study confirmed the presence of another opportunistic pathogen *klebsiella sp.* in ascitic fluid.

\*Corresponding author: Durgalakshmi R, Department of Veterinary Microbiology, Veterinary College and Research Institute, Orathanadu, Tamil Nadu, India, Tel: 044 2555 1583; E-mail: [durgabio14@gmail.com](mailto:durgabio14@gmail.com)

Received October 14, 2015; Accepted December 07, 2015; Published December 09, 2015

Citation: Durgalakshmi R, Sheeba A, Rajagopalan V, Masilamoni Ronald RS, Ananda Chitra M, et al. (2015) Antibiogram of *Klebsiella* Sp. Isolated from Ascitic Fluid of Canine Infected with Pyometra. J Med Microb Diagn 5: 212. doi:10.4172/2161-0703.1000212

Copyright: © 2015 Durgalakshmi R, et al.. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Figure 1:** Antibiogram of klebsiella species isolated ascitic fluid of rottweiler canine.

## References

1. Baithalu RK, Maharana BR, Mishra C, Sarangi L, Samal L (2010) Canine Pyometra. Veterinary World 3: 340-342.
2. Peden W, Michael, Zenoble R D (1982) Canine Ascites. Iowa State University Veterinarian 44: 3.
3. Wilson DE, Chalmers TC, Madoff MA (1967) The passage of cephalothin into and out of ascetic fluid. Am J Med Sci 253: 449-452.
4. Quinn PJ, Carter ME, Markey B, Carter GR (2004) Clinical Veterinary Microbiology, Mosby Elsevier Limited. Edinburgh, London, New York, Oxford, Philadelphia, St Louis, Sydney, Toronto.
5. Bauer AW, Kirby WMM, Sherris JC, Turck M (1966) Antibiotic susceptibility testing by a standardized single disk method. AM J Clin Pathol 45: 493-496.

**Citation:** Durgalakshmi R, Sheeba A, Rajagopalan V, Masilamoni Ronald RS, Ananda Chitra M, et al. (2015) Antibiogram of *Klebsiella* Sp. Isolated from Ascitic Fluid of Canine Infected with Pyometra. J Med Microb Diagn 5: 212. doi:10.4172/2161-0703.1000212

## Submit your next manuscript and get advantages of OMICS Group submissions

### Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

### Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsgroup.org/journals/submission/>