

Antibiotic Resistance Bacteria from Rivers Water in Algeria

Katia Djenadi*

Laboratory of Microbial Ecology, Faculty of Sciences of Nature and Life, University of Bejaia, 06000 Bejaia, Algeria

*Corresponding author: Katia Djenadi, Laboratory of Microbial Ecology, Faculty of Sciences of Nature and Life, University of Bejaia, 06000 Bejaia, Algeria, E-mail: djenadikatia@yahoo.fr

Received date: August 04, 2017; Accepted date: September 07, 2017; Published date: September 11, 2017

Copyright: © 2017 Djenadi K. This is an open-access article distributed under the terms of the creative commons attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The aim of the current investigation was to look on the presence of the antibiotic resistance bacteria in natural environment in Algeria. From fresh water, multidrug resistant bacteria were harvested, and then investigated for their resistance profile and extend spectrum Beta lactamase and metallo Beta lactamase production. From isolate bacteria only *Aeromonas hydrophila* catch our attention, because of their metallo Beta lactamase production and alimentary and clinical impact. This finding confirmed our hypothesis that natural environment could be colonized by multidrug resistant bacteria especially to carbapenem antibiotic who may be induced by mutation or horizontal gene transfer. The spread of these kinds of resistance organisms may become a serious issue for the public health and food security.

Keywords: *Aeromonas hydrophila*; Antibiotic resistance; River; Algeria

Introduction

Freshwater constitute favourable living ecosystem for numerous organisms such as *Aeromonas* spp [1]. Water constitutes not only a way of dissemination of antibiotic-resistant organisms among human and animal population, as drinking water is produced from surface water, but also the route by which resistance genes are introduced in natural bacterial ecosystems. Many of these bacteria harbour antibiotic resistance genes, mainly inserted in plasmid, transposant or intergen able to be transferred among water and soil bacteria community via horizontal gene transfer [2].

Aeromonas are Gram negative bacilli organisms. They occupied several environmental areas including: aquatic habitats, fish, foods, domesticated pets, invertebrate species, birds, ticks and insects and natural soils [3]. *Aeromonas* are involved in several human infections, mainly in gastroenteritis, cellulites, meningitis, bacteraemia, soft tissue infections, peritonitis and brochopulmonary infection [4]. *Aeromonas* genome species (*A. hydrophila*, *A. caviae* and *A. veronni* bv *sorbia*) are the main actor for the majority of human infections [5].

Against these infections; therapeutic molecule failed because of the resistance mechanism evolved by pathogenic agent. Antibiotic resistance is one of the major clinical issues. As predicted by Fleming the microbial organisms would evolve resistance mechanisms against the action of the antibiotics [6]. In fact, numerous resistance mechanisms have been reported since Fleming predicted this phenomenon. One of the most common mechanisms is against β lactam antibiotics, by production of β lactamases. Within *Aeromonas* genus, Beta lactam resistance principally mediated by three classes of β lactamases including, class C cephalosporinase, class D penicillinase and class B metallo Beta Lactamase [5].

Due to the overuse of broad spectrum antibiotics in clinical settings, agriculture, and fish hatcheries, antibiotic resistance has dramatically raised among disease causing *Aeromonas* [7].

There by, *Aeromonas* species received an increased attention from clinical investigators, because of their implication in serious human infections and antibiotic resistant rise within clinical [8].

Outside of the clinical environment, *Aeromonas* species express shy multidrug resistance. From aquatic samples, *Aeromonas* species have been reported to be susceptible to a variety of antimicrobial agents [9]

The current study was designed to evaluate the antibiotic resistance of bacilli Gram negative isolated from freshwater in Algeria. The investigations lunched with hypothesis that natural environment could be colonized by multidrug resistant bacteria especially to carbapenem antibiotic.

Materials and Methods

Sampling

Current study investigated on multidrug resistant Gram negatives bacteria. Around 15 samples were harvested from water river, sludge and wastewater in Setif region "Ait ourthilane" (Algeria) during winter period (2015).

Bacterial isolation and identification

From each sample serial dilution were prepared then enriched. One hundred micro litres from enrichment bacterial suspension was streaked on Mac Conkey (Himedia, USA) and Chromocult medium (Oxoid, UK) both supplemented with imipenem (0.5 μ l/mg) and vancomycin (16 μ g/ml). Then, isolates were purified on Chromocult medium (Oxoid, UK) with imipenem (0.5 μ l/mg) and conserved in agar slants [10].

The identification of isolates was carried out using API 20E (BIOMERIEUX, France). The obtained results were analysed using apiWeb BIOMERIEUX resources (<https://apiweb.biomerieux.com/servlet/Authenticate>).

Antimicrobial agent and resistance study

For this study we used the following antimicrobial agent: cefotaxim (CTX 30µg), ceftazidime (CAZ 30µg), aztreonam (ATM 30µg), ertapenem (ETP 10µg), meropenem (MEM10µg), imipenem (IMP 10µg) and amikacin (AK 30µg) (Oxoid, UK). The susceptibility and resistance of the isolates was performed following the disc diffusion method. Overnight culture resuspended in 0.9% of saline solution. Bacterial suspension with turbidity equivalent to 0.5 Mac Farland were inoculated by streaking the cotton swab on Muller Hinton agar (Oxoid, UK) plates then incubated at 37°C. After 24 h, the inhibition zones were measured and compared to guide line of the EUCAST (2014) [11].

From the isolates, multidrug resistant strains selected for the detection of extend spectrum beta lactamase (ESBLs) and metallo β lactamase (MBL) production. ESBL production was performed with double disc synergy test essay (DDST) [12]. Then revelation of the metallo β lactamase (MBL) carried out following the Ethylene diamine tetra acetic acid (EDTA), combined disc [13] and synergy test [14]. Then Carba NP [15] and Hodge test [12] were carried out.

Antibiotic	IMP (10 µg)	MEM (10 µg)	ETP (10 µg)	CAZ (30 µg)	CTX (30 µg)	AK (30 µg)	ATM (30 µg)	ESBL	MBL
Diameter zone (mm)	22	25	15	6	15	26	40	None	Yes

Table 1: Resistance profile of *Aeromonas hydrophila* RWBOCw3 isolate.

Discussion

Outside of the clinical environment, antibiotic molecules may be detected in natural environment. The antibiotic molecules are naturally produced by environmental micorbiota, though in concentration much lower compared to those used in therapy [16]. From nature, most potent molecule occurring β lactam were carbapenems and olixanic acids [17]. The natural environment may be the most important source of antibiotic resistance genes Czekalski et al. [18], reported that freshwater aquatic environment represent a potential reservoir of antibiotic genes such as sulfamides, tetracycline and fluroquinolone resistant genes [19].

From Seine River, France, Girlich and co-workers confirmed their hypothesis that *Aeromonas* spp could be an important environmental reservoir for ESBLs; a variety of ESBLs genes were detected from *Aeromonas* spp resistant to ceftazidime [9]. Other investigation from Eastern Basin of Lake Erie in Pennsylvania, *Aeromonas* spp (*A. hydrophila* and *A. veronii*) express resistant to tetracycline and ciprofloxacin [7].

Collection of water samples from different aquatic environments within an urban water cycle in the region of Northern Portugal followed by resistance profile determination reveal resistance to ceftazidime or meropenem in isolates from the drinking water treatment plant and waste water isolates were intrinsically resistant to nalidixic acid [20].

Within *Aeromonas* species, resistance to cephamycins and extend spectrum cephalosporins mediated by class C cephalosporinase (AmpC family). While, class B Metallo Beta Lactamase induced by CphA type within *Aeromonas hydrophila* and *Aeromonas veronii* clinical and environmental isolates express resistance to ampicillin, oxacillin, cephaloridine and imipenem [21]. Two other metallo beta lactamase (MBLs) (VIM and IMP) have identified in strains of

Results

In our investigation, from water and sludge river in Setif region "Ait Ourthilane" 30 isolates were obtained and only one strains from river water *Aeromonas hydrophila* RWBOCw3 with API 20 E profile: 1247521, showed interesting resistant profile against carbapenem antibiotic, especially imipenem, ertapenem cefotaxim and ceftazidime, while *Aeromonas hydrophila* RWBOCw3 were sensitive to meropenem, amikacin and Aztreonam Table 1. First investigation on the detection of ESBLs (DD test) did not show any formation of ghost inhibition zones between the central discs suggesting that carbapenem resistance could not be mediated by production of ESBL. While production of MBL was confirmed by positive results from Hodge and EDTA test; an distorted inhibition zone was observed on Hodge test plates and synergy image was formed and difference was observed between the inhibition zone of imipenem+EDTA and the inhibition zone of imipenem alone. With Carba NP test, red methyl changed the colour to yellow within 20min after incubation indicating possible carbapenemase carriage.

Aeromonas hydrophila and *Aeromonas caviae*, harboured on an integron and a plasmid, respectively [22,23].

Besides, misuse of antibiotic in clinical and veterinary set, anthropogenic human activity, spontaneous mutation and horizontal gene transfer are the principal actor in the dissemination and persistence of multidrug resistance in environment [24]. Resistant genes carried on mobile element such plasmid, integron and transposon led to spread among bacterial environment, even in the presence of selective pressure or antibiotic inducer at high or low concentration [9].

Depending on the bacterial population hosted in the environment, whether pathogens or no pathogens, the environment has an important influence on the development of multidrug resistant. Driven on chromosome or extra-chromosome or associated with mobile genetic element such as plasmids, integrons, genes cassettes or transposons, resistant genes are horizontal and vertical transferred from resistance to sensitive strains [25] causing the dissemination of antibiotic resistance within bacterial pathogen natural and hospital environment [19]. Spread of that kind of strains may create serious therapeutic problems in the future [26].

References

1. Carvalho MJ, Martínez-Murcia A, Esteves AC, Correia A, Saavedra MJ (2012) Phylogenetic diversity, antibiotic resistance and virulence traits of *Aeromonas* spp. from untreated waters for human consumption. Int J Food Microbiol 159: 230-239.
2. Messi P, Guerrieri E, Bondi M (2005) Antibiotic resistance and antibacterial activity in heterotrophic bacteria of mineral water origin. Sci Total Environ 346: 213-219.
3. Esteve C, Alcaide E, Blasco M D (2009) *Aeromonas hydrophila* subsp. dhakensis Isolated from Feces, Water and Fish in Mediterranean Spain. Microbes Environ 27: 367-373.

4. Nawaz M, Sung K, Khan SA, Khan AA, Steele R (2006) Biochemical and molecular characterization of tetracycline-resistant *Aeromonas veronii* isolates from catfish. Appl Environ Microbiol 72: 6461-6466.
5. Janda JM, Abbott SL (2010) The genus *Aeromonas*: Taxonomy, pathogenicity, and infection. Clin Microbiol Rev 23: 35-73.
6. Fleming A (1964) Penicillin. The Nobel Prize in Physiology or Medicine 1945.
7. Skwor T, Shinko J, Augustyniak A, Gee C, Andraso G (2014) *Aeromonas hydrophila* and *Aeromonas veronii* predominate among potentially pathogenic ciprofloxacin- and tetracycline-resistant *Aeromonas* isolates from Lake Erie. Appl Environ Microbiol 80:841-848.
8. Wu CJ, Wang HC, Chena PL, Chang MC, Sunny HS, et al. (2013) AQU-1, a chromosomal class C beta-lactamase, among clinical *Aeromonas dhakensis* isolates: Distribution and clinical significance. Int J Antimicrob Agents 42: 456-461.
9. Girlich D, Poirel L, Nordmann P (2011) Diversity of clavulanic acid-inhibited extended-spectrum Beta-lactamases in *Aeromonas* spp. from the Seine River, Paris, France. Antimicrob Agents Chemother 55: 1256-1261.
10. Choi JY, Kim Y, Ko EA, Park YK, Jheong WH, et al. (2012) *Acinetobacter* species isolates from a range of environments: species survey and observations of antimicrobial resistance. Diagn Microbiol Infect Dis 74: 177-180.
11. The European Committee on Antimicrobial Susceptibility Testing (2014) Breakpoint tables for interpretation of MICs and zone diameters.
12. Meletis G, Bagkeri M (2013) *Pseudomonas aeruginosa*: Multi-Drug-Resistance Development and Treatment Options. Infect Cont 33-56.
13. Jeong SH, Bae IK, Park KO, An YJ, Sohn SG, et al. (2006) Outbreaks of imipenem-resistant *Acinetobacter baumannii* producing carbapenemases in Korea. J Microbiol 44: 423-431.
14. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, et al. (2009) Characterization of a New Metallo β Lactamase gene, blaNDM-1, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence Type 14 from India. Antimicrob Agents Chemother 53: 5046-5054.
15. Bakour S, Garcia V, Loucif L, Brunel JM, Gharout-Sait A, et al. (2015) Rapid identification of carbapenemase-producing *Enterobacteriaceae*, *Pseudomonas aeruginosa* and *Acinetobacter baumannii* using a modified Carba NP test. New Microbes New Infect 7: 89-93.
16. Aminov RI (2009) The role of antibiotics and antibiotic resistance in nature. Environ Microbiol 11: 2970-2988.
17. Queenan AM, Bush K (2007) Carbapenemases: the versatile β Lactamases. Clin Microbiol Rev 20: 440-458.
18. O'Toole DK (2014) The Natural Environment May Be the Most Important Source of Antibiotic Resistance Genes. MBio 5: 285-314.
19. Czekalski N, Sigdel R, Birtel J, Matthews B, Bürgmann H (2015) Does human activity impact the natural antibiotic resistance background? Abundance of antibiotic resistance genes in 21 Swiss lakes. Environ Int 81: 45-55.
20. Figueira V, Vaz-Moreira I, Silva M, Manaia CM (2011) Diversity and antibiotic resistance of *Aeromonas* spp. in drinking and waste water treatment plants. Water Res 45: 5599-5611.
21. Walsh TR, Stunt RA, Nabi JA, MacGowan AP, Bennett PM (1997) Distribution and expression of beta-lactamase genes among *Aeromonas* spp. J Antimicrob Chemother 40:171-178.
22. Neuwirth C, Siebor E, Robin F, Bonnet R (2007) First occurrence of an IMP metallo- β -lactamase in *Aeromonas caviae*. IMP-19 in an isolate from France. Antimicrob Agents Chemother 51: 4486-4488.
23. Libisch B, Giske CG, Kovacs B, Toth TG, Füzi M (2008) Identification of the first VIM metallo beta lactamase-producing multiresistant *Aeromonas hydrophila* strain. J Clin Microbiol 46: 1878-1880.
24. Wellington EMH, Boxall AB, Cross P, Feil EJ, Gaze WH, et al. (2013) The role of the natural environment in the emergence of antibiotic resistance in Gram-negative bacteria. Lancet Infect Dis 13: 155-165.
25. Ashbolt NJ, Amézquita A, Backhaus T, Borriello P, Brandt KK, et al. (2013) Human Health Risk Assessment (HHRA) for Environmental Development And Transfer Of Antibiotic Resistance. Environ Health Perspect 121: 993-1001.
26. Hsieh WS, Wang NY, Jou-An F, Weng LC, Wu HH (2015) Identification of DHA-23, a novel plasmid-mediated and inducible AmpC beta-lactamase from *Enterobacteriaceae* in Northern Taiwan. Front Microbiol 6: 43-50.