

Anti-Inflammatory Activity and Cytotoxicity of the Starfish Extracts on Cancer Cells in Culture

Jumeri Mangun Wikarta^{1,2} and Sang Moo Kim^{2*}

¹Department of Agroindustrial Technology, Faculty of Agricultural Technology, Gadjah Mada University, Bulaksumur, Yogyakarta, Indonesia

²Department of Marine Food Science and Technology, Gangneung, Wonju National University, Gangneung 210-702, Republic of Korea

Abstract

Background: The starfishes were shown to contain the bioactive compounds which exhibit various biological activities including cytotoxicity, antiinflammatory, hemolytic, antibacterial, antiviral, and antifungal effects.

Objective: To determine the anti-inflammatory activity and cytotoxicity of the extracts of four different starfishes.

Materials and methods: The homogenized fresh body component of starfishes were extracted and concentrated. The EtOH extract of *A. pectinifera* with high extraction yield and total phenol was partitioned using different extraction solvents. The CHCl₃ fraction was further fractionated using thin-layer chromatography, silica-gel chromatography, gel filtration, and recycling preparative HPLC, in that order.

Results: Fraction SAP_{4,2,5} exhibited strong anti-inflammatory activity without cytotoxic effect up to a concentration of 25 µg/mL, and high cytotoxicity, with IC₅₀ values of 19.4, 34.8, and 32.7 µg/mL against AGS, DLD-1, and HeLa cells, respectively. These biological activities could be due to the presence of lathosterol, 13-tetradecen-1-ol acetate, and stigmast-7-en-3-ol (3β,5α,24S)-like compounds.

Conclusion: Starfish has a great potential as a resource for natural health products because of its strong anti-inflammatory and anticancer activities.

Keywords: Anti-inflammation; Cytotoxicity; Extract; Starfish; Steroid

Introduction

Many new drugs derived from secondary metabolites have been applied in the treatment and/or prevention of various diseases. However, these drugs have various and severe adverse effects such as nausea, vomiting, edema, and diarrhea. Therefore, naturally occurring agents with high effectiveness and no side-effects are desirable chemical therapeutics.

Inflammation is a complex pathophysiological process that is mediated by a variety of signaling molecules produced by leukocytes, macrophages, mast cells, and platelets, among other cells. Macrophages play an important role in inflammatory disease through releasing factors, such as the free radical nitric oxide (NO), prostaglandin mediators, and cytokines, which are involved in the immune response [1]. Excessive production of the inflammatory mediators is associated with many diseases, such as rheumatoid arthritis, chronic hepatitis, and pulmonary fibrosis [2]. Hence, inhibiting the production of inflammatory mediators in response to inflammatory stimuli might be a useful therapeutic strategy in inflammatory diseases.

Nitric oxide (NO) is a diatomic free radical produced from L-arginine by constitutive (cNOS) and inducible nitric oxide synthase (iNOS) in numerous mammalian cells and tissues. Nitric oxide (NO), superoxide (O₂⁻) and their reaction product peroxynitrite (ONOO⁻) may be generated in excess during the host response against viral and antibacterial infections and may contribute to pathogenesis by promoting oxidative stress, tissue injury and even cancer [3].

Marine organisms are excellent sources of structurally diverse molecules that are potentially valuable as drugs. Many natural marine products have been designated as sources of new anti-inflammatory factors and cancer chemopreventive agents [4]. In particular, the starfishes was considered as one of the most promising sources of such agents due to the variety of species and applications of its products.

Starfish (sea star) is an invertebrate belonging to the class Asteroidea, phylum Echinodermata, of which over 1500 species are

widely distributed in most of the oceans of the world. In recent years, starfish have attracted much attention due to their large scale outbreaks in coastal areas, which led to severe economic damage to the fishery and aquacultural grounds of benthic shellfishes, including mussels, oysters and scallops. Moreover, the industrial applications of the secondary metabolites of starfish have received attention worldwide. Steroidal glycosides, which are composed of a polyhydroxylated steroidal aglycone and a carbohydrate portion containing only one or two monosaccharide units, are a growing subgroup of the active glycoside compounds that have been isolated from starfishes [5]. Starfishes have also been reported to have antibacterial [6], antiviral [7], cytotoxic [8] and apoptotic [9], neurogenic [10], antifungal [11], antityrosinase [12], anti-inflammatory [13], and anticancer [14,15] activities.

The objective of this study was to determine and compare the anti-inflammatory activity and cytotoxicity against cancer cells of four starfishes, *Asterina pectinifera*, *Asterina amurensis* (the purple and yellow types) and *Distolasterias nipon*. Because *A. pectinifera* starfish is abundant and ubiquitous and has high levels of polyphenol and saponin, the anti-inflammatory and anticancer compounds of its species were purified, isolated, and identified.

Materials and Methods

Animal material

Specimens of *Asterina pectinifera*, purple-type *Asterina amurensis*,

*Corresponding author: Sang Moo Kim, Department of Marine Food Science and Technology, Gangneung, Wonju National University, Gangneung 25457, Republic of Korea, Tel: +82336402343; Fax: +82336402850; E-mail: smkim@gwnu.ac.kr

Received May 03, 2016; Accepted May 13, 2016; Published May 18, 2016

Citation: Wikarta JM, Kim SM (2016) Anti-Inflammatory Activity and Cytotoxicity of the Starfish Extracts on Cancer Cells in Culture. Med chem (Los Angeles) 6: 331-338. doi:10.4172/2161-0444.1000366

Copyright: © 2016 Wikarta JM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

yellow-type *Asterina amurensis*, and *Distolasterias nipon* were collected of the eastern coast off the Korean peninsula. The animals were brought to the laboratory in a fresh condition and were rapidly dissected and rinsed using tap water to eliminate contaminants, and were then stored at -40°C until use. Immediately before use, the starfishes were thawed at 4°C overnight and then were sliced into small pieces using a knife. The starfish samples were placed in a stainless-steel container, frozen using liquid nitrogen, and then ground using a blender.

Reagents

Dimethyl sulfoxide (DMSO), lipopolysaccharide (LPS) from *Escherichia coli* 0127:B8, N_ω-nitro-L-arginine methyl ester hydrochloride (L-NAME), Griess reagent (0.1% naphthylethylenediamine and 1% sulfanilamide in 5% H₃PO₄), and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazoliumbromide (MTT) were purchased from the Sigma Chemical Co. (St. Louis, MO, USA). RPMI 1640 medium and fetal bovine serum (FBS) were purchased from Lonza (Walkersville, MD, USA). All of the other commercially available chemicals and reagents that were used were of analytical grade.

Cell lines and culture

The macrophage cell line RAW 264.7 (Korean Cell Line Bank, Seoul, Korea) was grown in plastic culture flasks in RPMI-1640 medium containing L-glutamine and supplemented with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin under 5% CO₂ at 37°C. After 3 days, the cells were removed from the culture flask by scraping and centrifuging for 3 min at 6,574×g. The medium was removed and the cells were then resuspended using fresh RPMI-1640 medium. Three cancer cell lines, a human gastric carcinoma line (AGS), a human colon cancer line (DLD-1), and a human cervical cancer line (HeLa) (ATCC, Rockville, MD, USA) were grown in RPMI-1640 medium supplemented

with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin at 37°C in a humidified atmosphere of 5% CO₂. The concentration of RAW 264.7 cells was adjusted to 1 × 10⁶ cells/mL and that of the AGS cells, DLD-1 cells, and HeLa cells was adjusted to 1 × 10⁵ cells/mL using the same medium.

Preparation and purification of starfish extract

The flow chart for the purification of starfish bioactive compounds is shown in Figure 1. The homogenized fresh body component of the starfishes was extracted using 95% EtOH and H₂O in cold room for 6 h, which was repeated three times. The extracts were combined and filtered. The filtrate was concentrated in vacuo at 40°C to the point of dryness, yielding the crude extract. Because *A. pectinifera* is abundant and resulted in high extraction yield as well as high amount of total phenol (Table 1), this species was chosen for further analysis. The EtOH-extracted material of *A. pectinifera* was suspended in H₂O. This solution was partitioned using C₆H₁₄ to yield a C₆H₁₄ fraction. Next, the aqueous layer was successively partitioned using CHCl₃, EtOAc, and BuOH, in that order. After removing each solvent by vacuum evaporation, five fractions were obtained and were hereafter referred to as the C₆H₁₄, CHCl₃, EtOAc, BuOH, and H₂O fractions.

Because the CHCl₃ fraction exhibited the strongest anti-inflammatory and anticancer activities (Table 2), this fraction was coarsely fractionated using silica-gel (Merck KgaA, Darmstadt, Germany) column (3.0 × 50.0 cm) chromatography, using step-gradient elution with a CHCl₃:EtOH solvent system of increasing polarity (100:0 to 0:100, v/v, 400 mL) to yield 9 fractions (SAP₁-SAP₉). Each CHCl₃:EtOH fraction was concentrated using a vacuum evaporator (BUCHI Labortechnik AG, Postfach, Switzerland) and was finally vacuum-dried in a vacuum drying oven (Hanil, Gongju, Korea). The fraction with the highest yield, SAP₄, was further fractionated




Figure 1: The flow chart for the purification of starfish bioactive compounds.

Starfish	Extraction yield (%)		Total phenol (mg GAE/g)		Crude saponin (%)	
	EtOH	H ₂ O	EtOH	H ₂ O	EtOH	H ₂ O
<i>A. pectinifera</i>	6.5	10.6	47.6	64.3	17.2	13.4
<i>A. amurensis</i> (purple)	5.5	10.7	30.2	53.7	20.7	10.8
<i>A. amurensis</i> (yellow)	4.7	9.2	29.3	48.2	17.8	14.1
<i>D. nipon</i>	3.9	6.4	30.9	55.4	15.6	17.5

*Gallic acid equivalent

Table 1: The chemical composition of the starfish extracts.

Fraction	Yield (% w/w)	Anti-inflammatory (IC ₅₀ ^a µg/mL)	Anticancer (IC ₅₀ ^b µg/mL)		
			AGS	DLD	HeLa
EtOH		292.4	269.4	333.0	306.9
C ₆ H ₄	6.5 ^a	101.8	174.3	150.0	301.5
CHCl ₃	33.8 ^b	57.3	71.8	71.8	77.8
EtOAc		84.1	128.0	128.0	205.6
BuOH	11.3 ^b	115.0	129.0	129.0	159.4
H ₂ O	0.1 ^b	75.4	>400	>400	>400
L-NAME ^c	2.1 ^b	50.0	-	-	-
5 Fluorouracyl ^d	31.4 ^b	-	5.5	34.5	3.4
Paclitaxel ^d		-	3.2	24.6	2.2

^aPercentage of the fresh starfish *A. pectinifera*

^bPercentage of the dried EtOH extract

^cAnti-inflammatory positive control

^dAnticancer positive control

Table 2: The anti-inflammatory and anticancer activities of starfish extracts.

using LH-20 (3.5 × 36 cm) gel filtration chromatography using MeOH as the eluent to yield two fractions, SAP_{4.1} and SAP_{4.2}. Fraction SAP_{4.2} was further fractionated using a recycling preparative HPLC (Model LC-9104; Japan Analytical Industry Co. Ltd., Tokyo, Japan) equipped with a JAIGEL-GS310 column (21.5 × 500 mm). Elution was performed using CHCl₃:MeOH:H₂O (5:1:0.1) (v/v/v) with monitoring at 254 nm yielding five fractions (SAP_{4.2.1}-SAP_{4.2.5}). The solvent in each fraction was removed using a vacuum evaporator and the samples were finally vacuum-dried in a vacuum drying oven. The fractions were then utilized in anti-inflammatory and anticancer assays.

Identification of the purified compounds

The purified compounds were identified through GC/MS analysis using a GC 7890A system equipped with a quadrupole MS 5975C detector and an HP-35 column (30 m × 0.25 mm i.d., 0.25 µm film thickness) (Agilent Technologies, Waldbronn, Germany). The injection volume and port temperature were 2 µL and 250°C, respectively. The temperature of the column was started at 100°C for 2 min, raised to 300°C at 20°C/min and held for 6 min. Helium was used as the carrier gas, delivered at a linear flow rate of 1 mL/min. All of the spectra were scanned within the range of 33-600 m/z. The structures of the metabolites were deduced by comparing their spectroscopic data with those reported in the library.

Total phenolic and crude saponin contents

The total phenolic content was determined using the Folin-Ciocalteu method [16]. The extract was appropriately diluted using the extraction solvents. One hundred µL of each diluted extract was transferred into a test tube and brought to a volume of 500 µL using deionized water, after which 250 µL of Folin-Ciocalteu reagent (Sigma Chemical Co.) and 1.25 mL of a 12.5% aqueous sodium carbonate solution were added. The mixture was vortexed and then allowed to stand for 40 min at room temperature. Thereafter, the absorbance of each sample at 750 nm was measured against that of the control (500 µL of deionized water plus 1.5 mL of reagent mixture) and the blank (100 µL of sample diluent plus 1.9 mL of deionized water). A calibration curve was prepared using 10-50 µg/mL of gallic acid (Sigma Chemical Co.) dissolved in deionized water as the standard. The total phenolic

content was expressed as milligrams of gallic acid equivalent per gram of dry extract.

The crude saponin fraction was isolated according to the modified method of Ref. [17]. One gram of the starfish extract was dissolved in 125 mL of deionized water and the aqueous solution was transferred to a separating funnel. The solution was then treated twice using 125 mL of diethyl ether to remove the lipid components, and then was extracted three times using 125 mL of water-saturated butanol. The BuOH fraction was treated twice using 75 mL of deionized water to remove impurities. The crude saponin fraction was obtained by evaporating the BuOH under a reduced vacuum at 55°C and drying the sample at 105°C for 6 h; the sample was weighed after cooling in a desiccator. The yield of crude saponin was calculated as follows:

$$\text{Yield (\%)} = [W_{cs} / W_s] \times 100$$

where W_{cs} was the weight of the crude saponin sample and W_s was the weight of the powder sample used for extraction (1 g).

Anti-inflammatory activity

The anti-inflammatory activity was assayed according to the modified method of Choi and Hwang [18]. Briefly, cells were seeded in 96-well tissue culture plates (1 × 10⁵ cells/100 µL) and incubated at 37°C for 24 h in 5% CO₂. The extract sample was placed in phosphate-buffered saline (PBS) and was serially diluted. Cells were then stimulated with 200 µL solution containing the extract, 100 µL of LPS, and RPMI 1640 medium lacking phenol red for 24 h. One hundred microliters of the cell-culture supernatant was mixed with 100 µL of Griess reagent in the well of a 96-well plate, and the absorbance at 540 nm was measured 10 min later using a microplate spectrophotometer (EL-800; BioTek Instruments, Winooski, VT, USA). The percentage of nitric oxide inhibition was calculated based on the ability of the extract sample to inhibit nitric oxide formation by the cells compared with that of the control. The results were expressed as the percentage of inhibition of NO production compared with that of the control as follows:

$$\% \text{ inhibition} = [(NO_2^-)_c - (NO_2^-)_s] / (NO_2^-)_c \times 100$$

where $(NO_2^-)_c$ was the concentration of nitrite released by the cells in the absence of the extract and $(NO_2^-)_s$ was the concentration of nitrite released by the cells in the presence of the extract.

Cytotoxicity (cell viability) assay

To ensure that the observed inhibition of nitric oxide production of a sample was not false positive due to its cytotoxic effects, a cytotoxicity assay [19] was also performed following culture by measuring the extent of the mitochondrial-dependent reduction of MTT to formazan. To each well, 100 μ L of MTT (2 mg/mL) in PBS was added and the cells were incubated at 37°C for 4 h with 5% CO₂. The medium was then carefully discarded and the formed formazan salt was dissolved in 120 μ L of DMSO. The plates were placed at room temperature for 5 min by gently hand shaking to dissolve the formazan completely. A 100 μ L aliquot of the MTT formazan solution was transferred to a new 96-well tissue culture plate and then the absorbance at 540 nm was determined using a microplate reader (EL-800; BioTek Instruments). The absorbance of the control (untreated cells, without the extract) was taken as 100% viability. Viability was defined as the ratio (expressed as a percentage) of absorbance of the treated cells to that of the untreated cells.

Cytotoxicity against cancer cells

The cytotoxicity against cancer cells of the starfish extracts or fractions were assayed according to the modified method of Lee et al. [20]. Cells were cultured in RPMI-1640 medium supplemented with 10% heat inactivated FBS, 2mM glutamine, 100 μ g/mL streptomycin and 100 U/mL penicillin at 37°C in 5% CO₂. For the assays, the cells were plated in 96-well plates (10⁴ cells/well in 100 μ L of medium). After 4 h, the extract or fraction, which was dissolved in 1 \times PBS, was added to each well and the plates were incubated for 72 h. Afterward, PBS (100 μ L) containing 2 mg/mL of MTT was added to the culture medium. Three hours later, the formazan product resulting from the reduction of MTT was dissolved in DMSO, and the absorbance at 540 nm was measured using a microplate reader.

Statistical analysis

Each experiment was performed at least in triplicate, and the results were expressed as the mean values \pm SD. The differences among bioactivities of the fractions observed in the different assays were analyzed using an ANOVA, followed by least significance difference (LSD) test, with a p-value of <0.05 set as the level of significance.

Results and Discussion

Extract yield, total phenolic and crude saponin contents

The EtOH and H₂O extraction yields of the four different starfishes are shown in Table 1. The extraction yield and total phenol content of the H₂O extract of all of the starfishes were higher than those of the EtOH extracts. The higher level of yield of the H₂O extract might be due to starfishes containing more water-soluble substances, such as water-soluble lipids (ganglioside molecular species and a homogeneous ganglioside) and peptides [21]. Phenolics as well as non-phenolic compounds (sugars, organic acids, proteins and pigments) can be present in the extracts. The total phenolic concentration of the H₂O extracts of all of the starfishes was also higher than that of the EtOH extract (Table 1). This result is contrary to the finding of Nagai et al. [22], who reported that the total phenolic content of the EtOH extract of the horsetail plant *Equisetum arvense* L. was higher than that of the H₂O extract.

In contrast to the results regarding the content of total phenolics, the content of crude saponins of the EtOH extracts of the starfishes except for that of *D. nipon* was higher than that of the H₂O extracts (Table 1). The secondary metabolites of starfish include a remarkable diversity of various steroids, including sterols, polyhydroxysteroids, mono- and

biosides of polyhydroxysteroid, and steroid oligoglycosides called asterosaponins [23]. There are many reports of the anti-inflammatory and anticancer activities of bioactive compounds obtained from starfish, such as the anti-inflammatory activity of starfish steroids [13], and anticancer activity of polyhydroxylated steroids [24], cerebroside [8], and polysaccharides on human breast cancer [14] and colorectal adenocarcinoma cells [15]. Therefore, the starfish *A. pectinifera*, a carnivorous pirate invertebrate, was chosen for the extraction of anti-inflammatory and anticancer compounds using different organic solvents and for their subsequent purification.

Fractionation of the starfish extract

Because *A. pectinifera* is an abundant raw material and its EtOH extract had strong anti-inflammatory and anticancer activities, this extract was consecutively fractionated using organic solvents of increasing polarity (Figure 1). Extracting fresh starfish samples using EtOH resulted in a 6.5% (w/w) extraction yield (Table 2). It was found that partitioning the EtOH extract using C₆H₁₄ and H₂O led to relatively high yields (33.8 and 31.4%, respectively), whereas the CHCl₃, EtOAc, and BuOH fractions resulted in relatively low yields (11.3, 0.1, and 2.1%, respectively). Therefore, the compounds extracted from the starfish were distributed between polar and non-polar categories. This result is in good agreement with the findings of Higuchi et al. [10], in which ganglioside molecular species and a homogeneous ganglioside that were purified from the water-soluble lipid fractions of the chloroform/methanol extract of *A. pectinifera* starfish were characterized.

Anti-inflammatory activity

As shown in Figure 2A, incubating RAW 264.7 cells with the starfish extract had a significant inhibitory effect on LPS-induced NO production. The CHCl₃ fraction most effectively inhibited the LPS-induced NO production, with an IC₅₀ value of 57.3 μ g/mL, whereas the EtOH extract was the least effective, with an IC₅₀ value of 292.4 μ g/mL (Table 2). At 100 μ g/mL, the CHCl₃ fraction inhibited NO production by 98.4%, which was a much higher rate than those of the other fractions. These results agreed well with the results of Thao et al. [13], who reported that the CH₂Cl₂ extract of starfish *Astropecten polyacanthus* exerted anti-inflammatory effects on LPS-stimulated bone-marrow derived dendritic cells (BMDCs), with IC₅₀ values of 1.27, 8.82, and 11.48 μ g/mL for the inhibition of the production of IL-12 p40, IL-6, and TNF- α , respectively. Furthermore, 7 steroids with anti-inflammatory activity were purified from its CH₂Cl₂ extract and were identified. The decline in NO production may be attributed to the decrease in inducible NO synthase (iNOS) at the mRNA or protein level and the respective alterations of iNOS gene transcription [25]. The overexpression of pro-inflammatory cytokines, including IL-6 and TNF- α , is associated with the progression of inflammatory diseases. Hence, blocking the production of these cytokines and their signaling pathways is very important in preventing and treating inflammatory diseases [13].

Cell viability



The cytotoxic effects of the starfish extracts and the fractions on RAW 264.7 macrophages were evaluated using the MTT assay (Figure 2B). At 100 μ g/mL, the EtOH extract of starfish and the fractions except for CHCl₃ fraction, did not significantly alter cell viability over 24 h period. The viability rate of cells treated with the EtOH extract and the C₆H₁₄, EtOAc, BuOH, and H₂O fractions ranged from 81.3-98.2%, whereas that of the cells treated with the CHCl₃ fraction was 64.9%. The cytotoxicity rate of the starfish extract and the fractions was less than 93.7% that of L-NAME, the positive control, at same concentration. Steroids isolated from the EtOH extract of *A. pectinifera* exhibited

cytostatic activity [26]. Therefore, the lower viability rate of the cells treated with the CHCl₃ fraction of *A. pectinifera* compared with that of the cells treated with the other fractions might be due to its content of steroid compounds.

Cytotoxicity against cancer cells

As shown in Figure 3, the EtOH extract of starfish arrested the cell cycle progression of the AGS, DLD-1, and HeLa cancer cells in a concentration-dependent manner. At 400 µg/mL, the EtOH extract inhibited the growth of the AGS, DLD-1, and HeLa cells by 96.7, 96.5, and 92.5%, respectively. The cytotoxicity of the EtOH extract against HeLa cells was higher than the 0.7-7.8% rate of the EtOH extract of various edible seaweeds including *L. setchellii*, *M. integrifolia*, and *N. leutkeana* against the same cells at 1,000 µg/mL [27], but was lower than the 78.8 and 95.5% rate of the EtOH extract of *C. pilulifera* against HeLa cells at 200 µg/mL after treatment for 3 and 7 days, respectively [28]. The EtOH extract was then partitioned successively using C₆H₁₄, CHCl₃, EtOAc, and BuOH. The CHCl₃ fraction inhibited the growth of cancer cells significantly more strongly than did the EtOH extract or other fractions. This fraction may alter the cell-cycle regulation, thereby reducing the proliferation rate of cancer cells. The IC₅₀ values of the CHCl₃ fraction against the AGS, DLD-1, and HeLa cells were 47.2, 71.8, and 77.8 µg/mL, respectively (Table 3). This result is consistent with the

findings of Ref. [29], in which the CHCl₃ fraction of the herb *C. grandis* Osbeck (Dangyuja) had the highest anticancer activity. Because the CHCl₃ fraction had potent anti-inflammatory activity and cytotoxicity against cancer cells, it was conjectured that the active compounds responsible for both anti-inflammatory and anticancer activities of this fraction might be similar. Epidemiological studies have suggested that several human cancers are associated with chronically elevated levels of NO during infections [30]. Nitric oxide reacts rapidly with superoxide to form peroxynitrite, a powerful oxidant, which reacts with many biological molecules, possibly causing tissue damage. Therefore, chronic peroxynitrite exposure has been postulated as being associated with carcinogenesis due to its DNA damaging activities. Hence, DNA damage and inefficient DNA repair may initiate carcinogenesis [30]. Three polyhydroxysteroids among the 11 steroids that were isolated from three Pacific starfishes were found to be weakly cytotoxic against HeLa cells, with EC₅₀ values ranging from 53.0-174.4 µM, whereas the other steroids were inactive at up to 200 µM [24]. Cerebrocides purified from *A. pectinifera* starfish were also cytotoxic against two human cancer cells in a dose- and time-dependent manner up to 400 µg/mL [8]. A polysaccharide of *A. pectinifera* starfish also inhibited the growth of human breast [14] and colorectal [15] cancer cells. Therefore, steroids, spingolipids or polysaccharides obtained from starfish might act as anticancer compounds.





Purification of anti-inflammatory and anticancer compounds

The CHCl_3 fraction was further purified using a thin layer, silica-gel, and LH-20 gel filtration chromatography, and recycling preparative HPLC, in that order (Figure 1). Nine sub-fractions (SAP_1 - SAP_9) were obtained from the CHCl_3 fraction using silica-gel chromatography. Based on the extraction yield and activity of the subfractions, subfraction SAP_4 was chosen for further purification using Sephadex LH-20 gel filtration chromatography with EtOH as the eluent, which yielded two fractions, $\text{SAP}_{4.1}$ and $\text{SAP}_{4.2}$. The major fraction $\text{SAP}_{4.2}$ was then applied to recycling preparative HPLC, which yielded 5 fractions ($\text{SAP}_{4.2.1}$ - $\text{SAP}_{4.2.5}$) (Figure 4A). Fraction $\text{SAP}_{4.2.5}$ exhibited the highest anti-inflammatory activity and a high yield. Therefore, fraction $\text{SAP}_{4.2.5}$ was analyzed using gas chromatography/mass spectrometry (GC/MS) and the obtained spectra were compared with those in the Wiley7N library, thereby revealing at least 5 components (Figure 4B). The chromatogram of this fraction resulted in clear mass spectra, even for the minor peaks, with good resolution. The chemical composition of fraction $\text{SAP}_{4.2.5}$ obtained from *A. pectinifera* starfish as determined using GC/MS is presented in Table 3. Peak 2 (Figure 4B), as the

major component (57.9%), had a high (96%) amount of a compound corresponding to lathosterol ($\text{C}_{27}\text{H}_{46}\text{O}$, MW: 386.36). Lathosterol is a cholesterol-like molecule that is an example of a steroid. Peak 1 (6.8%) had a high (96%) amount of a compound corresponding to 13-tetradecen-1-ol acetate ($\text{C}_{16}\text{H}_{30}\text{O}_2$, MW: 254.22). Moreover, peak 5 (9.3%) consisted of 94% of a compound that matched stigmast-7-en-3-ol ($3\beta,5\alpha,24\text{S}$) ($\text{C}_{29}\text{H}_{50}\text{O}$, MW: 414.39).

Fraction $\text{SAP}_{4.2.5}$ exhibited strong anti-inflammatory activity and cytotoxicity against cancer cells in a dose-dependent manner (Figure 5). At 25 $\mu\text{g}/\text{mL}$, fraction $\text{SAP}_{4.2.5}$ inhibited NO release by 92.4% (IC_{50} , 14.6 $\mu\text{g}/\text{mL}$) without causing cytotoxicity (cell viability >90%) (Figure 5A), which suggested that this fraction contained NO-inhibitory compounds. However, at 50 $\mu\text{g}/\text{mL}$, fraction $\text{SAP}_{4.2.5}$ was cytotoxic (cell viability approximately 10%). Ergost-5-en-3-ol (3β), stigmasta-5,22E-dien-3 β -ol, and stigmast-5-en-3-ol (3β) purified from plant leaves exerted anti-inflammatory activity, while cholest-5-en-3-ol (3β) did not [31]. Furthermore, ergosta-7,22-dien-3-ol from spiny sea-star showed stronger anti-inflammatory activity by reducing more than 20% of NO levels than two fatty acids, *cis* 11-eicosenoic and *cis* 11,14 eicosadienoic



Peak	RT (min)	Library	Molecular Formula	Molecular weight (Dalton)	Match (%)	Area (%)
1	12.138	13-Tetradecen-1-ol acetate	C ₁₆ H ₃₀ O ₂	254.22	96	6.8
2	12.746	Lathosterol	C ₂₇ H ₄₆ O	386.36	96	57.9
3	13.599	Ergosta-7,22-dien-3-ol	C ₂₈ H ₄₆ O	398.36	70	13.9
4	15.022	Androstan-11-one,3-(acetyloxy)-17-iodo-,(17 α)-	C ₂₁ H ₃₁ IO ₃	458.13	58	5.6
5	16.802	Stigmast-7-en-3-ol, (3 β ,5 α ,24S)	C ₂₉ H ₅₀ O	414.39	94	9.3

Table 3: The chemical compositions of subfraction SAP_{4,2,5} obtained from starfish extract.

acids, in RAW 264.7 cells. However, maximum activity was obtained when both compounds tested in combination, thus suggesting a potentially synergistic activity of both classes of metabolites [32].

Fraction SAP_{4,2,5} also suppressed the proliferation of AGS, DLD-1, and HeLa cells in a dose-dependent manner (Figure 5B). At 50 μ g/mL, fraction SAP_{4,2,5} significantly inhibited the growth of AGS, DLD-1, and HeLa cells by 95.2 (IC₅₀, 19.4 μ g/mL), 82.1 (IC₅₀, 34.8 μ g/mL), and 88.5% (IC₅₀, 32.7 μ g/mL), respectively. Hence, fraction SAP_{4,2,5} inhibited the growth of AGS cancer cells more efficiently than it did that of DLD-1 and HeLa cells. Many researchers have reported that starfishes contain steroids, including sterols, polyhydroxysteroids, and saponins, which appear to have anticancer activities [33,34]. Two polyhydroxysteroids of *A. pectinifera* starfish were also cytotoxic to the HL-60 cells, with IC₅₀ values of 80.3 and 40.5 μ M [26]. A polar steroid isolated from *Evasterias* genus starfish were weakly cytotoxic to HeLa cancer cells [24]. Lemon essential oil containing 13-tetradecen-1-ol acetate showed inhibit the growth of HeLa cancer cells in a dose-dependent manner [35], whereas 5 α -cholest-7-en-3 β -ol purified from *A. pectinifera* starfish exhibited antimutagenic activity against the mutagens MNNG and NQO [36]. The polyhydroxy steroids and saponins of *A. pectinifera* starfish were also cytotoxic to the human liver carcinoma HepG2 cell line [34]. A ganglioside obtained from *A. pectinifera* starfish was neurotogenic in the rat pheochromocytoma cell line [10]. Therefore, steroids such as lathosterol and stigmast-7-en-3-ol (3 β ,5 α ,24S), and 13-tetradecen-1-ol acetate are considered the main anti-inflammatory and anticancer compounds of starfish *A. pectinifera*.

The anticancer effect of fraction SAP_{4,2,5} might be mediated by growth inhibition of cancer cells through dysregulated tubulin polymerization causing cell death by disrupting the normal microtubule dynamics required for cell division and vital interphase processes [37], and the inhibition of STAT3 proteins (signal transducers and activators of transcription), which are important for cancer cell survival and proliferation [38]. The mechanism underlying the anticancer activity of bioactive compounds might be the activation of the host immune response, which stimulates T-cell subsets and the production of cytokines (TNF- α and IFN- γ) that participate in anticancer effects [39]. Moreover, Ref. [40] reported that a reduction in the fraction of cells in the G1 phase and the appearance of a fraction of cells with a hypodiploid DNA content was associated with delphinidin treatment of human uterine carcinoma and colon adenocarcinoma cells. Another possible anticancer mechanism of the starfish extract is the effect of the steroidal constituents of starfish on the induction of apoptosis, in which steroids in the CH₂Cl₂ fraction induced apoptosis through the down-regulation of Bel-2 expression, the up-regulation of Bax expression, the cleavage of caspase-9 and -3, and the cleavage of poly (ADP-ribose) polymerase in HL-60 cells [13].

In the present study, the starfish extract exhibited strong anti-inflammatory activity and cytotoxicity against cancer cells. Overall, these findings confirmed the utility of clinical investigations of the efficacy of starfish extracts in preventing inflammation and in cancer chemotherapy. These data can be considered an important basis for proceeding to patient studies or to the formulation of drug products.

However, further *in vivo* studies using animal models and human patients are necessary to develop and exploit this nascent promise.

Conclusions

The anti-inflammatory and anticancer effects of starfish extracts were evaluated to identify the beneficial effects of this species under conditions related to inflammation and cancer prevention. The starfish extracts exhibited strong anti-inflammatory and anticancer activities against human cancer cells. These biological activities of the starfish extract could be partially explained by the presence of lathosterol, stigmast-7-en-3-ol,(3 β ,5 α ,24S)-like, and 13-tetradecen-1-ol acetate-like compounds. Based on the biofunctional activities of the starfish extract, *A. pectinifera* starfish could be utilized as a functional ingredient in food or pharmaceutical products to promote human health.

Acknowledgements

This research was supported by the Korea Sea Grant Program (GangWon Sea Grant) funded by the Ministry of Oceans and Fisheries in Korea.

References

- Hsieh SC, Fang SH, Rao YK, Tzeng YM (2008) Inhibition of pro-inflammatory mediators and tumor cell proliferation by *Anisomeles indica* extracts. *J Ethnopharmacol* 118: 65-70.
- Heinzel FP (1990) The role of IFN-gamma in the pathology of experimental endotoxemia. *J Immunol* 145: 2920-2924.
- Menichini F, Conforti F, Rigano D, Formisano C, Piozzi F, Senatore F (2009) Phytochemical composition anti-inflammatory and antitumor activities of four *Teucrium* essential oils from Greece. *Food Chem* 115: 679-686.
- Donia MS, Wang B, Dunbar DC, Desai PV, Patny A, et al. (2008) Mollamides B and C, cyclic hexapeptides from the Indonesian tunicate *Didemnum molle*. *J Nat Prod* 71: 941-945.
- Iorizzi M, Minale L, Riccio R, Debray M, Menou JL (1986) Steroidal glycosides from the starfish *Haliyle regularis*. *J Nat Prod* 49: 67-78.
- Go HJ, Jo M J, Kim GD, Park NG (2014) Biological activities of acidic extracts of the starfish *Asterina pectinifera*. *Kor J Fish Aquat Sci* 47: 122-128.
- Wang W, Li F, Alam N, Liu Y, Hong J, et al. (2002) New saponins from the starfish *Certonaroda semiregularis*. *J Nat Prod* 65: 1649-1656.
- Zuo H, Wang T, Qi Y, Zhao Q, Li Z (2013) Extraction, purification and cytotoxic activity of cerebrocides from *Asterina pectinifera*. *Adv Mat Res* 781: 682-686.
- Tang HF, Cheng G, Wu J, Chen XL, Zhang SY, et al. (2009) Cytotoxic asterosaponins capable of promoting polymerization of tubulin from the starfish *Culcita novaeguineae*. *J Nat Prod* 72: 284-289.
- Higuchi R, Inoue S, Inagaki K, Sakai M, Miyamoto T, et al. (2006) Biologically active glycosides from asteroidea, 42. Isolation and structure of a new biologically active ganglioside molecular species from the starfish *Asterina pectinifera*. *Chem Pharm Bull (Tokyo)* 54: 287-291.
- Chludil HD, Seldes AM, Maier MS (2002) Antifungal steroidal glycosides from the patagonian starfish *Anasterias minuta*: structure-activity correlations. *J Nat Prod* 65: 153-157.
- Jeong MH, Yang KM, Kim JK, Nam BH, Kim GY, et al. (2013) Inhibitory effects of *Asterina pectinifera* extracts on melanin biosynthesis through tyrosinase activity. *Int J Mol Med* 31: 205-212.
- Thao NP, Cuong NX, Luyen BT, Quang TH, Hanh TT, et al. (2013) Anti-inflammatory components of the starfish *Astropecten polyacanthus*. *Mar Drugs* 11: 2917-2926.

14. Lee KS, Shin JS, Nam KS (2011) Cancer chemopreventive effects of starfish polysaccharide in human breast cancer cells. *Biotechnol Bioproc E* 16: 987-991.
15. Lee KS, Shin JS, Nam KS (2012) Inhibitory effect of starfish polysaccharides on metastasis in HT-29 human colorectal adenocarcinoma. *Biotechnol Bioproc E* 17: 764-769.
16. Mamelona J, Pelletier E, Girard-Lalancette K, Legault J, Karboune S, et al. (2007) Quantification of phenolic contents and antioxidant capacity of Atlantic sea cucumber *Cucumaria frondosa*. *Food Chem* 104: 1040-1047.
17. Kwon JH, Bélanger JM, Paré JR (2003) Optimization of microwave-assisted extraction (MAP) for ginseng components by response surface methodology. *J Agric Food Chem* 51: 1807-1810.
18. Choi EM, Hwang JK (2005) Screening of Indonesian medicinal plants for inhibitor activity on nitric oxide production of RAW 2647 cells and antioxidant activity. *Fitoterapia* 76: 194-203.
19. Abas F, Lajis NH, Israf DA, Khozirah S, Kalsom YU (2006) Antioxidant and nitric oxide inhibition activities of selected Malay traditional vegetables. *Food Chem* 95: 566-573.
20. Lee JY, Hwang WI, Lim ST (2004) Antioxidant and anticancer activities of organic extracts from *Platycodon grandiflorum* A. De Candolle roots. *J Ethnopharmacol* 93: 409-415.
21. Lou H, Gao Y, Zhai M, Qi Y, Chen L, et al. (2008) A novel peptide from alpha5 helix of *Asterina pectinifera* cyclin B conjugated to HIV-Tat(49-57) with cytotoxic and apoptotic effects against human cancer cells. *Bioorg Med Chem Lett* 18: 4633-4637.
22. Nagai T, Myoda T, Nagashima T (2005) Antioxidative activities of extract and ethanol extract from field horsetail (tsukushi) *Equisetum arvense* L. *Food Chem* 91: 389-394.
23. Kicha AA, Ivanchina NV, Gorshkova IA, Ponomarenko LP, Likhatskaya GN, et al. (2001) The distribution of free sterol polyhydroxysteroid and steroid glycosides in various body components of the starfish *Patiria* (= *Asterina*) *pectinifera*. *Comp Biochem Physiol B Biochem Mol Biol* 128: 43-52.
24. Levina EV, Aminin DL, Koval'chuk SN, Kozhemiako VB, DyshlovoĀ SA, et al. (2010) Polar steroids from *Solaster endeca* starfish and the physiological activity of polar steroids from three starfish species. *Bioorg Khim* 36: 251-258.
25. Wu LC, Fan NC, Lin MH, Chu IR, Huang SJ, et al. (2008) Anti-inflammatory effect of spilanthol from *Spilanthes acmella* on murine macrophage by down-regulating LPS-induced inflammatory mediators. *J Agric Food Chem* 56: 2341-2349.
26. Li Z, Chen G, Lu X, Wang H, Feng B, et al. (2013) Three new steroid glycosides from the starfish *Asterina pectinifera*. *Nat Prod Res* 27: 1816-1822.
27. Yuan YV, Walsh NA (2006) Antioxidant and antiproliferative activities of extracts from a variety of edible seaweeds. *Food Chem Toxicol* 44: 1144-1150.
28. Kwon H, Bae S, Kim K, Han C, Cho S, et al. (2007) Induction of apoptosis in HeLa cells by ethanolic extract of *Corallina pilulifera*. *Food Chem* 104: 196-201.
29. Kim H, Moon JY, Mosaddik A, Cho SK (2010) Induction of apoptosis in human cervical carcinoma HeLa cells by polymethoxylated flavone-rich *Citrus grandis* Osbeck (*Dangyuja*) leaf extract. *Food Chem Toxicol* 48: 2435-2442.
30. Liu RH, Hotchkiss JH (1995) Potential genotoxicity of chronically elevated nitric oxide: a review. *Mutat Res* 339: 73-89.
31. Mujeeb F, Bajpai P, Pathak N (2014) Phytochemical evaluation, antimicrobial activity, and determination of bioactive components from leaves of *Aegle marmelos*. *Biomed Res Int* 2014: 497606.
32. Pereira DM, Correia-Da-Silva G, Valentao P, Teixeira N, Andrade PB (2014) Anti-Inflammatory Effect of Unsaturated Fatty Acids and Ergosta-7,22-dien-3-ol from *Marthasterias glacialis*: Prevention of CHOP-Mediated ER-Stress and NF-κB Activation. *PLOS ONE* 9: e88341.
33. Alla AK, Natalia VI, Trinh TTH, Anatoly IK, Pavel SD, et al. (2010) Two new asterosaponins, archasterosides A and B, from the Vietnamese starfish *Archaster typicus* and their anticancer properties. *Bioorg Med Chem* 20: 3826-3830.
34. Peng Y, Zheng J, Huang R, Wang Y, Xu T, et al. (2010) Polyhydroxy steroids and saponins from China Sea starfish *Asterina pectinifera* and their biological activities. *Chem Pharm Bull (Tokyo)* 58: 856-858.
35. Huang CY, Pu H, Zhang ZY, Song J (2010) Evaluation of antioxidant and antitumor activities of lemon essential oil. *J Med Plant Res* 4: 1910-1915.
36. Han YH, Ham JH, Lee NJ, Park CH, Shin YH, et al. (2000) Antimutagenic activity of 5alpha-cholest-7-en-3beta-ol, a new component from the starfish *asterina pectinifera*. *Biol Pharm Bull* 23: 1247-1249.
37. Cragg GM, Newman DJ (2005) Plants as a source of anti-cancer agents. *J Ethnopharmacol* 100: 72-79.
38. Pathak AK, Bhutani M, Nair AS, Ahn KS, Chakraborty A, et al. (2007) Ursolic acid inhibits STAT3 activation pathway leading to suppression of proliferation and chemosensitization of human multiple myeloma cells. *Mol Cancer Res* 5: 943-955.
39. Chen R, Meng F, Liu Z, Chen R, Zhang M (2010) Antitumor activities of different fractions of polysaccharide purified from *Ornithogalum caudatum* Ait. *Carbohydr Polym* 80: 845-851.
40. Yi W, Fischer J, Akoh CC (2005) Study of anticancer activities of muscadine grape phenolics in vitro. *J Agric Food Chem* 53: 8804-8812.

Citation: Wikarta JM, Kim SM (2016) Anti-Inflammatory Activity and Cytotoxicity of the Starfish Extracts on Cancer Cells in Culture. Med chem (Los Angeles) 6: 331-338. doi:10.4172/2161-0444.1000366

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>