

Antimicrobial Activity of the Crude Extracts and Fractions of Three *Baccharis* Species

Tatiana Zuccolotto¹, Félix Lourenço AV¹, Estevan Bruginiski¹, Bruna Alves¹, Andressa Veiga², Fabio Seigi Murakami² and Francinete Ramos Campos^{1*}

¹Department of Pharmacy, Biosciences Laboratory with Emphasis on Mass Spectrometry, Federal University of Paraná, Curitiba, Brazil

²Department of Pharmacy, Quality Control Laboratory Microbiology, Federal University of Paraná, Curitiba, Brazil

Abstract

The antimicrobial activities of crude extracts and fractions from three *Baccharis* L. species were tested against *Staphylococcus aureus* (ATCC 6538), *Escherichia coli* (ATCC 8738), *Pseudomonas aeruginosa* (ATCC 9027), and *Candida albicans* (ATCC 10231) using the microdilution plate method. The results showed that the crude extract from female *B. burchellii* Baker had moderate activity against *S. aureus* (minimum inhibitory concentration [MIC], 0.9 mg.mL⁻¹). Among the fractions obtained from this extract, the dichloromethane fraction showed the highest activity against *S. aureus* (MIC, 0.4 mg.mL⁻¹). The ethyl acetate fractions from female *B. burchellii* (MIC, 0.6 mg.mL⁻¹ and 1.2 mg.mL⁻¹, respectively) and *B. aracatubaensis* Malag (MIC, 1.1 mg.mL⁻¹ for both) were moderately effective against *S. aureus* and *P. aeruginosa*. The extracts from *B. organensis* Baker showed no significant activity against any organism tested. None of the extracts from *Baccharis* species showed any activity against *C. albicans*. In addition, the chemical investigation of the dichloromethane and ethyl acetate fractions from female *B. burchellii* was carried out, resulting in the identification of *trans*-ferulic acid, ethyl caffeoate, naringenin, and 7-hydroxy-benzaldehyde compounds. These phenolic compounds were found in other species of *Baccharis* and have been shown to possess antimicrobial activity. The results obtained in this work with respect to *B. burchellii* indicate that this species is a promising source of compounds with antimicrobial activities.

Keywords: Asteraceae; Baccharis; Antimicrobial; Phenolic compounds

Introduction

The *Baccharis* L. genus consists of about 500 species distributed exclusively in the Americas, found in the southern United States to southern Argentina and Chile [1,2]. There are about 178 described species in Brazil, mainly located in the southeastern and southern regions [3]. Species of this genus are well known for their use in folk medicine, especially in South America. These plants are used for the treatment of various diseases such as ulcers, gastritis, inflammation, diabetes, and skin infections [4-6]. Numerous biological activities have been attributed to essential oils, extracts, and compounds isolated from the *Baccharis* genus [7-9]. Campos et al. noted that several species of this genus have shown anti-inflammatory, anti-diabetic, anti-ulcer, or anti-microbial activities. However, there are very few reports on the antimicrobial activities of the genus *Baccharis* [10]. In this context, the aim of this study was to evaluate the antimicrobial activities of crude extracts (male and female specimens) and fractions (female specimens) from *Baccharis organensis* Baker, *Baccharis burchellii* Baker, and *Baccharis aracatubaensis* Malag, as well as to perform a chemical analysis of fractions obtained from female *B. burchellii*.

Materials and Methods

Chemicals and reagents

Dimethyl sulfoxide (DMSO), methanol (MeOH), ethyl acetate (EtOAc), and dichloromethane (CH₂Cl₂) were purchased from Tedia (Fairfield, OH, USA). Deuterium solvents (CDCl₃, DMSO-d₆, and CD₃OD) (≥ 99.9% D) were purchased from Cambridge Isotope Laboratories, Inc. (Andover, MA, USA). Mass spectrometry (MS)-grade methanol and acetonitrile (ACN) were obtained from Sigma-Aldrich (St. Louis, MO, USA). Distilled and deionized water was obtained using a Millipore system (Millipore Milli-RO plus, MA, USA). Tryptic soy broth (TSB), Mueller-Hinton broth (MHB), sabouraud dextrose broth (SDB), tryptic soy agar (TSA), sabouraud agar, chloramphenicol, 2,3,5-triphenyltetrazolium chloride (TTC), and ketoconazole were purchased from Millipore-Sigma (Darmstadt, Germany).

Chemical analysis

NMR data were acquired at 303 K in CDCl₃ for all compounds by using a Bruker AVANCE 600 NMR spectrometer operating at 14.1 Tesla, and ¹H and ¹³C spectra were recorded at 600.13 and 150.61 MHz, respectively. The spectrometer was equipped with a 5-mm quadrinuclear inverse detection probe with z-gradient. One-bond and long-range ¹H-¹³C correlations from the HSQC and HMBC NMR experiments were obtained with average coupling constants ¹J_(H,C) and ¹RJ_(H,C) optimized for 140 and 8 Hz, respectively. The ¹H and ¹³C NMR chemical shifts are given in ppm relative to the tetramethylsilane (TMS) signal as the internal reference, and the coupling constants (J) in Hz. Low-resolution electrospray ionization mass spectrometry (LRESIMS) experiments were performed on a Thermo LTQ XL Ion Trap, equipped with an ESI source. Silica gel 60 (70-230 mesh) and sephadex LH-20 (25-100 μm) were used for column chromatography (CC), and precoated silica gel plates (60 F₂₅₄ Merck, 0.2 mm, aluminum) were used for analytical thin layer chromatography (TLC). Gel plates were sprayed with p-anisaldehyde and heated, followed by exposure to UV_{254/366} light for visualization of compounds.

Plant material collection

Botanical materials of male and female specimens of *Baccharis* were collected separately and randomly along a transect within the same population in November 2013 in the “Morro do Canal”, Municipality

***Corresponding author:** Ramos Campos F, Department of Pharmacy, Federal University of Parana, Av. Lothario Meissner, Botanical Gardens, Curitiba, Brazil, Tel: +554133604162; E-mail: francampos@ufpr.br

Received August 24, 2016; **Accepted** August 27, 2016; **Published** August 31, 2016

Citation: Zuccolotto T, Félix Lourenço AV, Bruginiski E, Alves B, Veiga A, et al. (2016) Antimicrobial Activity of the Crude Extracts and Fractions of Three *Baccharis* Species. Med Chem (Los Angeles) 6: 557-560. doi:10.4172/2161-0444.1000399

Copyright: © 2016 Zuccolotto T, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

of Piraquara, Paraná State, Brazil. The *B. aracetubaensis* (leaves) and *B. organensis* (leaves) samples were collected at [25°30'52-48'' S/48°59'10-41'' O] and [25°30'52-39'' S/48°59'10-78'' O], respectively, at an elevation of 1200-1300 m. The *B. burchellii* (cladodes) samples were collected in the proximity of one river in [25°31'11-54'' S/49°00'21-17'' O] at an elevation of 906 m. The species were identified by Osmar dos Santos Ribas, Dr. Gustavo Heiden, and Dr. Angelo Alberto Schneider. The voucher specimens were deposited in the Botanical Museum of Curitiba (MBM), under the registration numbers: (MBM-286268/MBM-286267), (MBM-386275/MBM-386266), and (MBM-386257/MBM-386256), respectively.

The access to the botanical material was authorized and licensed by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and the Conselho de Gestão do Patrimônio Genético (CGEN/MMA) and registered as N° 010304/2013-4.

Collection and purification of the extracts

The air-dried botanical materials from the male and female specimens of *B. aracetubaensis* (0.986 kg and 1.0 kg, respectively), *B. burchellii* (1.1 kg and 1.2 kg, respectively), and *B. organensis* (0.490 kg and 0.560 kg, respectively) were extracted successively with a solution of ethanol:water (90:10, v/v), at room temperature. The solvent was removed from the extracts under reduced pressure to obtain the crude extracts Ba-M (129.1 g) and Ba-F (121.6 g), Bb-M (211.4 g) and Bb-F (230.4 g), and Bo-M (158.2 g) and Bo-F (184.7 g), respectively. The crude extracts from all the three female species were defatted with n-hexane; then, the crude extracts Bb-F and Bo-F were subjected to liquid-liquid partitioning with the solvents: CH₂Cl₂ (3 × 500 mL) to yield Bb-D (6.1 g) and Bo-D (14.5 g) fractions; EtOAc (3 × 500 mL) to yield Bb-Ae (26.7 g) and Bo-Ae (9.6 g) fractions; and remaining aqueous to yield Bb-Aq (30.5 g) and Bo-Aq (18.9 g) fractions, respectively. The crude extract of *B. aracetubaensis* was subjected to liquid-liquid partitioning with the solvents EtOAc (3 × 500 mL) to yield Ba-Ae (6.2 g) and remaining aqueous residue to yield Ba-Aq (26.2 g) fractions.

Part of the Bb-Ae fraction (4.5 g) was subjected to silica gel CC and was eluted with increasing concentrations of CH₂Cl₂ in n-hexane (100:0 to 10:90, v/v), followed by EtOAc in CH₂Cl₂ (100:0 to 30:70, v/v), and MeOH in EtOAc (100:0 to 70:30, v/v), affording 181 sub-fractions (30 mL each) that were pooled into 10 groups according to TLC analysis. Groups 2 (48.8 mg), 5 (131.8 mg), and 6 (171.9 mg) resulted in compounds 1, 2, and 3, respectively. Part of the Bb-D fraction (3.2 g) was subjected to silica gel CC and was eluted with increasing concentrations of CH₂Cl₂ in n-hexane (100:0 to 10:90, v/v), followed by EtOAc in CH₂Cl₂ (100:0 to 30:70, v/v), and MeOH in EtOAc (100:0 to 70:30, v/v), affording 66 sub-fractions (30 mL each) that were pooled into 6 groups according to TLC analysis. Group 2 (639.5 mg) resulted in compounds 2 and 4, and from of group 3 (191.3 mg) resulted in compounds 2 and 3.

Trans-Ferulic acid (1): C₁₀H₁₀O₄ - ¹H-NMR (600 MHz, CDCl₃) δ 7.03 (1H, *d*, *J* = 1.9 Hz, H-2), 7.07 (1H, *dd*, *J* = 1.9, 8.2 Hz, H-6), 6.91 (1H, *d*, *J* = 8.2 Hz, H-5), 6.28 (1H, *d*, *J* = 15.9 Hz, H-8), 7.60 (1H, *d*, *J* = 15.9 Hz, H-7), 3.92 (3H, *s*, OCH₃); ¹³C-NMR (600 MHz, CDCl₃) δ 167.1 (C-9), 148.9 (C-3), 146.7 (C-4), 127.4 (C-1), 123.0 (C-6), 114.7 (C-5), 115.8 (C-8), 109.1 (C-2), 144.6 (C-7), 56.2 (OCH₃); LRESIMS *m/z* 195 [M+H]⁺, 177 [M - H₂O]⁺ (100%).

Caffeate ethyl (2): C₁₁H₁₂O₄ - ¹H-NMR (600 MHz, CDCl₃) δ 6.87 (1H, *d*, *J* = 2.1 Hz, H-2), 7.00 (1H, *dd*, *J* = 2.1, 8.1 Hz, H-6), 7.07 (1H, *d*, *J* = 8.1 Hz, H-5), 6.25 (1H, *d*, *J* = 15.9 Hz, H-8), 7.56 (1H, *d*, *J* = 15.9 Hz, H-7), 4.25 (2H, *q*, *J* = 7.1 Hz, H-12), 1.33 (3H, *t*, *J* = 7.1 Hz, H-11).

¹³C-NMR (600 MHz, CDCl₃) δ 167.4 (C-9), 146.3 (C-3), 144.9 (C-4), 127.4 (C-1), 122.4 (C-6), 114.6 (C-5), 144.4 (C-8), 116.2 (C-7), 115.6 (C-2), 14.1 (C-12), 60.7 (C-11); LRESIMS *m/z* 207 [M - H]⁻, 179 [M - C₂H₅]⁻ (100%).

Naringenin (3): C₁₅H₁₂O₅ - ¹H-NMR (600 MHz, CDCl₃) δ 12.03 (1H, *s*, H-5), 7.32 (2H, *d*, *J* = 8.5, H-2' H-6'), 6.88 (2H, *d*, *J* = 8.5, H-5' H-3'), 5.98 (1H, *d*, *J* = 2.1, H-6), 6.00 (1H, *d*, *J* = 2.1, H-8), 5.35 (1H, *dd*, *J* = 13.1, 3.0, H-2), 3.08 (1H, *dd*, *J* = 17.2, 13.1, H-3α), 2.78 (1H, *dd*, *J* = 17.2, 3.0, H-3β); ¹³C-NMR 196.0 (C-4), 163.8 (C-5), 103.4 (C-9), 156.0 (C-4'), 127.9 (C-2' C-6'), 115.1 (C3' C-5'), 179.8 (C-10), 95.4 (C-6), 96.9 (C-8), 79.0 (C-2), 43.4 (C-3); LRESIMS *m/z*: 271 [M - H]⁻, 177 [M - C₆H₆O]⁻ (24%), 151 [M - C₇H₆O₂]⁻ (100%).

7-Hydroxy-benzaldehyde (4): C₇H₆O₂ - ¹H-NMR (600 MHz, DMSO-d₆) δ 7.76 (2H, *d*, *J* = 8.5, H-2, H-6), 6.93 (2H, *d*, *J* = 8.5, H-3, H-5), 9.76 (1H, *s*). ¹³C-NMR (600 MHz, DMSO-d₆) 115.8 (C-3, C-5), 128.3 (C-1), 132.0 (C-2, C-6), 163.2 (C-4), 191.0 (CHO). LRESIMS *m/z*: 121 [M - H]⁻, 106 [M - H₂O]⁻ (24%), 93 [M - CHO]⁻ (100%), 77 [M - H₂O - CHO]⁻ (22%).

Antimicrobial assay: The antimicrobial assays of the crude extracts (male and female specimens) and fractions (female specimens) from *Baccharis organensis*, *B. burchellii*, and *B. aracetubaensis* were performed using Clinical and Laboratory Standards Institute (CLSI) microdilution method [11]. The samples were tested against *Staphylococcus aureus* (ATCC 6538), *Escherichia coli* (ATCC 8738), *Pseudomonas aeruginosa* (ATCC 9027), and *Candida albicans* (ATCC 10231). The microbial suspensions used for inoculation were prepared at 10⁵ CFU (colony forming unit/ml) by diluting fresh cultures at McFarland 0.5 density. Positive controls used were 100 µg.mL⁻¹ chloramphenicol for bacteria and 500 µg.mL⁻¹ ketoconazole for yeast. The crude extracts and the CH₂Cl₂ fractions were solubilized in 20% MeOH and 5% DMSO, and the EtOAc and aqueous fractions were solubilized in water.

In this assay, the crude extracts were used at concentrations between 0.78 µg.mL⁻¹ and 100 µg.mL⁻¹, and the fractions were used at concentrations between 0.39 µg.mL⁻¹ and 50 µg.mL⁻¹. In each well of the microplate, was added 100 µL MHB for bacterial strains or 100 µL of SDB for yeast strain. In the first well, was added 100 µL of extracts or essential oils, and then performed serial dilutions (1:1, v/v), followed by addition of 10 µL of the inoculum into each well, and incubated at 35 °C for 20 h. After the incubation period, was added 20 µL of 0.125% TTC solution to all wells of the plates, followed by two hours of incubation. The absorbance was measured using a spectrophotometer at 540 nm. The minimal inhibitory concentration (MIC) was defined as the lowest concentration of the extract showing no visible bacterial growth after the incubation period.

Results and Discussion

Identification of the compounds

The chromatographic fractionation was achieved only for the fractions from female *B. burchellii* (Bb-Ae and Bb-D), resulting in the identification of the four compounds by LRESIMS and 1D and 2D NMR and upon comparison with previous literature. In the analysis of the Bb-Ae fraction from group 2 was identified, *trans*-ferulic acid (1) [12]; from group 5 resulted in isolation of caffeate ethyl (2) [13]; from group 6, a mixture of caffeate ethyl (2) and naringenin (3) [14]. In the analysis of the Bb-D fraction was identified from group 2, a mixture of caffeate ethyl (2) and 7-hydroxy-benzaldehyde (4) [15]; from group 3 was identified caffeate ethyl (2) and naringenin (3).

All compounds have been identified for the first time from the female cladodes of *B. burchellii* and are commonly found in this genus.

Antimicrobial activity

The antimicrobial activities of the samples from *B. organensis*, *B. burchellii*, and *B. aracetubaensis* were evaluated according to the microdilution method described by CLSI [11]. According to the results summarized in Table 1, the crude extract from female *B. burchellii* showed a MIC of 0.9 mg.mL⁻¹ against *S. aureus*, which was the highest activity among all the crude extracts analyzed. This extract was fractionated and the dichloromethane fraction (Bb-D) showed the highest antimicrobial activity against *S. aureus*, with a MIC of 0.4 mg.mL⁻¹ [16]. Caffeate ethyl, naringenin, and 7-hydroxy-benzaldehyde compounds were identified upon chemical analysis of this fraction. Furthermore, the ethyl acetate fractions from *B. burchellii* (Bb-Ae) and *B. aracetubaensis* showed moderate activity, with MIC values ranging between 0.6 and 1.2 mg.mL⁻¹. Ethyl caffeate, naringenin, and *trans*-Ferulic acid compounds were identified in the Bb-Ae fraction. Campos et al. had reported that the extracts and/or fractions from this genus are constituted mainly of phenolic compounds such as flavonoids, phenolic acids, and terpenes, and these possess antimicrobial activity [10], corroborating with the results obtained in this work. According to a survey conducted by Coppo and Marchese, the antibacterial activity of polyphenols can be attributed mainly to flavonols, flavones, isoflavones, flavanones, and flavan-3-ol [17]. Among the compounds tested against *S. aureus*, naringenin demonstrated strong antimicrobial activity [18-20]. Rangel observed antimicrobial activity of the extracts from *Baccharis nitida* against *S. aureus* strains [21]. Other compound classes, such as diterpenes, identified from *B. dracunculifolia*, *B. grisebachii*, *B. trimera*, *B. incarum*, and *B. dentata*, also showed activity against *S. aureus* strains [22-26].

In tests conducted using samples from *Baccharis* against *Pseudomonas aeruginosa*, MIC between 1.1 and 26.5 mg.mL⁻¹ was obtained (Table 1). The fractions that showed moderate activities were

Ba-Ae (MIC, 1.1 mg.mL⁻¹) and Bb-Ae (MIC, 1.2 mg.mL⁻¹), which were from *B. aracetubaensis* and *B. burchellii*, respectively [16]. Previous studies have reported antimicrobial activities against *P. aeruginosa* in other species of *Baccharis* such as *B. dracunculifolia*, *B. articulata* [27,28], and *B. nitida* [21]. In the plant kingdom, phenolic compounds are involved in the plant defense; and since they are synthesized in response to microbial infections [29], they can also be effective antimicrobials against a wide variety of microorganisms [30,31].

Conclusion

The crude extracts and fractions from *Baccharis aracetubaensis*, *B. burchellii*, and *B. organensis* showed significant antibacterial activity against tested strains. The highest activity against *S. aureus* was exhibited by the dichloromethane fraction from female *B. burchellii*, and moderate activity was observed in the crude extract from its male specimens and the ethyl acetate fraction from its female specimens. The ethyl acetate fractions from female *B. burchellii* and *B. aracetubaensis* showed moderate activity against *P. aeruginosa*. Extracts from *B. organensis* showed no significant activity against any organism tested. Neither the extracts nor fractions from *B. organensis*, *B. aracetubaensis*, or *B. burchellii* showed antifungal activity. Phenolic derivate compounds identified in the dichloromethane and ethyl acetate fractions from *B. burchellii* were the *trans*-ferulic, ethyl caffeate, naringenin, and 7-hydroxy-benzaldehyde compounds. These phenolic compounds were found in other species of the *Baccharis* and have been shown to possess antimicrobial activity. In this context, the results obtained in this work, with respect to *B. burchellii*, indicate that this species is a promising source of compounds with antimicrobial activity.

Acknowledgements

The authors thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), UFPR, Fundação Araucária for financial support and fellowships, the MBM for collection of botanical material, and Dr. Angelo Alberto Schneider (UNIPAMPA-RS) for *B. burchellii* identification. CGEN/CNPq (Conselho de Gestão do Patrimônio Genético) by authorization (nº 010304/2013-4).

Species	Extract and Fraction	Antimicrobial Activity (MIC mg.mL ⁻¹)			
		<i>S. aureus</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>C. albicans</i>
<i>Baccharis aracetubaensis</i>	Ba-M	8.0	16.0	8.0	---
	Ba-F	9.8	19.7	19.7	---
	Ba-Ae	1.1	4.5	1.1	---
	Ba-Aq	13.2	26.5	26.5	---
<i>Baccharis burchellii</i>	Bb-M	4.0	14.8	3.7	---
	Bb-F	0.9	8.1	4.0	---
	Bb-D	0.4	2.9	2.9	---
	Bb-Ae	0.6	4.9	1.2	---
	Bb-Aq	3.2	26.0	6.5	---
<i>Baccharis organensis</i>	Bo-M	3.6	14.4	14.4	---
	Bo-F	4.0	16.0	16.0	---
	Bo-D	2.6	10.4	5.2	---
	Bo-Ae	5.6	11.7	5.6	---
	Bo-Aq	6.6	26.4	26.4	---

Ba-M and Ba-F: crude extract from *B. aracetubaensis* male and female, respectively; Ba-Ae and Ba-Aq: fractions ethyl acetate and aqueous from *B. aracetubaensis* female, respectively; Bb-M and Bb-F: crude extract from *B. burchellii* male and female, respectively; Bb-D, Bb-Ae and Bb-Aq: fractions dichloromethane, ethyl acetate and aqueous from *B. burchellii* female, respectively; Bo-M and Bo-F: crude extract from *B. organensis* male and female, respectively; Bo-D, Bo-Ae and Bo-Aq: fractions dichloromethane, ethyl acetate and aqueous from *B. organensis* female, respectively; ---: Not activity; Positive control for antifungal activity: ketoconazole (500 µg.mL⁻¹); Positive control for antibacterial activity: chloramphenicol (100 µg.mL⁻¹); Negative control: Methanol/DMSO/H₂O (20:5:75, v/v) or H₂O.

Table 1: Antimicrobial activity of crude extracts and fractions from *B. aracetubaensis*, *B. burchellii* and *B. organensis*.

References

- Verdi LG, Brighente IMC, Pizzolatti MG (2005) Gênero *Baccharis* (Asteraceae): Aspectos químicos, econômicos e biológicos. *Quim Nova* 28: 85-94.
- Karam TK, Dalposso LM, Casa DM, Freitas GBR (2013) Carqueja (*Baccharis trimera*): utilização terapêutica e biossíntese. *Rev Bras Plantas Med* 15: 280-286.
- Heiden G, Pirani JR (2014) Two new species of *Baccharis* subgen. *Baccharis* (Asteraceae, Astereae) with single-flowered female capitula from the Serra do Cipó, Minas Gerais, Brazil. *Phytotaxa* 164: 141-148.
- Gamberini MT, Skorupa LA, Souccar C, Lapa AJ (1991) Inhibition of gastric secretion by a water extract from *Baccharis triptera*, Mart. *Mem Inst Oswaldo Cruz* 86: 137-139.
- Abad MJ, Bermejo P (2007) *Baccharis* (Compositae): a review update. *Arkivoc* 7: 76-96.
- Trojan-Rodrigues M, Alves TLS, Soares GLG, Ritter MR (2012) Plants used as antidiabetics in popular medicine in Rio Grande do Sul, southern Brazil. *J Ethnopharmacol* 139: 155-163.
- Florão A, Budel JM, Duarte MDR, Marcondes A, Rodrigues RAF, et al. (2012) Essential oils from *Baccharis* species (Asteraceae) have anti-inflammatory effects for human cells. *J Essent Oil Res* 24: 561-570.
- Rezende TP, Corrêa JOA, Aarestrup BJV, Aarestrup FM, Sousa OV, et al. (2014) Protective Effects of *Baccharis dracunculifolia* Leaves Extract against Carbon Tetrachloride- and Acetaminophen-Induced Hepatotoxicity in Experimental Animals. *Molecules* 19: 9257-9272.
- Toyama DO, Ferreira MJP, Romoff P, Fávero OA, Gaeta HH, et al. (2014) Effect of Chlorogenic Acid (5-Caffeoylquinic Acid) Isolated from *Baccharis oxyodonta* on the Structure and Pharmacological Activities of Secretory Phospholipase A2 from *Crotalus durissus terrificus*. *Biomed Res Int* 2014: 1-10.
- Campos FR, Bressan J, Jasinski VCG, Zuccolotto T, Silva LE, et al. (2016) *Baccharis* (Asteraceae): Chemical Constituents and Biological Activities. *Chem Biodivers* 13: 1-17.
- CSLI (2012) Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically. Approved Standard. 9th edn.
- Prachayasittikul S, Suphamong S, Worachartcheewan A, Lawung R, Ruchirawat S, et al. (2009) Bioactive metabolites from *Spilanthes acmella* Murr. *Molecules* 14: 850-867.
- Uwai K, Osanai Y, Imaizumi T, Kanno S, Takeshita M, et al. (2008) Inhibitory effect of the alkyl side chain of caffeic acid analogues on lipopolysaccharide-induced nitric oxide production in RAW264.7 macrophages. *Bioorg Med Chem* 16: 7795-7803.
- Frontana-Urbe BA, Escárcega-Bobadilla MV, Estrada-Reyes R, Morales-Serna JA, Salmón M, et al. (2011) A new languidulane diterpenoid from *Salvia mexicana* var. *mexicana*. *Molecules* 16: 8866-8873.
- Wang YS, Huang R, Yang JH (2011) Chemical Constituents of *Litsea szemaonis*. *Chem Nat Compd* 47: 122-123.
- Alianni N, Kalpoutzakis E, Mitaku S, Chinou IB (2010) Composition and antimicrobial activity of the essential oils of two *Origanum* species. *J Agric Food Chem* 49: 4168-4170.
- Coppo E, Marchese A (2014) Antibacterial activity of polyphenols. *Curr Pharm Biotechnol* 15: 380-390.
- Rauha JPP, Remes S, Heinonen M, Hopia A, Kähkönen M, et al. (2000) Antimicrobial effects of Finnish plant extracts containing flavonoids and other phenolic compounds. *Int J Food Microbiol* 56: 3-12.
- Kosalec I, Pepeljnjak S, Bakmaz M, Vladimir-Knezević S (2005) Flavonoid analysis and antimicrobial activity of commercially available propolis products. *Acta Pharm* 55: 423-430.
- Scazzocchio F, D'Auria FD, Alessandrini D, Pantanella F (2006) Multifactorial aspects of antimicrobial activity of propolis. *Microbiol Res* 161: 327-333.
- Rangel D, Garcia I, Velasco J, Buitrago D (2001) Actividad antimicrobiana de los extractos etanólico, acetónico y acuoso de *Baccharis nitida* (Ruiz et Pavon) Pers. *Rev La Fac Farm* 42: 43-46.
- Feresin GE, Tapia A, Gimenez A, Ravelo AG, Zacchino S, et al. (2003) Constituents of the Argentinian medicinal plant *Baccharis grisebachii* and their antimicrobial activity. *J Ethnopharmacol* 89: 73-80.
- Zampini IC, Isla MI, Schmeda-Hirschmann G (2009) Antimicrobial and Antioxidant Compounds From the Infusion and Methanolic Extract of *Baccharis incarum* (Wedd.) Perkins. *J Chil Chem Soc* 54: 477-481.
- Diaz MAN, Rossi CC, Mendonça VR, Silva DM, Ribon AOB, et al. (2010) Screening of medicinal plants for antibacterial activities on *Staphylococcus aureus* strains isolated from bovine mastitis. *Rev Bras Farmacogn* 20: 724-728.
- Aleixo AA, Herrera KMS, Ribeiro RIMA, Lima LARS, Ferreira JMS (2013) Antibacterial activity of *Baccharis trimera* (Less.) DC. (carqueja) against bacteria of medical interest. *Rev Ceres* 60: 731-734.
- Sartor T, Xavier VB, Falcão MA, Mondin CA, Santos MA, et al. (2013) Seasonal changes in phenolic compounds and in the biological activities of *Baccharis dentata* (Vell.) GM Barroso. *Ind Crops Prod* 51: 355-359.
- Fabri RL, Nogueira MS, Dutra LB, Bouzada MLM, Scio E (2011) Potencial antioxidante e antimicrobiano de espécies da família Asteraceae. *Rev Bras Plantas Med* 13: 183-189.
- Vivot EP, Sánchez C, Cacik F, Sequin C (2012) Actividad antibacteriana en plantas medicinales de la flora de Entre Ríos (Argentina). *Cienc Exactas y Nat* 45: 165-185.
- Farah A, Donangelo CM (2006) Phenolic compounds in coffee. *Brazilian J Plant Physiol* 18: 23-36.
- Daglia M, Papetti A, Dacarro C, Gazzani G (1998) Isolation of an antibacterial component from roasted coffee. *J Pharm Biomed Anal* 18: 219-225.
- Zimmer KR, Blum-Silva CH, Souza ALK, Wulff-Schuch M, Reginatto FH, et al. (2014) The Antibiofilm Effect of Blueberry Fruit Cultivars Against *Staphylococcus epidermidis* and *Pseudomonas aeruginosa*. *J Med Food* 17: 324-331.

Citation: Zuccolotto T, Félix Lourenço AV, Bruginski E, Alves B, Veiga A, et al. (2016) Antimicrobial Activity of the Crude Extracts and Fractions of Three *Baccharis* Species. *Med Chem (Los Angeles)* 6: 557-560. doi:10.4172/2161-0444.1000399

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit>