

Antimicrobials in Boar Semen Extenders – A Risk/Benefit Analysis

Morrell JM*

Clinical Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden

Abstract

Antimicrobial agents are added to semen extenders to control the growth of microbes contaminating semen during collection. However, such a non-therapeutic use of antibiotics is not without problems, since it may contribute to the development of antibiotic resistance. This review analyses the risks and benefits of using antibiotics to control bacteria in boar semen extenders, including understanding how semen becomes contaminated and the effects of these contaminants on sperm quality. In addition, the advantages and disadvantages of using antibiotics to control bacteria will be considered, as well as possible alternatives.

How do Samples become Contaminated and with What?

Boar semen is usually collected by the gloved hand method, allowing possibilities for bacterial contamination during the collection process as well as during the subsequent processing of the semen. Bacteria such as *Staphylococcus* and *Streptococcus* spp., are found on the skin and mucosa of health animals as well as in the gut (e.g. enteric bacteria) and respiratory tract [1]. Personnel also represent a source of contamination for the semen. Moreover, environmental bacteria are found on surfaces and in the air in the processing laboratory, as well as in water for washing equipment, originating from water tanks and pipes.

Effects of Bacteria on Sperm Quality

The presence of bacteria in extended semen creates competition for nutrients [2] and also results in the production of metabolic byproducts that may harm the spermatozoa. In addition, some bacteria contain lipopolysaccharides (LPS) in their cell walls which are released when they die; these LPS also cause damage to spermatozoa. The presence of contaminant bacteria in extended boar semen is associated with a decrease in sperm motility and viability [3,4], premature acrosome reaction [5] or sperm agglutination [6]. Furthermore, bacteria may cause the production of antibodies directed against the sperm glycocalyx complex [7,8].

Why Should this be a Problem for the Female?

Since bacteria are transferred to the female reproductive tract during natural mating, several physiological mechanisms have developed to protect the female. Thus surplus semen is removed from the uterus via backflow and various immune mechanisms are activated to attack and remove bacteria before the arrival of the embryo in the uterus for implantation. The problem arises when stored semen is used (as is very common in the pig breeding industry), because the semen extenders used to maintain sperm viability and functionality also serve as nutrient media for bacterial growth. Moreover, conventional storage of extended boar semen at 16–18°C allows for some bacterial multiplication since the temperature is not sufficiently low to prevent growth [2]. Thus inseminating a stored semen dose may result in a higher bacterial load being deposited in the uterus than would be the case in natural mating. The insemination of sows with contaminated semen may be associated with vulvar discharge and returns to oestrus [9], or embryonic or foetal death, endometritis, systemic infection and/or disease in recipient female [1], or reduced litter size [10]. Deteriorating sperm motility and membrane integrity was associated with increasing bacterial loads [11].

Addition of Antibiotics to Semen Extenders

Bacterial contamination can be controlled by adding antibiotics to semen extenders. The antibiotics used and the concentrations recommended

for commercial semen doses are stipulated by national and international guidelines [12]. However, there are disadvantages to this practice, both for sperm quality and for the environment. The antimicrobials may themselves have a detrimental effect on sperm survival, thus limiting the choice of agents that can be added to semen extenders. In an effort to reduce this toxicity, a cocktail of broad spectrum, highly potent antibacterial agents are used. However, any use of antibiotics can contribute to the development of antibiotic resistance, and this resistance can, in turn, be transferred to other bacteria in other host species [13]. Thus, unfortunately, even though combinations of antibiotics may reduce sperm toxicity, they may actually contribute more to antibiotic resistance than single agents. It is essential that any extender containing antibiotics and any unused semen doses should be disposed of in an approved manner that results in the inactivation of the antibiotics before they reach the environment.

Risk/Benefit Analysis

Risk/Benefit analysis is given in Table 1.

	Risk	Benefit
No antibiotics	Bacteria compete with sperm for nutrients in semen extender, produce toxic byproducts and LPS; may cause disease in females after AI.	No antibiotic toxicity to spermatozoa; no risk of contributing to the spread of antibiotic resistance
Antibiotics	May be toxic to spermatozoa; bacteria found in semen may be resistant to the antibiotics; can lead to contamination of the environment; may lead to the development of antibiotic resistance; can release LPS.	Kill contaminating bacteria before they have a chance to grow or compete for nutrients in semen extender; bacteria are not transferred to the female via AI (unless resistant),

Table 1: Risk/Benefit analysis.

*Corresponding author: Jane M Morrell, Clinical Sciences, Swedish University of Agricultural Sciences (SLU), Box 7054, SE.75007, Uppsala, Sweden, E-mail: jane.morrell@slu.se

Received January 27, 2016; Accepted February 15, 2016; Published February 25, 2016

Citation: Morrell JM (2016) Antimicrobials in Boar Semen Extenders – A Risk/Benefit Analysis. J Antimicro 2: 107. doi:10.4172/2472-1212.1000107

Copyright: © 2016 Morrell JM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Non-antibacterial Methods of Controlling Bacterial Contamination

Strict attention to hygiene during semen collection and processing may reduce bacterial contamination [3]. The normal flora of the skin, hair and respiratory tract of the male cannot be reduced; however, personnel can minimise their own contribution through personal hygiene and the sterility of the collection equipment. Protective clothing and footwear should always be worn; it should be laundered regularly or disinfected as appropriate. Personnel with respiratory infections should preferably avoid collecting semen or should wear disposable masks. The semen collection area and equipment should be thoroughly cleaned and disinfected. However, since soaps and disinfectants can be spermicidal, residues of these substances should be avoided on surfaces that will be in contact with semen. Local practice regarding cleanliness and hygiene should be followed at all times. The use of closed semen collection devices will help to prevent contamination from the environment.

These measures will certainly reduce bacterial contamination but will not remove it completely. Physical removal of the bacteria from sperm samples, e.g. by colloid centrifugation [14,15], could be an effective alternative to adding antibiotics to semen extenders. In the reported studies, separation of spermatozoa and bacteria was more effective for some bacteria than others, depending on whether the bacteria possessed a flagellum or could attach to the spermatozoa, but also on their size and whether they could aggregate or swarm. Another determining factor appeared to be whether the colloid centrifugation was carried out immediately after semen collection or after several hours of storage. Interestingly there was some indication that bacteria that had succeeded in passing through the colloid into the sperm pellet might show reduced ability to multiply, despite the lack of antibiotics in the colloid formulation. This is an interesting observation and is in keeping with other studies on pasture contamination with faecal bacteria where it was seen that bacteria that have bound to soil particles are unable to grow. The possibility that the colloid formulation may act as a natural bacterial inhibitor warrants further investigation.

Although this review focuses on extenders for boar semen, the same principles apply to other species too, although in other species e.g. bull, the use of frozen semen may reduce the opportunities for bacterial multiplication in semen doses. Therefore, efforts to improve cryopreservation methods for boar semen to give comparable fertility results to stored semen, might also contribute to a reduction in antibiotic usage. Alternatively, the development of cold storage methods for boar

semen instead of the conventional storage at 16 to 18°C could reduce bacterial multiplication and therefore reduce antibiotic usage.

References

1. Maes D, Nauwynck H, Rijsselaere T, Mateusen B, Vyt P, et al. (2008) Diseases in swine transmitted by artificial insemination: an overview. *Theriogenology* 70: 1337-1345.
2. Althouse GC (2008) Sanitary Procedures for the Production of Extended Semen. *Reprod Dom Anim* 43: 374-378.
3. Althouse GC, Pierdon MS, Lu KG (2008) Thermotemporal dynamics of contaminant bacteria and antimicrobials in extended porcine semen. *Theriogenology* 70: 1317-1323.
4. Bussalleu E, Yeste M, Sepúlveda L, Torner E, Pinart E, et al. (2011) Effects of different concentrations of enterotoxigenic and verotoxigenic *E. coli* on boar sperm quality. *Anim. Reprod. Sci* 127: 176-182.
5. Köhn FM, Erdmann I, Oeda T, el Mulla KF, Schiefer HG, et al. (1998) Influence of urogenital infections on sperm functions. *Andrologia* 30: 73-80.
6. Monga M, Roberts JA (1994) Spermagglutination by bacteria: receptor-specific interactions. *J. Androl* 15: 151-156.
7. Auroux MR, Jacques L, Mathieu D (1991) Is the sperm bacterial ratio a determining factor in impairment of sperm motility: an in-vitro study in man with *Escherichia coli*. *Int J Androl* 47: 264-270.
8. Kurpisz M, Alexander NJ (1995) Carbohydrate moieties on sperm surface: physiological relevance. *Fertil. Steril* 63: 158-165.
9. Althouse G, Kuster C, Clark S, Weisiger R (2000) Field investigations of bacterial contaminants and their effects on extended porcine semen. *Theriogenology* 53: 1167-1176.
10. Maroto Martín LO, Muñoz EC, De Cupere F, Van Driessche E, Echemendia-Blanco D, et al. (2010) Bacterial contamination of boar semen affects the litter size. *Anim Reprod Sci* 120: 95-104
11. Sepúlveda L, Bussalleu E, Yeste M, Bonet S (2014) Effects of different concentrations of *Pseudomonas aeruginosa* on boar sperm quality. *Anim Reprod Sci* 150: 96-106.
12. (2014) Council Directive, European Union, 90/429/EEC.
13. Johansson A, Greko C, Engström BE, Karlsson M (2004) Antimicrobial susceptibility of Swedish, Norwegian and Danish isolates of *Clostridium perfringens* from poultry, and distribution of tetracycline resistance genes. *Vet Microbiol* 99: 251-257.
14. Morrell JM, Wallgren M (2011) Removal of bacteria from boar ejaculates by Single Layer Centrifugation can reduce the use of antibiotics in semen extenders. *Anim Reprod Sci* 123: 64-69.
15. Morrell JM, Klein C, Lundeheim N, Erol E, Troedsson MH (2014) Removal of bacteria from stallion semen by colloid centrifugation. *Anim Reprod Sci* 145: 47-53.

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/medicaljournals/default.aspx>

Citation: Morrell JM (2016) Antimicrobials in Boar Semen Extenders – A Risk/Benefit Analysis. *J Antimicro* 2: 107. doi: [10.4172/2472-1212.1000107](https://doi.org/10.4172/2472-1212.1000107)