

Antioxidant Activity of Methanol Extract/Fractions of Senggani Leaves (*Melastoma candidum* D. Don)

Marjoni MR* and Zulfisa A

Pharmacy Academy of Dwi Farma Bukittinggi, Bukittinggi 26121, West Sumatera, Indonesia

*Corresponding author: Marjoni MR, Pharmacy Academy of Dwi Farma Bukittinggi, Bukittinggi 26121, West Sumatera, Indonesia, Tel: +62-752-625164; Fax: +62-752-625164; E-mail: mhdriza.marjoni@gmail.com

Received date: Jun 10, 2017; Accepted date: July 31, 2017; Published date: August 04, 2017

Copyright: © 2017 Marjoni MR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Senggani (*Melastoma candidum* D. Don) leaf is one of many plants in hilly areas around the city of Bukittinggi and one of biological potencies that can be utilized primarily as antioxidant and used to treat various diseases. Some active compounds in *Melastoma* genus that has activity as an antioxidant such as flavonoids and tannins.

This study aims to examine the activity of free radical scavenging by methanol extract of senggani leaves and their fractions. The antioxidant activity of methanol extract of senggani leaves and its fractions was evaluated in vitro assays using the 2,2-diphenyl-1-picrylhydrazine (DPPH) method. The amount of antioxidant activity is characterized by the value of IC₅₀ which is concentration of extract and fractions required to inhibit 50% of DPPH free radicals. The results showed that the antioxidant activity of senggani leaves were very strong for ethyl acetate and chloroform fractions with IC₅₀ values of 43,1301 µg/ml and 43,8924 µg/ml, respectively, while for methanol fraction was strong with IC₅₀ value of 65,6521 µg/ml. While the fraction of n hexane was categorized moderate with IC₅₀ value of 122,3880 µg/ml. Senggani leaves have potential to be used as natural antioxidant in preventing free radicals.

Keywords: *Melastoma candidum* D. Don; Antioxidant; Fractionation; DPPH

free radicals, forming chelates with catalytic metal compounds, and capturing oxygen [7].

Introduction

Free radicals are atoms, groups of atoms or molecules that have one or more unpaired electrons in the outermost orbital, so they are chemically highly reactive and tend to seek electron pairs to be able to bind to achieve stability. Free radical atoms are constantly attacking body's cells including normal cells for obtaining electron pairs [1,2]. Free radicals react with biological compounds in body continuously and if they do not stop, they can damage body cells and provides harmful effects to body health. This free radical reaction will also cause various diseases such as cancer, heart, cataract, premature aging, and other degenerative diseases [3].

Based on the source, free radicals can be sourced from outside body such as motor vehicle pollution, cigarette smoke, and ultraviolet light. In addition, free radicals can also come from within body itself during the process of respiration, oxidative enzyme, as well as oxidative phosphorylation in mitochondria. Thus, even if we avoid free radicals from outside the body, free radicals will automatically be generated from normal biological processes of body. To overcome this, antidote of free radicals or antioxidants are needed that serve to inhibit or stop the negative effects caused by free radicals [4,5].

Secondary metabolite compounds derived from plants, especially phenolic compounds has potential to be natural antioxidants. Many studies suggest that phenolic group compounds such as flavonoids, phenolic acids, lignins, cinnamic acids, coumarins, tocopherols and tannins have their activities as natural antioxidants [6]. Antioxidants derived from plants are highly recommended in this case, they are safer for body and can block oxidative damage through reduction with

One of the plants that have potential to be developed as natural antioxidant is senggani leaves (*Melastoma candidum* D. Don). The preliminary test showed that the extract of senggani leaves contains phenolic compounds and strong antioxidant activity. Based on the result, this study was conducted, with the aim to determine the antioxidant activity of methanol extract of senggani leaves and its fractions.

Materials and Methods

Research methodology

The research method used is explanatory research approached by regression analysis. Explanatory research is a study conducted to determine the value of independent variables, either one or more (independent) variables without making a comparison, or relate with other variables [8].

Materials

Senggani leaves used were taken from Panta Sub Regency, Agam Province of West Sumatera Indonesia. Sampling was done purposively i.e., without comparing with other regions. Chemicals: ethanol (E. Merck), n-hexane (E. Merck), ethyl acetate (E. Merck), chloroform (E. Merck) and aquadest. The material for the radical scavenging 2,2-Diphenyl-1-Picrylhydrazil (DPPH) (Sigma. Co.).

Extraction and fractionation

250 g of fresh sengгани leaves were macerated with methanol p.a for 3 × 3 days at room temperature. The maceration results were then filtered using Whatman no 1 filter paper. The filtrate was concentrated using rotary evaporator, to obtain a pourable extract. The methanol extract (partially) was added with aquadest in volume ratio of 2: 1 and parted liquid-liquid in the Liquid-Liquid Continuous Extraction (LLCE) appliance using n-hexane, chloroform and ethylacetate solvents in volume ratio of 1: 3. The extract of the fractionation was concentrated with rotary evaporator to obtain thick extension of each fraction.

DPPH radical scavenging activity

In this method, activity of free radical scavenging extract or fractions is determined by measuring the intensity of the purple color from the DPPH methanol solution [9].

Each extract and fraction was weighed 10 mg then dissolved with methanol up to 10 ml, so we got concentration of 1 mg/ml. Then dilution was performed by adding aquadest so that the samples with concentrations of (10, 30, 50, 70, 90 µg/ml) were obtained.

For the determination of antioxidant activity, each concentrate was pipetted of 0.2 ml with a micro pipette inserted into vial, added 3.8 ml of 50 µM DPPH solution. The mixture was homogenized and left for 30 min in darkness, measured the absorption with a UV-Vis spectrophotometer at 514 nm. The antioxidant activity of the samples was determined by the amount of DPPH radical absorption resistance by calculating the percentage of DPPH absorption inhibition using the formula:

Where, A₀: DPPH radical absorption of 50 µM at 514 nm wavelength; A_s: Samples absorptions in DPPH radical of 50 µM at 514 nm wavelength.

IC₅₀ values of each extract and fractions concentrations were calculated using the linear regression formula.

Results and Discussion

The extraction of sengгани leaves was performed using maceration method with ethanol solvent. This maceration method was chosen because the process is easy and does not require high temperature so it can reduce the possibility of damage to chemical compounds that have antioxidant activity contained in sengгани leaves [10,11].

Maseration results in the form of brownish green maceration were then concentrated using a rotary evaporator equipped with a vacuum pump. The vacuum process occurred during the evaporation of solvents which allowed the solvent to evaporate at temperature below its boiling point and the process of work can take place more quickly. Evaporation of methanol solvent can be carried out below its boiling point at 55°C. This process was carried out at these temperatures to keep the active compounds contained undamaged by heating.

Sengгани leaves are macerated with methanol for 3 × 3 days. The obtained macerate evaporated using a rotary evaporator to obtain viscous methanol extract. The viscous methanol extract was taken as much as 20 g of diffraction with solvents with increasing polarity starting from n-hexane, chloroform and ethyl acetate. Each fraction then evaporated with a rotary evaporator to obtain the viscous extract from each fraction.

The first fractionation was made using n-hexane solvent. The n-hexane solvent is non-polar so that it removes fat and attracts non-polar compounds such as fatty acids, sterols, coumarins, and some terpenoids. This is followed by chloroform and ethyl acetate. Ethyl acetate is used to extract compounds with intermediate polarities such as flavonoids, tannins and some alkaloids. The viscous methanol extract which is more polar than other solvents was used to extract polar compounds such as flavonoids, glycosides, tannins, and some alkaloids [12].

Screening of phytochemical extract was performed by using tube test i.e., reacting the sample with specific reagent solution to find out the secondary metabolite content. Table 1 shows phytochemical screening results of sengгани leaves.

S. No.	Secondary Metabolite	Reactor	Description
1	Alkaloids	Mayer & Dragendorff	
2	Flavonoids	HCL and Mg	+
3	Tanin	FeCl ₃	+
4	Phenolic	FeCl ₃	+
5	Saponin	Aquades	+
6	Glycosides	Lieberman-Burchad	+

Table 1: Phytochemical screening results.

Based on the results of phytochemical screening examination, it was known that methanol extract of sengгани leaves has secondary metabolite namely flavonoida, phenolic, saponin, tannin, and glikosida compounds.

The antioxidant activity of each extract and fraction was performed using DPPH method. This method is a common and simple method used to test antioxidant activity *in vitro*. According to Hanani in 2005 the DPPH free radical absorption method is a simple, easy and can be done using a small amount of sample and a relatively short working time [13]. In addition to the simplicity of the method, this DPPH method requires few samples and chemicals and takes time not too long. Measurement of antioxidant activity was performed using UV-vis spectrophotometer at a maximum DPPH wavelength of 514 nm which is the maximum DPPH wavelength at a concentration of 50 µM. The presence of antioxidant activity from the sample resulted in a change of color of the original DPPH solution which initially in viscous purple (violet) into pale yellow [14]. The antioxidant activity of each fraction is also carried out at the same wavelength.

The principle of DPPH method is the antioxidant interaction with DPPH as free radical both by electron transfer or hydrogen radical at DPPH and will neutralize free radical from DPPH. Free radicals are highly reactive molecules because they have unpaired electrons in their outer orbitals that can react with cellular molecules by binding to cell's electrons, resulting in a chain reaction that produces new free radicals [15].

If all the electrons in DPPH free radicals become paired then color of the solution would change from dark purple to bright yellow and the absorbance measured at 514 nm wavelength [16,17]. The absorbance measurements were performed after incubation for 30 min to allow the reaction between DPPH as free radical and the tested samples. The determination of antioxidant activity values is performed

quantitatively by measuring the activity of DPPH radical capture by a compound having antioxidant activity using UV-vis spectrophotometry. The result of spectrophotometer measurement is UV-Vis spectra at visible wavelength range.

The results of this study indicate the presence of antioxidant activity seen from the decrease of absorption values of DPPH radical caused by the test samples at various concentrations and the increasing value of the percentage of inhibition produced. The change in color of DPPH from purple to a slightly yellowish after 30 min incubated was visible and can also be observed. This DPPH color change is caused by the presence of compounds in the sample used to donate its hydrogen atom to free radical of DPPH, so reducing it to a more stable form of DPPH-H (1,1-diphenyl-2-picrylhydrazine). From the measurements of methanol extract and fractions of senggangi leaves of, it showed that the higher concentration used, the lower absorbance value obtained.

The reaction scheme between free radicals and antioxidants can be seen by the following Figure 1.




Figure 1: Schematic of the reaction between free radicals and antioxidants.

The determination results of antioxidant activity of senggangi leaves extract and can be seen in Table 2.

Extract Fractions and	Concentration (µg/ml)	Inhibition average (%)	Linear Regression Equation	IC50
Methanol	5	32.317	$\hat{y}=0.299x + 30.37$	65,6521 µg/ml
	15	34.603		
	25	37.079		
	35	41.46		
	45	43.873		
Chloroform	5	26,98	$\hat{y}=23,9235 + 0,5941x$	43,8924 µg/ml
	15	32,57		
	25	38,54		
	35	45,64		
	45	50,15		
Ethyl Acetate	5	35.81	$\hat{y}=0.361x + 34.43$	43,1301 µg/ml
	15	40.635		
	25	43.556		
	35	46.286		
	45	51.048		
N Hexana	5	0.762	$\hat{y}=0.420x - 1.403$	122,3880 µg/ml
	15	5.079		
	25	8.952		
	35	12.825		
	45	17.905		

Table 2: Antioxidant activity of senggangi leaves extract and fractions.

According to Sudjana [8], the linear regression equation has positive b value, thus indicating that the curve of antioxidant inhibition values is an increasing curve. The coefficient b is the linear regression coefficient and denotes that the average change of y variable for each

variable change is by one unit. From the data on the methanol fraction, we got the value of b=+ 30.37, so it can be said that for every x (sample concentration) increased BH 1 µg/ml, then y (% inhibition) increased by 0.299.

Charts of antioxidant activity measurements of extract and fractions are shown in Figure 2.




Figure 2: IC50 values obtained based on linear regression equation for (a) Methanol (b) Chloroform (c) Ethyl acetate and (d) N hexane fractions.

The regression curve also shows that there is a close correlation between concentration and % inhibition. This is shown by the values of R (correlation coefficient), and R² (coefficient of determination) above 0.90. The values of R indicate that there is a correlation between the samples concentrations and % inhibition. From the values of R² (R square) it can be seen that there is a significant correlation between the concentration of solvent with % inhibition observed with the degree of closeness of 0, 0.986. This indicates that more than 98% of inhibition degree is affected by the concentration of the ingredients, while less than 2% is affected by other factors.

The result of methanol fraction test shows that the higher the solvent concentration, the higher the percentage of inhibition, this is due to the more samples, the higher the antioxidant content so that it also affects the level of free radical inhibition by the antioxidant agent.

The results of antioxidant activity test of Chloroform fraction by linear regression equation also has positive b value, thus indicating that curve of antioxidant inhibition value at chloroform fraction also is an increasing curve. From the data seen on the fraction of chloroform, the value obtained of $b = +23,9235$ so, it can be said for every x (samples concentrations) increased by 1 µg/ml, then y (% inhibition) increased by 0.5941.

So also with the antioxidant activity test of Ethyl Acetate Fraction, the data of linear regression equation also has positive b value, thus

indicating that the curve of antioxidant inhibition values on the ethanol fraction is also an increasing curve. From the data seen in the methanol fraction, we get the value of $b = +34,43$, so it can be said that for every x (samples concentrations) increases by 1 µg/ml, then y (% inhibition) increases by 0.361.

The regression curve also shows that there is a close correlation between concentrations and % inhibition. This is shown by the values of r (correlation coefficient), and R² (coefficient of determination) above 0.90. The values of r indicate that there is a correlation between the sample concentrations with % inhibition. From the values of R² (R square) it can be seen that there is a significant correlation between the concentration of solvent with % inhibition observed with the degree of closeness of 0.988. The test results show that the higher of the solvent concentration, the higher of the percentage of inhibition, this is due to the more samples, the higher the antioxidant content so that it also affects the level of free radical inhibition by the antioxidant agent

The parameter used to show the antioxidant activity is the values of inhibition concentration (IC₅₀) which is the concentration of antioxidant substances that can reduce 50% of free radicals. According to Jun et.al 2003, antioxidant activity is classified into 5 groups: highly active, active, moderate, weak and inactive. For more detail can be seen in the Table 3.

S. No.	Intensity	IC ₅₀ Values
1	Highly active	<50 µg/ml
2	Active	50-100 µg/ml
3	Moderate	101-250 µg/ml
4	Weak	250-500 µg/ml
5	Inactive	>500 µg/ml

Table 3: Power levels of antioxidant by DPPH method [18].

Meanwhile, according to Molyneux, IC₅₀ value of 200-1000 µg/ml was declared less active but still has potential as an antioxidant [19]. Based on the results of the research, IC₅₀ ethanol extract value is 5.030 µg/ml so it falls into a very strong antioxidant category based on the average mean of IC₅₀ indicated in Table 1, it is known that IC₅₀ values obtained show that methanol extract and fractions of senggani leaves show the existence of antioxidant activity. The fractions of chloroform and ethyl acetate have very strong antioxidant activity because it has IC₅₀ value less than 50 µg/ml. A compound is said to be very active antioxidant if its IC₅₀ value <50 µg/ml. The methanol fraction belongs to the active antioxidant category because it has IC₅₀ value between 50-100 µg/ml (65.6521 µg/ml). While for the n-hexane fraction is included into weak antioxidant activity because a substance is said to have weak antioxidant activity when it has IC₅₀ value of between 151-200 µg/ml.

Based on IC₅₀ values obtained, it is known that the fractions of ethyl acetate and chloroform have the greatest antioxidant activity. This is because the fractions of ethyl acetate and chloroform allegedly contain phenolic compound namely flavonoids in the form of aglycons, where the solubility of aglycons in flavonoids is found in the more nonpolar fraction. Possible antioxidant compounds of senggani leaves will potentially be antioxidants when extracted with semi-polar and non-polar solvents then flavonoid components attached to the glycine will be readily absorbed in a polar solvent such as water. This will lead to the occurrence of steric barriers that may inhibit reactions of free radical capturing and are less effective as antioxidants [20,21].

According to Gusrav, flavonoid compounds act as antioxidants because they have hydroxyl groups that can release protons in the form of hydrogen ions. The hydrogen ions only have one proton and do not have electrons, so in the radical electrons found on the nitrogen atoms in DPPH compound bound to hydrogen ions and produce reduced DPPH [22]. Radicals in DPPH can be reduced when reacting with a hydrogen donor contained in a phenolic compound [23].

The results of antioxidant extract test showed that antioxidant activity of Senggani leaves is very strong for ethyl acetate and chloroform fractions with the IC₅₀ values of 43,1301 µg/ml and 43,8924 µg/ml, respectively, while for methanol fraction is strong with IC₅₀ value of 65, 6521 µg/ml. While the n hexane fraction categorized moderate with IC₅₀ value of 122,3880 µg/ml. The antioxidant potency of the less active n-hexane fraction is thought to be due to the presence of intruders such as proteins, fats and other soluble compounds in non-polar solvents, in this case n-hexane solvent, thereby blocking the free radical capturing process. The presence of protein or fat compounds in the fraction may interfere the free radical capturing process by flavonoid compounds. Protein or fats in plants can provide hydrogen atoms they possessed that will bind to the hydroxyl radicals in DPPH [24].

Ethyl acetate and chloroform fractions have very strong antioxidant activity because they contain more phenolic derivatives such as tannins and flavonoids. It is suspected that the classes of compounds that provide antioxidant activity in ethyl acetate and chloroform fractions are phenolic and flavonoids. Flavonoid compounds act as antioxidants because they have hydroxyl groups that can donate protons in the form of hydrogen ions. This hydrogen ion will bind to free radical electrons present in the nitrogen atoms of DPPH compound resulting in reduced DPPH [22].

Conclusion

Senggani leaves have a very powerful antioxidant activity. Fractions with antioxidant activity ranging from the strongest to the weakest are ethyl acetate fraction, chloroform fraction, methanol and n hexane fractions.

References

1. Donatus, IA (1994) Interaction of curcumin with paracetamol study on pharmacology aspects of biological change, dissertation, faculty of post graduate UGM, Yogyakarta.
2. Gitawati R (1995) Free radicals: Nature and role in making cell death. *Mirror World Med* 102: 33-35.
3. Kikuzaki H, Hisamoto M., Hirose K, Akiyama K, Taniguchi H (2002) Antioxidant properties of ferulic acid and its related compounds. *J Agric Food Chem* 50: 2161-2168.
4. Halliwell B, Gutteridge JMC (1992) Free radical in biology and medicine (3rd edn.). Oxford University Press, NY, USA pp: 796-798.
5. Lestariana W (2003) Biochemical reviews of diet to prevent disease deficiency and degenerative disease.
6. Kahkonen MP, Hopia AI, Fuorella HC (1999) Antioxidant activity of extract containing phenolic compound. *J Agri Food Chem* 47: 3954-3962.
7. Sayuti K, Yenrina R (2015) Antioxidants, natural and synthetic. Andalas University Press, Padang.
8. Sudjana (2002) Method of statistics. Tarsito: Bandung.
9. Abe N, Murata T, Hirota A (1998) Novel DPPH Radical Scavengers, Bisorbicillinol and Demethyltrichodimerol, from a Fungus Biosci Biotech Biochem 62: 661-666.
10. Ansel HC (2005) Introduction to pharmaceutical dosage form. Jakarta: UI Press, pp: 605-606, 608.
11. Voigt R (1995) Pharmacy technology, translated by Soendani NS, Yogyakarta. UGM Press, pp: 561; 567-569; 577.
12. Sarker, Satyajit D, Latif Z, Gray AI (2006) Natural products isolation. Totowa: Humana Press.
13. Hanani EA (2005) Identification of antioxidant compounds in sponges *Callyspongia* sp. from thousand islands. *Pharmaceutical Science Magazine* 2: 127-133.
14. Permana D, Hj Lajis N, Abas F, Ghafar Othman A, Ahmad R, et al. (2003) Antioxidative constituents of *hedotis diffusa* wild. *Nat Pro Sci* 9: 7-9.
15. Ketaren S (1986) Oil processing technology and food fat. UI-Press: Jakarta.
16. Green RJ (2004) Antioxidant activity of peanut plant tissues. North Carolina State University, Raleigh.
17. Gurav S, Deshkar N, Gulkari V, Duragkar N, Patil A (2007) Free radical scavenging activity of *polygala chinensis* linn. *Pharmacologyline* 2: 245-253.
18. Jun M, Fu HY, Hong J, Wan X, Yang CS, et al. (2003) Comparison of antioxidant activities of isoflavones from kadzu root (*puerari lobata ohwi*). *J Food Sci* 68: 2117-2122.
19. Molyneux P (2004) The use of the stable free radical diphenylpicrylhydrazyl for estimating antioxidant activity. *SJST* 26: 211-219.

-
20. Kahkonen MP, Hopia AI, Fuorella HC (1999) Antioxidant activity of extract containing phenolic compound. J Agri Food Chem 47: 3954-3962.
 21. Pokorni J, Yanishlieva N, Gordon M (2001) An antioxidant in food practical application. Woddhead Publishing Ltd., England, pp: 2, 10-12, 17, 44-45, 101, 107-108.
 22. Gusrav S, Deshkar N, Gulkari V, Duragkar N, Patil A (2007) Free radical scavenging activity of polygala chinensis linn. Pharmacology 2: 245-253.
 23. Arazo M, Bello A, Rastrelli L, Montelier M, Delgado L, et al. (2011) Antioxidant properties of pulp and peel of yellow mangosteen fruits. Emir J Food Agri 23: 517-524.
 24. Pine HS (1988) Free radicals. Bandung: ITB. Organic Chemistry pp: 23-26.