

Appendicectomy for Appendicitis and Laparoscopy Training

Ray-Offor E^{*} and Jebbin NJ

Department of Surgery, University of Port Harcourt Teaching Hospital, Alakahia Rivers State, Nigeria

^{*}Corresponding author: Ray-Offor E, Department of Surgery, University of Port Harcourt Teaching Hospital, Alakahia Rivers State, Nigeria, Tel: 2348481794; E-mail: erayoffor@yahoo.com

Received date: October 12, 2016; Accepted date: January 23, 2017; Published date: January 30, 2017

Copyright: © 2017 Ray-Offor E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Globally, acute appendicitis is the most common cause of acute surgical abdomen with significant morbidity and mortality. Laparoscopy is a notable stride in the improvement of surgical care.

Aims: To study the practice of appendicectomy in the tertiary hospital setting of a developing country to know who performs it, when and how.

Patients and Methods: A retrospective study of all appendicectomies for acute Appendicitis performed in University of Port Harcourt Teaching Hospital (UPTH) from January 2007 to December 2012. Data were extracted from the theatre records on demographics, anaesthesia, cadre of surgeon, method of operation and operating time. Statistical analysis of data was done using SPSS version 20.

Results: A total of 432 appendicectomies for acute appendicitis were performed during the study period. There were 205 males and 217 females with 10 unspecified cases. Four hundred and eight (98.6%) were by open method while 6 (1.4%) underwent laparoscopic appendicectomy. Senior registrars performed most appendectomies 255 (59%) while consultant surgeons performed 83 (19.0%) cases.

Conclusions: Surgical trainees perform most of the emergency operations for acute appendicitis. The open method is routinely used however a structured training in laparoscopic appendicectomy especially for surgical trainees is recommended.

Keywords: Acute appendicitis; Appendicectomy; Laparoscopy

Introduction

Acute appendicitis is the most common surgical emergency in our environment and other parts of the world [1-2]. The common underlying pathology is occlusion of the narrow lumen which may result from the following: lymphoid hyperplasia, faecolith, adhesions and kinking from repeated episodes of catarrhal appendicitis, foreign bodies, worms and carcinoma of the base of the appendix. This inflammatory condition of the vestigial appendix remains a significant cause of morbidity and occasional mortality despite extensive studies. A diagnosis can be made by clinical assessment however accuracy is aided by additional laboratory tests, scoring systems, radiological imaging and laparoscopy.

Laparoscopic appendicectomy is gaining prominence as a preferred operative method especially with uncomplicated appendicitis. Meta-analysis of randomized controlled trials concluded that laparoscopic appendicectomy was better than the open method in terms of post-operative pain, recovery, length of hospital stay and wound infection [3-6]. Aesthetics is better achieved with the small scars of port site incisions. In addition laparoscopy allows a panoramic examination of the abdominal viscera. This is especially important in the female patient where pelvic diseases can present diagnostic dilemma.

In the general surgery practice of our institution, laparoscopic intervention has commenced but is not yet routine practice. The high

prevalence of acute appendicitis with the need for appendicectomy provides a valuable opportunity to gain proficiency in basic therapeutic laparoscopy principles before graduating to advanced laparoscopic procedures. This study in the tertiary hospital of a developing country aims to study who performs appendicectomy, when and how?

Patients and Methods

Study settings

This study was conducted in the University Teaching Hospital of the metropolitan city of Port Harcourt in Nigeria-an 840 bed tertiary health care facility that serves states of the South- south region of Nigeria especially Rivers and Bayelsa states.

Study population

All consecutive cases of appendectomy for acute appendicitis performed during the study period were included.

Study design

This is a retrospective study of all appendicectomies for acute appendicitis in University of Port Harcourt Teaching Hospital Port Harcourt Rivers State performed between January 2007 and December 2012. The theatre records of the hospital were used to extract data on cases. Data included demographics, date/time of surgery, cadre of

performing surgeon, anaesthesia, method of operation and operating time.

Statistical analysis

Statistical analysis was done using IBM SPSS (Chicago ILUSA) version 20. Values were calculated in simple percentages. The mean of age and operation time were calculated with standard deviation.

Results

A total of 432 appendicectomies for acute appendicitis were performed during the study period. There were 205 male patients and 217 female patients with 10 unspecified cases (Table 1), giving a male to female ratio of 1:1.1.

Age group(years)	Male	Female	Unspecified sex	Total	Percentage (%)
1-10	5	14	1	20	4.6
11-20	46	60	0	106	24.5
21-30	78	104	0	182	42.1
31-40	48	30	3	81	18.8
41-50	19	5	0	24	5.5
51-60	6	2	2	10	2.3
61-70	1	2	0	3	0.7
>70	0	0	0	0	0
Unspecified age	2	0	4	6	1.5
Total	205	217	10	432	100

Table 1: Age and sex distribution of cases of acute appendicitis.

The incidence of acute appendicitis was highest in the third decade of life 182 (42.1%) followed by the second decade of life 106 (24.5%). There was a mean age of 26.18 years SD 10.8. There was an average of 72 appendicectomies per annum mostly performed during the evening shift (Table 2).

Year of study	Morning shift (7.00 am -2 pm)	Evening shift (2.00 pm-9.00 pm)	Night shift (9.00 pm-7.00 am)	Unspecified	Total
2007	17	26	23	0	66
2008	20	30	20	0	70
2009	24	34	16	3	77
2010	15	27	7	2	51
2011	24	43	21	0	88
2012	21	40	19	0	80
Total	121	200	106	5	432

Table 2: Yearly incidence of acute appendicitis and time of surgery.

Three hundred and forty nine cases (81.0%) were performed by surgical trainees with senior registrars performing over half of the case load (Figure 1). There is no record of a House officer as the performing surgeon. General anaesthesia with endotracheal intubation was predominantly used.




Figure 1: Cadre of surgeon performing appendicectomies.

Most of the cases were uncomplicated acute appendicitis 375(86.8%) and appendicectomy was performed by open method using Lanz incision (Table 3). Fifty one cases of complicated appendicitis with suspected perforated appendicitis with peritonitis needed laparotomy by midline incision. Six laparoscopic appendicectomies were successfully performed by a consultant surgeon(ER-O).

Pathology (method of surgery)	Frequency	Percentage(%)
Uncomplicated appendicitis (Open-Lanz incision)	375	86.8
Complicated appendicitis (Midline laparotomy)	51	11.8
Uncomplicated appendicitis (Laparoscopic)	6	1.4
Total	432	100

Table 3: Types of surgeries performed.

The operating time ranged from 18 minutes to 102 minutes in appendicectomy for uncomplicated cases with a mean of 60.6minutes SD 33.2. For laparotomy, the mean operating time was 100.5 minutes SD 52.8.

Discussion

Appendicectomy for acute appendicectomy is one of the most common emergency surgical procedures performed worldwide. In our study there is an average annual incidence of 72 cases in comparison to studies on acute appendicitis showing 38.2 and 49.3 from tertiary hospitals in Northern and South-western parts of Nigeria respectively. The peak age incidence of acute appendicitis observed was the third decade followed by the second decade. This peak age incidence was not observed to vary with the sex of the patient however varying reports have been documented from different parts of the country [7,8]. A slight predilection for females of 1.1 : 1 was observed in this study. This

is however dissimilar to the trend of male predominance widely documented [7,9,10].

Trainee surgeons performed the highest number of appendectomies in this study with senior registrars accounting for 59% and registrars 22%. This is in line with the global trend. A UK audit of 557 hospitals observed that appendicectomies in adults were delegated to junior doctors and 53% were performed by senior house officers and 42% by registrars [11]. There was no record of a House Officer (Intern) as a performing surgeon but as an assistant surgeon. It was observed that one quarter of all appendectomies in this study were performed during the night shift (Table 2) as the major elective surgeries are performed during the morning shift. The consultants mostly perform these major surgeries or assist the senior registrars as part of their training. This trend may explain the long mean operating time as junior residents try to gain competency.

Appendicectomies were traditionally performed by the open method in 98.6% of cases and mostly under general anaesthesia. In recent times spinal anaesthesia is more frequently used by the new crop of anaesthetists except in cases of sepsis for perforated appendicitis when general anaesthesia is the preferred choice. The practice of laparoscopic appendectomy is new in our environment. A pilot study of 6 (1.4%) cases was successfully performed during the study period with extracorporeal ligation of appendix base. None was performed during the night shift. A major reason for this was the logistic challenge of nurses shift duty. In the face of competing need of emergency cases for limited theatre space, daytime shifts which are better staffed are more convenient for laparoscopic appendectomy. A dedicated theatre space for laparoscopy is ideal [12].

This study suggests the need to emphasize laparoscopy in the post-graduate training curriculum of trainee surgeons. A structured training program involves a competency framework design for preoperative assessment of a surgeon's skills before an independent laparoscopic surgery. This training includes learning anatomy specific to laparoscopic surgery for each organ, physiology of pneumoperitoneum, hand-eye coordination with a two-dimensional monitor and knowledge of instruments and equipment specific to laparoscopic surgery. The use of virtual reality simulator, porcine *in vivo* or *ex vivo* set ups and trainer boxes play vital roles in overcoming the crux of the learning curve [13,14]. It has been observed that simultaneous training of open and endoscopic surgery has a synergistic effect in the education of surgical residents as fine anatomy is learned through the magnified image on the monitor [15].

Conclusion

Appendectomy for acute appendicitis is a common abdominal surgery in our environment with trainee surgeons performing the bulk

of surgeries. A well-structured training in laparoscopic appendicectomy for trainee surgeons in line with current standard in developed countries is recommended.

References

1. Datubo-Brown DD, Adotey JM (1990) Pattern of surgical acute abdomen in the University of Port Harcourt Teaching Hospital. West Afr J Med 1: 59-62.
2. Alagoa PJ, Jebbin NJ (2010) The changing pattern of acute abdomen in Port Harcourt, Nigeria. PH Med J 4: 122-127.
3. Chung RS, Rowland DY, Li P, Diaz J (1999) A meta-analysis of randomized controlled trials of laparoscopic versus conventional appendectomy. Am J Surg 77: 250-253.
4. Garbutt JM, Soper NJ, Shannon WD, Botero A, Littenberg B (1999) Meta-analysis of randomized trials comparing laparoscopic versus open appendectomy. Surg Laparosc Endosc 9: 17-26.
5. Golub R, Siddiqui F, Pohl D (1998) Laparoscopic versus open appendectomy: a meta-analysis. J Am Coll Surg 186: 543-553.
6. Temple LK, Litwin DE, McLeod RS (1999) A meta-analysis of laparoscopic versus open appendectomy in patients suspected of having acute appendicitis. Can J Surg 42: 377-383.
7. Oguniola AS, Adeoti ML, Oyemolade TA (2010) Appendicitis: Trends in incidence, age, sex, and seasonal variations in South-Western Nigeria. Ann Afr Med 9: 213-217.
8. Ahmed SA, Makama JG, Mohammed U, Sanda RB, Shehu SM, et al. (2014) Epidemiology of appendicitis in Northern Nigeria: A 10-year review. Sub-Saharan. Afr J Med 1: 185-190.
9. Agboola JO, Olatoke SA, Rahman GA (2014) Pattern and presentation of acute abdomen in a Nigerian Teaching Hospital. Niger Med J 55: 266-270.
10. Chamisa I (2009) A clinicopathological review of 324 appendices removed for acute appendicitis in Durban, South Africa: a retrospective analysis. Ann R Coll Surg Engl 91: 688-692.
11. Lansdown MR, Gray AJ, Treasure T, Layer GT (2006) Appendicectomy: who performs it, when and how? Ann R Coll Surg Engl 88: 530-534.
12. Ray-Offor E, Okoro PE, Allison AB (2014) Pilot study on laparoscopic surgery in Port-Harcourt, Nigeria. Niger J Surg 20: 23-25.
13. Nguyen NT, Mayer KL, Bold RJ, Larson M, Foster S, et al. (2000) Laparoscopic suturing evaluation among surgical residents. J Surg Res 3: 133-136.
14. Roberts KE, Bell RL, Duffy AJ (2006) Evolution of surgical skills training. World J Gastroenterol 12: 3219-3224.
15. Kano N, Takeshi A, Kusanagi H, Watarai Y, Mike M, et al. (2010) Current surgical training in open and laparoscopic surgery. Surg Endosc 24: 2927-2929.