



Aspects of Cognition/Health Failure by HIV-infected Individuals: Amelioration through Exercise

Trevor Archer*

Department of Psychology, University of Gothenburg, Box 500, S-405 30 Gothenburg, Sweden

Editorial

A profile of progressive neurological symptoms referred to collectively as HIV-associated neurocognitive disorder (HAND) has been found to be linked to human immunodeficiency virus type 1 (HIV-1) with symptoms implicating several cognitive domains, including attention, learning, memory, affect and motor functioning [1-3]. Neuropsychological testing indicates that around fifty percent of HIV patients, in antiretroviral therapy, present evidence of mild cognitive impairment (MCI)/HIV dementia which implies that prevalence was unchanged from the pre- to the potent antiretroviral period, although incidence of MCI had increased and HIV dementia decreased [4]. In children and adolescents presenting HIV, the status of stimulant treatment for cognitive, behavioral and quality-of-life deficits remains an expectancy, rather than a guarantee of long-term improvement [5]. The syndrome, HAND, presents a spectrum of neurocognitive-neurologic deficits characterized, on the one hand, by poor concentration, memory impairments and psychological sluggishness, and on the other hand, by psychomotor retardation, deficits in fine motor control, balance and posture problems with accompanying clumsiness, and tremors. The behavioral deficits include apathy, flatness of affect, lethargy and loss of spontaneity [6]. HIV-induced metaboencephalopathy engenders HAND symptoms with immune activation arising from macrophages and microglia [7], with accompanying neurotoxin release, dendritic abnormality and neuron loss [8] and cerebellar and subcortical gray matter atrophy [9]. Su et al. [10] have shown the HIV-infected males displayed greater white matter hyperintensities that was associated independently with higher age levels, higher diastolic blood pressure and D-dimer levels and longer periods with lower CD₄₊ cell counts. Although HAND is linked generally with HIV diagnosis, for most infected individuals receiving antiretroviral therapy combined with virologic suppression the disorder is not progressive during a four-year follow-up period [11,12]; nevertheless, certain variants of HIV-linked neurocognitive deficits were shown to be related to attentional bias that included motor deficits commonly associated with HIV-induced damage to the basal ganglia and previous inability to avoid risk-taking behaviors [13].

It has been observed that age-related co-morbidities exacerbate the risk for cognitive deficits in patients with HIV diagnosis, a reality that requires consideration when examining individual-based factor affecting progression and expression, such as age, gender, ethnicity and genetics. With advancing age, individual presenting HAND are at greater for progressive deterioration of daily life activities, medical decision-making and quality-of-life [14-16]. In a study deriving an estimate of 'cognitive reserve from the Brief Intelligence Test and the Cognitive Reserve Index, it was observed that lifestyle choices affected coping-capacity to deal with HIV in aging and HIV-related neurodegeneration [17,18]. It has been shown that in a cohort of HIV patients (aged 60 years and above) that the adherence to antiretroviral therapy was related to cognitive impairment [12], apolipoprotein E4 genotype, monocyte efficacy scores, with trend associations for diabetes and low CD4+ cell count [19,20]. Plasma microRNA profiling of genes, applied to estimate neural development, cell death, neuroinflammation, cell signaling and cytokine functions [21], has provided useful biomarkers

for HAND through predicting HIV-associated cognitive impairments [22]. The Veterans Aging Cohort Study Index, developed as a risk index for HIV, in relation to global cognition, was found to vary as a function of ethnic/racial grouping [23]. In a study of HIV patients in Nigeria, it was shown that the risk of developing neurocognitive impairment differed between genders possibly due to the effects of higher plasma levels of HIV coercing the activation of circulating monocytes [24]. Male transgenic mice, inducibly expressing HIV-1 Tat, the HIV-1 transactivator of transcription, or non-expressing control mice, displayed spatial memory impairments and other cognitive deficits, together with increased vulnerability of parvalbumin-expressing and somatostatin-expressing interneurons in the hippocampus [25] and other markers of neuronal damage [26].

Physical exercise has been described as a combination of planned and organized physical activities that fulfill the intent for development and advancement several domains that include health, physical fitness, quality-of-life and functional sufficiency [27]. Exercise/activity encompasses any and all physical activity that provokes force muscular force dissembling existing inertia [28]. Muscular exertion through physical activity bestows marked benefits for cerebral integrity and neurocognition [29-31], in addition to the development and proliferation of brain structure and function, improved quality-of-life and physical endurance and strength [32]. Simonik et al. [33] have shown that the propensity and compliance to engage in exercise among individuals enduring HIV presents a dynamic and fluctuating construct that may be influenced by the episodic nature of HIV and multi-morbidity with the recommendation that enhancement of physical activity/exercise ought to maximize health outcomes. Combinations of endurance (aerobic) and resistance exercise three times weekly well-tolerated by HIV-patients and brought about improvements in cardiorespiratory fitness, strength, body composition and quality-of-life [34]. Higher levels of education, socioeconomic status, cognitively-challenging occupations and intelligence indices predict superior cognitive function and daily functional activity in HIV afflicted individual. Certainly, exercise interventions for Nigerian HIV-patients (18 years and above) improved markedly both pulmonary functions and alleviated respiratory and depressive symptoms [35]. Physical activity engagement and employment are linked to greater performance in neurocognitive tasks [36-38]. Cross-sectional studies have indicated that activity and active engagement in daily activities bolsters neurocognitive functioning [39]. As implied by Mattson [40]

*Corresponding author: Archer T, Department of Psychology, University of Gothenburg, Box 500, S-405 30 Gothenburg, Sweden, Tel: +46 31 7864694; E-mail: trevor.archer@psy.gu.se

Received May 23, 2016; Accepted May 24, 2016; Published May 31, 2016

Citation: Archer T (2016) Aspects of Cognition/Health Failure by HIV-infected Individuals: Amelioration through Exercise. HIV Curr Res 1: e101.

Copyright: © 2016 Archer T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and others [41-46], HAND patients ought to benefit from the plethora of health advantages bestowed by regular physical exercise: accelerated brain-derived factor and other trophic factor production, enhanced synaptic (even axonal) plasticity, neurogenesis and neuron arborization, increased production of anti-inflammatory cytokines and other aspects of improved neuroimmune functioning, and marked dividends for cognitive performance and affective balance. Furthermore, exercise intervention was applied to fifteen HIV-infected individuals between 35 and 51 years-of-age, and improvements in hand strength, scapular force, and several quality-of-life domains, including environment, perception of quality-of-life and global health [47,48]. Thirty HIV patients, undergoing highly active antiretroviral therapy, were either assigned to a nonlinear resistance exercise group or to a group with the exercise intervention interval covering 12 weeks [49]. After the 12-week intervention, the exercise group had augmented levels of lean body mass, with a reduction in body fat mass and body fat percentage whereas the levels of total cholesterol, low density lipoprotein cholesterol, triglyceride, and c-reactive protein were reduced, with the high density lipoprotein cholesterol level increased. Thus, the exercise intervention brought about alleviative changes in the body composition, lipid profile, and inflammation markers of HIV patients. Finally, in a study of the aerobic (endurance) plus resistance exercise training combination upon self-reported mood disturbances, e.g. reported depression and profiles of mood states, perceived stress, frequency of self-reported symptoms, and symptom-distress in Exercised or Non-exercised HIV-infected individuals, obtained significant decreases in reported depression scores and total profile of mood state. Taken together, the consensus of current findings and notions concerning intervention implies that physical exercise holds manifest benefits, certainly as an adjunctive therapy, for HIV-infected individuals presenting HANDS.

References

- Antinori A, Arendt G, Becker JT, Brew BJ, Byrd DA, et al. (2007) Updated research nosology for HIV-associated neurocognitive disorders. *Neurology* 69: 1789-99.
- Libertone R, Balestra P, Lorenzini P, Pinnetti C, Ricottini M, et al. (2014) APRI and FIB-4 scores are not associated with neurocognitive impairment in HIV-infected persons. *J Int AIDS Soc* 17: 19658.
- McArthur JC, Steiner J, Sacktor N, Nath A (2010) Human immunodeficiency virus-associated neurocognitive disorders: mind the gap. *Ann Neurol* 67: 699-714.
- Ellis R, Langford D, Masliah E (2007) HIV and antiretroviral therapy in the brain: neuronal injury and repair. *Nat Rev Neurosci* 8: 33-44.
- Sirois PA, Aaron L, Montepiedra G, Pearson DA, Kapetanovic S, et al. (2016) Stimulant Medications and Cognition, Behavior and Quality of Life in Children and Youth with HIV. *Pediatr Infect Dis J* 35: e12-8.
- Focà E, Magro P, Motta D, Compostella S, Casari S, et al. (2016) Screening for Neurocognitive Impairment in HIV-Infected Individuals at First Contact after HIV Diagnosis: The Experience of a Large Clinical Center in Northern Italy. *Int J Mol Sci* 17: 434.
- Joseph J, Colosi DA, Rao VR (2016) HIV-1 Induced CNS Dysfunction: Current Overview and Research Priorities. *Curr HIV Res*.
- Jensen BK, Monnerie H, Mannell MV, Gannon PJ, Espinoza CA, et al. (2015) Altered Oligodendrocyte Maturation and Myelin Maintenance: The Role of Antiretrovirals in HIV-Associated Neurocognitive Disorders. *J Neuropathol Exp Neurol* 74: 1093-118.
- Kallianpur KJ, Shikuma C, Kirk GR, Shiramizu B, Valcour V, et al. (2013) Peripheral blood HIV DNA is associated with atrophy of cerebellar and subcortical gray matter. *Neurology* 80: 1792-9.
- Su T, Wit FW, Caan MW, Schouten J, Prins M, et al. (2016) White matter hyperintensities in relation to cognition in HIV-infected men with sustained suppressed viral load on cART. *AIDS*.
- Gates TM, Cysique LA, Siefried KJ, Chaganti J, Moffat KJ, et al. (2016) Maraviroc-intensified combined antiretroviral therapy improves cognition in virally suppressed HIV-associated neurocognitive disorder. *AIDS* 30: 591-600.
- Sacktor N, Skolasky RL, Seaberg E, Munro C, Becker JT, et al. (2016) Prevalence of HIV-associated neurocognitive disorders in the Multicenter AIDS Cohort Study. *Neurology* 86: 334-40.
- Anderson BA, Kronemer SI, Rilee JJ, Sacktor N, Marvel CL (2015) Reward, attention, and HIV-related risk in HIV+ individuals. *Neurobiol Dis*.
- Moore RC, Fazeli PL, Jeste DV, Moore DJ, Grant I, et al. (2014) Successful cognitive aging and health-related quality of life in younger and older adults infected with HIV. *AIDS Behav* 18: 1186-97.
- Nyirenda M, Evandrou M, Mutevedzi P, Hosegood V, Falkingham J, et al. (2015) Who cares? Implications of care-giving and -receiving by HIV-infected or -affected older people on functional disability and emotional wellbeing. *Ageing Soc* 35: 169-202.
- Solomon P, O'Brien K, Wilkins S, Gervais N (2014) Aging with HIV: a model of disability. *J Int Assoc Provid AIDS Care* 13: 519-25.
- Milanini B, Ciccarelli N, Fabbiani M, Limiti S, Grima P, et al. (2016) Cognitive reserve and neuropsychological functioning in older HIV-infected people. *J Neurovirol*.
- Ciccarelli N, Grima P, Fabbiani M, Baldonero E, Borghetti A, et al. (2015) Baseline CD4(+) T-cell count and cardiovascular risk factors predict the evolution of cognitive performance during 2-year follow-up in HIV-infected patients. *Antivir Ther* 20: 433-40.
- Valcour VG (2013) HIV, aging, and cognition: emerging issues. *Top Antivir Med* 21: 119-23.
- Valcour VG, Sithinamsuwan P, Letendre S, Ances B (2011) Pathogenesis of HIV in the central nervous system. *Curr HIV/AIDS Rep* 8: 54-61.
- Kim W, Zekas E, Lodge R, Susan-Resiga D, Marcinkiewicz E, et al. (2015) Neuroinflammation-Induced Interactions between Protease-Activated Receptor 1 and Proprotein Convertases in HIV-Associated Neurocognitive Disorder. *Mol Cell Biol* 35: 3684-700.
- Asahchop EL, Akinwumi SM, Branton WG, Fujiwara E, Gill JM, et al. (2016) Plasma microRNA profiling predicts HIV-associated neurocognitive disorder. *AIDS*.
- Marquine MJ, Sakamoto M, Dufour C, Rooney A, Fazeli P, et al. (2016) The impact of ethnicity/race on the association between the Veterans Aging Cohort Study (VACS) Index and neurocognitive function among HIV-infected persons. *J Neurovirol*.
- Royal W, Cherner M, Burdo TH, Umlauf A, Letendre SL, et al. (2016) Associations between Cognition, Gender and Monocyte Activation among HIV Infected Individuals in Nigeria. *PLoS One* 11: e0147182.
- Marks WD, Paris JJ, Schier CJ, Denton MD, Fitting S, et al. (2016) HIV-1 Tat causes cognitive deficits and selective loss of parvalbumin, somatostatin, and neuronal nitric oxide synthase expressing hippocampal CA1 interneuron subpopulations. *J Neurovirol*.
- Fitting S, Knapp PE, Zou S, Marks WD, Bowers MS, et al. (2016) Interactive HIV-1 Tat and morphine-induced synaptodendritic injury is triggered through focal disruptions in Na⁺ influx, mitochondrial instability, and Ca²⁺ overload. *J Neurosci* 34: 12850-64.
- Morris M, Schoo A (2004) Optimizing exercise and physical activity in older adults. *Butterworth Heinemann, Edinburgh*.
- McArdle WD, Katch FI, Katch VI (1978) *Essentials of exercise physiology*.
- Bediz CS, Oniz A, Guducu C, Ural Demirci E, Ogut H, et al. (2016) Acute Supramaximal Exercise Increases the Brain Oxygenation in Relation to Cognitive Workload. *Front Hum Neurosci* 10: 174.
- Marks BL, Katz L, Styner M, Smith JK (2010) Aerobic fitness and obesity: relationship to cerebral white matter integrity in the brain of active and sedentary older adults. *Br J Sports Med* 45: 1208-1215.
- Segalowitz SJ (2016) Exercise and Pediatric Brain Development: A Call to Action. *Pediatr Exerc Sci* 28: 217-25.
- Deschamps A, Diolez P, Thiaudier et al (2010) Effects of exercise programs to prevent decline in health related quality of life in highly deconditioned institutionalized elderly persons: a randomized controlled trial. *Arch Intern Med* 170: 162-169.

33. Simonik A, Vader K, Ellis D, Kesbian D, Leung P, et al. (2016) Are you ready? Exploring readiness to engage in exercise among people living with HIV and multimorbidity in Toronto, Canada: a qualitative study. *BMJ Open* 6: e010029.
34. O'Brien KK, Tynan AM, Nixon SA, Glazier RH (2016) Effectiveness of aerobic exercise for adults living with HIV: systematic review and meta-analysis using the Cochrane Collaboration protocol. *BMC Infect Dis* 16: 182.
35. Aweto HA, Aiyegbusi AI, Ugonabo AJ, Adeyemo TA (2016) Effects of Aerobic Exercise on the Pulmonary Functions, Respiratory Symptoms and Psychological Status of People Living With HIV. *J Res Health Sci* 16: 17-21.
36. Chernoff RA, Martin DJ, Schrock DA, Huy MP (2010) Neuropsychological functioning as a predictor of employment activity in a longitudinal study of HIV-infected adults contemplating workforce reentry. *J Int Neuropsych Soc* 16: 38-48.
37. Dufour CA, Marquine MJ, Fazeli PL, Henry B, Ellis RJ, et al. (2013) Physical exercise is associated with less neurocognitive impairment in HIV-infected adults. *J Neurovirol* 19: 410-417.
38. Fazeli PL, Marquine MJ, Dufour C, Henry BL, Montoya J, et al. (2015) Physical Activity is Associated with Better Neurocognitive and Everyday Functioning Among Older Adults with HIV Disease. *AIDS Behav* 19: 1470-7.
39. Fazeli PL, Woods SP, Heaton RK, Umlauf A, Gouaux B, et al. (2014) An active lifestyle is associated with better neurocognitive functioning in adults living with HIV infection. *J Neurovirol* 20: 233-42.
40. Mattson MP (2013) Exercise and the brain: a slap on the HAND. *J Neurovirol* 19: 1-4.
41. Archer T (2011) Physical exercise alleviates debilities of normal aging and Alzheimer's disease. *Acta Neurol Scand* 123: 221-238.
42. Archer T (2012) Influence of physical exercise on traumatic brain injury deficits: scaffolding effect. *Neurotox Res* 21: 418-34.
43. Archer T (2014) Health benefits of physical exercise for children and adolescents. *J Novel Physiother* 4: 203-206.
44. Archer T, Garcia D (2014) Physical exercise improves academic performance and well-being in children and adolescents. *Int J School Cogn Psychol* 1: e102.
45. Archer T, Garcia D (2015) Exercise and dietary restriction for promotion of neurohealth benefits. *Health* 7: 136-152.
46. Archer T, Fredriksson A, Schütz E, Kostrzewa RM (2011) Influence of physical exercise on neuroimmunological functioning and health: aging and stress. *Neurotox Res* 20: 69-83.
47. DE Medeiros Guerra LM, Galvão DE Souza HA, Mesquita Soares TC, Gomes DA Silva J, DA Rocha Morgan DA, et al. (2016) Resisted exercise, morphological and functional standards, and quality of life of people living with HIV/AIDS. *J Sports Med Phys Fitness* 56: 470-5.
48. Mesquita Soares TC, Galvão De Souza HA, De Medeiros Guerra LM, Pinto E, Pipolo Milan E, et al. (2011) Morphology and biochemical markers of people living with HIV/AIDS undergoing a resistance exercise program: clinical series. *J Sports Med Phys Fitness* 51: 462-6.
49. Zanetti HR, Cruz LG, Lourenço CL, Ribeiro GC, Leite MA, et al. (2016) Nonlinear Resistance Training Enhances the Lipid Profile and Reduces Inflammation Marker in People Living With HIV: A Randomized Clinical Trial. *J Phys Act Health*.

Citation: Archer T (2016) Aspects of Cognition/Health Failure by HIV-infected Individuals: Amelioration through Exercise. HIV Curr Res 1: e101.

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission/