

# Association of Self-reported Height Loss and Kyphosis with Loss of Teeth in Japanese Elderly

Akira Taguchi<sup>1</sup>, Mikio Kamimura<sup>2</sup>, Noriyuki Sugino<sup>1</sup>, Keiichi Uchida<sup>1</sup>, Yutaka Kitamura<sup>3</sup>, Shota Ikegami<sup>4</sup>, Yukio Nakamura<sup>4</sup>, Shigeharu Uchiyama<sup>4</sup>, Hiroyuki Kato<sup>4</sup>

<sup>1</sup>Department of Oral and Maxillofacial Radiology, School of Dentistry, Matsumoto Dental University, Shiojiri 399-0781, Japan. <sup>2</sup>Center of Osteoporosis and Spinal Disorders, Kamimura Orthopaedic Clinic, Matsumoto 399-0021, Japan. <sup>3</sup>Center of Oral and Maxillofacial Surgery and Dental Implant in Shinshu, Nagano 382-0201, Japan. <sup>4</sup>Department of Orthopaedic Surgery, Shinshu University School of Medicine, Matsumoto 390-8621, Japan

## Abstract

**Study background:** Height loss and kyphosis are useful surrogate markers of osteoporotic vertebral fractures in the elderly. Loss of teeth in the elderly also is associated with osteoporosis. These imply the possibility that self-reported these indices may be associated with loss of teeth in the elderly. This study aimed to clarify the associations of self-reported height loss and kyphosis with number of teeth lost in Japanese elderly.

**Subjects and Methods:** Among patients who visited dispensing pharmacies in Matsumoto, Japan, 307 patients (75 men and 232 women) aged 50–97 years participated in the study. They completed a structured questionnaire including covariates related to loss of teeth. Self-reported height loss and kyphosis were simply defined as three categories: no; mild-to-moderate; severe.

**Results:** Analyses of covariance adjusted for covariates revealed that there were no significant differences in the numbers of teeth lost in total, or during the past 1 year among the three self-reported height loss categories. Significant differences were observed in the total numbers of teeth lost among the three self-reported kyphosis categories ( $p < 0.001$ ). Subjects who were conscious of severe kyphosis had significantly larger number of teeth lost (mean  $\pm$  SEM,  $16.1 \pm 1.8$ ) than those who were conscious of no kyphosis ( $8.7 \pm 0.6$ ,  $p < 0.001$ ) and mild-to-moderate kyphosis ( $8.3 \pm 0.7$ ,  $p < 0.001$ ). Furthermore, there were significant differences in the number of teeth lost during the past 1 year among the three self-reported kyphosis categories ( $p = 0.031$ ). Subjects who were conscious of severe kyphosis had significantly greater number of teeth lost during the past 1 year ( $0.9 \pm 0.2$ ) than those who were conscious of no kyphosis ( $0.3 \pm 0.1$ ,  $p = 0.03$ ).

**Conclusions:** Our results suggest that Japanese elderly with self-reported severe kyphosis may lost more teeth than those without self-reported severe kyphosis.

*Key Words: Height loss, Kyphosis, Tooth loss, Osteoporosis, Fracture*

## Introduction

Loss of teeth may be associated with changes in nutritional intake, progression of cognitive impairment, increased risk of disability, and increased mortality risk in elderly populations [1-4]. Dental caries and periodontal diseases are major causes of tooth loss in these populations. Retention of teeth is very important for maintenance of general health in the elderly. Although it remains unknown whether future tooth loss can be certainly predicted by evaluating dental caries and periodontal disease, smoking is significantly associated with greater progression of periodontal disease, resulting in tooth loss [5,6].

An association between tooth loss and osteoporosis determined by the percent cortical area at the metacarpal midshaft was firstly reported in postmenopausal women aged 60–69 years [7]. He concluded that middle-aged women may be more likely to retain their teeth if they avoid smoking and undertake an effective program to prevent the progression of osteoporosis. Since then, many investigators have demonstrated the association of periodontal disease progression and/or tooth loss with skeletal bone mineral density (BMD) measured by dual X-ray absorptiometry, especially in postmenopausal women [8-15]. Furthermore, a potential association of osteoporotic spine fractures with tooth loss has been reported, although the relationship between periodontitis and spine fractures remains unclear [16,17].

Two-thirds of patients with vertebral fractures have no symptoms like back pain; therefore, vertebral fractures are accurately determined by lateral radiographs. However, if such radiographs are used to evaluate the association between vertebral fractures and tooth loss, the radiation doses during the acquisition of lateral radiographs may be harmful for individuals without these fractures, making radiology hard to justify. Height loss and kyphosis from younger ages are considered to be surrogate markers or screening tools for vertebral fractures in the elderly [18,19]. Conversely, several thresholds for height loss to distinguish individuals with and without vertebral fractures have been suggested. However, it is likely that many elderly people may not accurately remember their heights at younger ages. Patients' estimated current height tended to be incorrect, with a mean difference of  $-2.5$  cm from the current measured height, in a total of 8,610 patients with a mean age of 70.9 years [20]. Regarding kyphosis, after adjustment for age, a  $15^\circ$  increase in kyphosis was associated with the presence of a vertebral fracture (OR, 1.57; 95% CI, 1.46–1.69) [21]. Kyphosis was more strongly related to thoracic fractures than to lumbar fractures, and kyphosis was most prominent in women with multiple thoracic wedge fractures in their study. However, untrained investigators are unlikely to be able to easily measure the degree of kyphosis.

If height loss and kyphosis are associated with prevalent





kyphosis categories ( $p=0.001$ ). Subjects who were conscious of severe kyphosis had significantly greater number of teeth lost during the past 1 year ( $0.9 \pm 0.2$ ) than those who were conscious of no kyphosis ( $0.3 \pm 0.1$ ,  $p=0.03$ ) (Figure 3).

### Discussion

Height loss and kyphosis are considered to be surrogate markers of vertebral fractures [18,19]. Our hypothesis was that these self-reported markers may be associated with number of teeth lost, because several investigators have reported associations between tooth loss and osteoporosis [8-17]. In fact, we found that self-reported kyphosis was significantly associated with number of teeth lost in our study, while self-reported height loss was not. It was reported that patients' estimated current height tended to be incorrect [20]. In addition, height loss can

be caused not only by vertebral fractures, but also to certain extents by intervertebral disk degeneration that decreases disk height, osteoarthritic conditions of the spine, hip, or knee, various inflammatory and structural/congenital spinal deformities, and weakness of the back muscles. No significant relationship was noted between vertebral fractures and height loss in 400 Japanese men and women [23]. Measurement of kyphosis may be similar to that of height loss. However, 10% of women with the most severe kyphosis had 7–17% lower BMD compared with the rest of their cohort [24]. Although we found no difference in numbers of teeth lost between subjects with self-reported no kyphosis and mild-to-moderate kyphosis, individuals with self-reported severe kyphosis may have lost more teeth in total than those with self-reported no or mild-to-moderate kyphosis.

Subjects who were conscious of severe kyphosis had significantly greater number of teeth lost (both total numbers of teeth lost and number of teeth lost during the past 1 year) than those who were conscious of no kyphosis. This implies the possibility that individuals who are conscious of severe kyphosis may lose a greater number of teeth than those without self-reported kyphosis in the future. Women who lost teeth during a 7-year follow-up period experienced less favorable changes in BMD at all sites compared with 144 women who lost no teeth among 189 healthy, white, dentate, postmenopausal women [25]. A significant relationship also was found between changes in skeletal BMD and number of teeth lost during a 5-year study period in Japanese postmenopausal women [15].

The numbers of teeth lost in total and during the past 1 year were self-reported by the participants in this study. It is likely that self-reported numbers of teeth lost may be inaccurate measures; however, many investigators have used self-reported tooth counts in large epidemiological studies to assess the associations between tooth loss and general diseases like arterial diseases and cancer [26-28]. Self-reported numbers of remaining teeth, fillings, root canal therapies, and prostheses were strongly correlated with clinical records ( $r=0.74-1.0$ ) [29]. In the present study, the correlation coefficient between the actual and self-reported numbers of remaining teeth was 0.82.

This questionnaire-based survey had some limitations. First, all study subjects were patients who visited dispensing pharmacies in Matsumoto, rather than healthy subjects. In addition, some orthopedic clinics were near the dispensing pharmacies in this study, resulting in including many subjects (49.8%) who had been treated with osteoporosis medications. Therefore, our subjects are not representative of normal Japanese elderly. Our sample size of subjects also was relatively small. Further investigations including large numbers of community-dwelling individuals or those who attend medical check-ups will be necessary to clarify our findings.

Second, the self-reported periodontal status in this study was only at the time of visit to the dispensing pharmacy. The validity of a simple question asking whether subjects have periodontal symptoms is expected to be relatively good according to our previous study [22]; however, it is unknown whether self-reported periodontal status at the time of visit to a dispensing pharmacy is representative of previous or future




Figure 1. Total numbers of teeth lost among the three self-reported height loss categories. Data are shown as means  $\pm$  SEM.




Figure 2. Total numbers of teeth lost among the three self-reported kyphosis categories. Data are shown as means  $\pm$  SEM.




Figure 3. Numbers of teeth lost during the past 1 year among the three self-reported kyphosis categories. Data are shown as means  $\pm$  SEM.



