

Atypical HUS with Dysphagia and Palatal Palsy Responding to IVIG

Vivek Kumar^{1*}, Kiran Sodhi², Madhuri Kanitkar³, Deepak⁴, Anupama⁴, Ajay Bajpai⁴ and Shonenedra⁴¹Department of Pediatrics, Command Hospital, Lucknow, India²Department of Pediatrics, Air Force Hospital, Kanpur, India³Department of Pediatrics and Pediatric Nephrology Base Hospital, Delhi, India⁴Division of Pediatrics, Pediatric Residents, Command Hospital, Lucknow, India

Abstract

D negative/Atypical HUS (Hemolytic Uremic Syndrome) is a difficult condition to diagnose and treat. It has been shown to be a disease of complement dysregulation. Treatment is difficult and revolves around plasma exchange. We report one case of an Atypical HUS in a toddler girl who developed multi organ dysfunction associated with never reported dysphagia and palatal paralysis which responded to IVIG (Intravenous Immunoglobulin) given over five days

Keywords: Atypical HUS; Dysphagia; Palatal palsy; IVIG

Introduction

The classical triad of microangiopathic hemolytic anemia, renal failure and thrombocytopenia seen in Typical HUS may not present acutely and simultaneously in an atypical form. Atypical HUS may involve multi organ system. CNS manifestations comprising of irritability, encephalopathy and thrombotic stroke is seen in 20-50% of the cases [1]. We present one case of an Atypical HUS in a toddler girl with multi organ dysfunction and never reported manifestation of dysphagia and palatal paralysis. Atypical HUS main line of treatment is plasma exchange. Due to non availability of plasma exchange therapy our patient was managed with FFP (Fresh Frozen Plasma) and supportive treatment while her dysphagia and palatal paralysis responded to IVIG.

Case Report

2 yr old female toddler was admitted with a history of (a) high grade intermittent fever without chills and rigors 15 days duration, (b) swelling over whole body starting over face and periorbital area 7 days duration (c) decreased urine output 1 day duration. On admission patient weight was 10.5 kg (85% of 50 centile) and height was 85 cm (92% of 50 centile NCHS standard). She was irritable, febrile-100°F, Pulse 120/minute regular normal volume, Blood Pressure 120/98 mm Hg (more than 99 centile), Respiratory rate 30/minute. Pallor and edema (periorbital and limb) was present. Systemic examination revealed firm hepatomegaly with ascites. Patient was started on restricted iv fluids (insensible loss and urine output), intravenous antibiotics (cefoperazone @ 150 mg/kg/day in three divided doses), diuretics, 5% albumin (single dose @ 1 gm/kg) and antihypertensives (nifedipine @ 1.5 mg/kg/day, atenolol 0.5 mg/kg/day and finally prazosin @ 0.1 mg/kg/day) and furosemide @ 2 mg/kg/day. Initial investigations revealed hb 7.9 gm%, Total counts 10200/cmm Differential count polymorphs 60% Lymphocytes 35% eosinophils 4% monocytes 5% platelets 3.1 lac/cumm, urine routine/microscopic 7-8 pus cells/hpf RBC 15-17/HPF protein 2+ blood urea/serum creatinine 196/3.2 mg%, serum sodium /potassium 136/4.4 meq/dl, serum albumin 2.8 gm%, serum bilirubin 0.8 mg% SGOT/SGPT 192/103 IU/L, serum cholesterol 170 mg%, C3, C4 was normal, serum amylase was normal, ANA, dsDNA, ANCA, DCT, HbsAg, HIV and anti HCV were negative. USG abdomen showed bilateral medical renal disease and ascites while Chest X-ray was normal. At this stage of the illness possibilities were AKI (Acute Kidney Injury) secondary to (a). RPGN (Rapidly Progressive Glomerulonephritis) (b). D negative HUS (Diarrhea negative Hemolytic Uremic Syndrome) (c). Interstitial Nephritis.

She improved in next three days so we started her on oral feeds.

On day four of admission she developed hyperglycemia (486 mg%) with ketonuria 4+. For this she was managed with intravenous fluids, insulin and potassium. This transient DKA (Diabetic Ketoacidosis) state recovered in next 48 hours. Next three days her edema and renal function improved. But on day 10 of admission she developed tachycardia with increased pallor. Her hemoglobin had dropped to 5.4 gm% while PBS (Peripheral Blood Smear) showed features of intravascular hemolysis with thrombocytopenia (Table 1). Her LDH (Lactate Dehydrogenase) was 2000IU/L. She was transfused PRBC (Packed Red Blood Cell) @ 25 ml/kg over next three days. She was also given Fresh frozen plasma @ 10 ml/kg for the next three days, due to non availability of plasma exchange, guided by serum LDH level. Her condition stabilised and urine output improved. On Day 14 of admission she had hypokalemic (serum potassium 1.9 meq/dl) flaccid quadriplegia with bulbar paralysis which was managed with iV potassium followed by oral potassium supplements (Figure 1). Her paralysis recovered in limbs with normalisation of serum potassium but dysphagia, nasopharyngeal regurgitation and palatal palsy persisted. MRI brain was normal. CH50, factor H and I were normal. Serum and urine amino acid chromatography for cobalamin metabolism was normal. She was given IVIG @400 mg/kg for five day, with this she started showing improvement in next three days and her palsy completely recovered in ten days (Table 2). During this period she was kept on tube feeds her kidney function and liver enzymes normalised but her hypertension persisted. On follow up at one month there was no edema and blood pressure was between 90-95 centile with three antihypertensives. Renal function was normal. Pediatric nephrologist consultation was taken and it was advised not to do a renal biopsy at this stage. ADAMTS13 and complement gene mutations studies could not be done due to non availability.

Discussion

Central nervous system manifestations in the form of irritability,

***Corresponding author:** Vivek kumar, 7 Air Force Hospital, Nathu Singh Road, Kanpur Cantt-208004, Uttar Pradesh, India, Tel: +919532295066; E-mail: Vk3532@gmail.com

Received July 12, 2013; **Accepted** November 04, 2013; **Published** November 06, 2013

Citation: Kumar V, Sodhi K, Kanitkar M, Deepak, Anupama, et al. (2013) Atypical HUS with Dysphagia and Palatal Palsy Responding to IVIG. *Pediat Therapeut* 3: 176. doi:[10.4172/2161-0665.1000176](http://dx.doi.org/10.4172/2161-0665.1000176)

Copyright: © 2013 Kumar V, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

| | Day 16 of illness | Day 20 | Day 26 | Day 29 |
|-------------------------------------|---|------------------|--------------------------------------|-----------------|
| Hb gm% | 7.9, no evidence of hemolysis | 9.1 | 5.4, intravascular hemolysis present | 8.2 |
| TLC /cmm | 10,200 | 16,300 | 6,800 | 7000 |
| Platelets | 3.1 lac/cu mm | 2 lac/cu mm | 40,000/cu mm | 1 lac/cu mm |
| Blood Urea mg% | 196 | 106 | 50 | 30 |
| Serum creat mg% | 3.2 | 0.9 | 0.8 | 0.5 |
| Serum sodium meq/dl | 136 | 134 | 130 | 124 |
| Serum potassium meq/dl | 4.4 | 4.9 | 3.9 | 1.9 |
| Serum. Albumin gm% | 2.8 | 3.2 | 3.5 | 3.4 |
| Serum. Bilirubin mg% | 0.8 | 0.9 | 0.8 | 0.7 |
| SGOT/SGPT IU/L | 192/103 | 203/86 | 67/45 | 45/40 |
| Urine RE/ME | Pus cells 7-8/HPF 15RBC/HPF Urine Protein 2+ | Urine protein 1+ | NAD | Urine protein1+ |
| RBC-15-17/HPF, Protein 2+ | Urine protein 1+ | NAD | Urine protein1+ | |
| Lactate Dehydrogenase IU/L | | | | 800 |
| Direct coombs test (DCT) | | | Negative | |
| PBS(Peripheral Blood Smear) present | Normocytic Normochromic | | Schistocytes, burr cells | |
| Blood culture | | Sterile | | |

Table 1: Showing details of investigations: features of intravascular hemolysis with thrombocytopenia.




Figure 1: Bar chart depicting clinical Problems with duration.

| | | |
|----|------------------------------------|---|
| 1. | Initial Presentation | Restricted fluids, multiple antihypertensives, diuretics and supportive treatment |
| 2. | DKA | Insulin, intravenous fluids and potassium |
| 3 | Intravascular Hemolysis | PRBC and FFP |
| 4 | Hypokalemia/quadriparesis | IV Potassium followed by oral potassium |
| 5 | Dysphagia and palatal palsy | IVIG |

Table 2: Depicting various interventions: Improvement and recovery of Palsy.

drowsiness, seizures, cortical blindness, hemiparesis and coma are known extra renal complications of Atypical or D negative HUS [2]. These manifestations could be due to microangiopathic lesions or RPLS (Reversible Posterior Leukoencephalopathy Syndrome) secondary to hypertension. Both could be documented in MRI Brain. Complications of dysphagia and palatal palsy with a normal MRI brain are reported for the first time to the best of our knowledge. These complications getting cured using IVIG itself is a novel finding.

A typical HUS has shown to be due to alternative complement pathway dysregulation with 50% cases involving complement regulatory genes, factor H, factor I and MCP (Membrane Cofactor Protein) [3].

Citation: Kumar V, Sodhi K, Kanitkar M, Deepak, Anupama, et al. (2013) Atypical HUS with Dysphagia and Palatal Palsy Responding to IVIG. *Pediat Therapeut* 3: 176. doi:10.4172/2161-0665.1000176

Plasma exchange therapy though cumbersome and invasive still forms the main modality of treatment. However Watt et al. had reported the only case of Atypical HUS responding to a combination of steroids and IVIG [4]. Our case who had developed a never reported finding of dysphagia and palatal palsy following Hypokalemia responded dramatically to five day therapy of IVIG (total 2 gm) alone. This could be a good alternative to plasma therapy apart from very promising and upcoming Eculizumab (monoclonal antibody against complement factor) [5].

References

- Eriksson KJ, Boyd SG, Tasker RC (2001) Acute neurology and neurophysiology of haemolytic-uraemic syndrome. *Arch Dis Child* 84: 434-435.
- Koehl B, Boyer O, Biebuyck-Gougé N, Kossorotoff M, Frémeaux-Bacchi V, et al. (2010) Neurological involvement in a child with atypical haemolytic uremic syndrome. *Pediatr Nephrol* 25: 2539-2542.
- Kavanagh D, Goodship TH, Richards A (2006) Atypical haemolytic uraemic syndrome. *Br Med Bull* 77-78: 5-22.
- Watt T, Warshaw B, Katzenstein HM (2009) A typical hemolytic uremic syndrome responsive to steroids and intravenous immune globulin. *Pediatr Blood Cancer* 53: 90-91.
- Loirat C, Babu S, Furman R, Sheerin N, Cohen D, et al. (2011) Eculizumab Efficacy and Safety in Patients With Atypical Hemolytic Uremic Syndrome (aHUS) Resistant to Plasma Exchange/Infusion, London, UK.

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

User friendly/feasible website-translation of your paper to 50 world's leading languages
Audio Version of published paper
Digital articles to share and explore

Special features:

300 Open Access Journals
25,000 editorial team
21 days rapid review process
Quality and quick editorial, review and publication processing
Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, and Google Scholar etc
Sharing Option: Social Networking Enabled
Authors, Reviewers and Editors rewarded with online Scientific Credits
Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>

