

Audit of Screening for Diabetic Nephropathy in a Teaching Hospital in Nigeria

Okafor UH^{1*}, Ezeala A^{1,2} and Aneke E¹

¹Department of Medicine, Enugu State University Teaching Hospital (ESUTH) Parklane Enugu, Nigeria

²Department of Medicine, University of Nigeria Teaching Hospital Enugu, Nigeria

Abstract

Background: Diabetes nephropathy is the commonest cause of end stage kidney disease worldwide. The high incidence of the disease highlights the need to detect it early and commence prompt intervention. The aim of this study is to audit the screening for diabetic nephropathy in patients with diabetes mellitus attending a teaching hospital in Nigeria.

Methodology: This a retrospective study and the study population was adult patients with diabetes mellitus who attended medical outpatient clinic (MOP) of ESUTH from 1st January 2010 to 31st December 2011. Their medical records were retrieved from the hospital data bank and their biodata, clinical and laboratory parameters were documented. Data was analysed using SPSS Vs 15.

Results: Two hundred and three patients were selected for the study, 32% were males, mean age of 60.0 ± 13.5 years and range of 27 to 94 years. The duration of DM ranges from 2 to 312 months, 4.1% had type 1 DM, 88.1% were hypertensive. The eGFR ranges from 11 to 205 ml/min/1.73 m², 37.6% of the patients had eGFR less than 60 ml/min/1.73 m² and 22.3% had proteinuria. Duration of diabetes mellitus and duration of hypertension were assessed in 91 and 25% of the patients respectively; weight, oedema and sensation were assessed in less than 7%. BMI and fundoscopy was not assessed in any of the patients. 92.6% and 97% of the patients had records of their blood sugar on the first and last follow up. Less than 50% of the patients had record of serum electrolyte, urea, creatinine, urinalysis and packed cell volume. None of the patients had records of HbA1C or eGFR.

Conclusion: The patients were poorly screened for diabetic nephropathy.

Keywords: Diabetic nephropathy; Diabetes mellitus; Hypertension; Proteinuria

Introduction

Diabetes mellitus (DM) is a serious and growing public health problem affecting 2% to 3% of the adult population, and up to 20% to 25% of the elderly population [1]. It results in reduced life expectancy and increased morbidity resulting from diabetes-related complications including diabetic nephropathy.

Diabetic nephropathy is a micro vascular and common chronic complication of diabetes mellitus. It is the commonest cause of end stage kidney disease worldwide [1]. However, in spite of being a common co morbid condition of type 2 diabetes mellitus, it is often remains undetected and consequently untreated [2]. The high incidence of the disease highlights the importance of detecting it early so that treatment to delay its progression can begin.

Therapeutic interventions and efforts to prevent the progression of nephropathy have the greatest impact if instituted very early in the course of the disease. Therefore, it is recommended that all patients with DM should receive routine screening tests for diabetic nephropathy so that appropriate treatments can be instituted as early as possible [2]. Despite this evidence, implementation rates of recommended interventions are low. This frequently leads to ineffective or delayed treatment of complications [2,3].

Screening for diabetic nephropathy must be initiated at the time of presentation, since many of the adult diabetic patients especially patients with type 2 DM already have evidence of kidney disease at that time [4,5].

The aim of this study is to audit the screening for diabetic

nephropathy in patients with diabetes mellitus attending a teaching hospital in Nigeria.

Methodolgy

This was a retrospective study. The study location was medical outpatient clinic of Enugu State University Teaching Hospital (ESUTH) Parklane Enugu in Nigeria. The hospital was initially a multispecialty specialist hospital and was changed to a teaching hospital in 2005. It is situated in Enugu metropolis in Enugu State in Nigeria. The hospital caters for patients from Enugu State and also receives referral from the neighboring states of Anambra, Imo, Abia, Ebonyi, Benue, Delta, Cross Rivers and Kogi. It is a 400 bedded hospital offering emergency and outpatient services in all specialties. It also offers training program for students in undergraduate and post graduate medicine.

The study population was adult patients with diabetes mellitus who attended medical outpatient clinic (MOP) of ESUTH from 1st January 2010 to 31st December 2011. Patients that were with gestational diabetes

***Corresponding author:** Okafor UH, Department of Medicine, Enugu State University Teaching Hospital (ESUTH) Parklane Enugu, Nigeria, Tel: +2348033130014; E-mail: umehughes@yahoo.com

Received January 19, 2015; **Accepted** March 16, 2015; **Published** March 19, 2015

Citation: Okafor UH, Ezeala A, Aneke E (2015) Audit of Screening for Diabetic Nephropathy in a Teaching Hospital in Nigeria. J Diabetes Metab 6: 525. doi:10.4172/2155-6156.1000525

Copyright: © 2015 Okafor UH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

mellitus and newly diagnosed (less than a month or 2 follow up visit) of diabetes mellitus were excluded from the study.

The subjects were randomly selected from the DM patients who attended MOP during the study period. Their medical records were retrieved from the hospital data bank. The biodata, clinical and laboratory parameters of the subjects as in their medical records were documented. The onset and duration diabetes mellitus in these patients were determined from onset of clinical features/or elevation of the blood glucose till time of presentation.

The estimated glomerular filtration rate was calculated using Cockcroft and Gault formula.

The data obtained was entered into a spread sheet and analysed using SPSS Vs 15. Continuous variables were reported as mean \pm standard deviation and categorical variables as proportions. P value less than 0.05 was significant.

Results

Patient's characteristics

The records of 203 patients that were selected among the DM patients that attended MOP during the study period were retrieved. Using the available records, the characteristics of the patients were as follows. Sixty five (32%) of the subjects were males, with male: female ratio of 1:2.1. Their mean age was 60.0 ± 13.5 years with a range of 27 to 94 years. The duration of DM ranges from 2 to 312 months with a median of 60 months, 4.1% had type 1 DM and they were on insulin, 85.9% had type 2 DM, non-had type 3 DM and patients with type 4 (gestational) DM were excluded.

One hundred and seventy nine (88.1%) patients were hypertensive and the mean systolic blood pressure was 139 ± 29 mmHg, and diastolic blood pressure was 84 ± 16 mmHg. 62.1% of the patients with hypertension were on angiotensin converting enzyme inhibitor. The estimated GFR (eGFR) ranges from 11 to 205 ml/min/1.73 m² with a median of 79.1 ml/min/1.73 m², 37.6% of the patients had eGFR less than 60 ml/min/1.73 m² and 22.3% had proteinuria.

Audit of clinical parameters

All the patients had records of their age, sex and blood pressure. Duration of diabetes mellitus and hypertension were assessed in 91 and 25% of the patients respectively. Less than 7% of the patients had their weight, oedema and sensation assessed. None of the patients studied was assessed for body mass index (BMI) or fundoscopy. The details of the clinical parameters assessed were shown in Table 1.

Audit laboratory parameters

One hundred and eighty nine (92.6%) and 197 (97%) patients had records of their blood sugar on the first and last clinic visit respectively. Less than 50% of the patients had record of serum electrolyte, urea and creatinine, urinalysis and packed cell volume. None of the patients had records of glycosylated haemoglobin (HbA1C), microalbuminuria and glomerular filtration rate (eGFR). The details are shown in Table 2.

Discussion

The 21st century has the most diabetogenic environment in human history with increasing incidence and prevalence globally [5,6]. This increase in prevalence of diabetes mellitus had been estimated to be greater in the developing countries including Nigeria. Diabetes mellitus especially type 2 had been reported to be more preponderant in females and middle age [1]. This study revealed that 68% of the

studied population was females and their mean age was 60 ± 13.5 years. The National Kidney Foundation Kidney Disease Outcomes Quality Initiative (KDOQI) clinical practice guidelines recommend screening for CKD in at risk individuals using proteinuria and the eGFR [7]. A 2009 study from the United States found that almost 40% of type 2 DM patients have CKD to some degree [8] it is found in this study that among patients with available records, more than a third of them have evidence of kidney dysfunction as detected by eGFR less than 60 ml/min/1.73 m² and/or proteinuria.

A detailed physical assessment of patients with diabetes mellitus is very important and will encourage early detection and prompt management of antecedent complications including diabetic nephropathy. This will forestall poor morbidity and lead to better outcome. Thus comprehensive history and physical examination including duration of diabetes mellitus, hypertension, smoking, obesity and assessment of other complications of diabetes mellitus including diabetic neuropathy and especially diabetic retinopathy at presentation are important. These parameters may give clue to early indicators to the development of diabetic nephropathy. Diabetic nephropathy is a microvascular complication and has been noted to be associated with other microvascular injuries like diabetic retinopathy and neuropathy. Thus early detection of these markers will assist in prevention, early diagnosis and subsequently prompt intervention to slow progression of diabetic nephropathy and other complications of diabetes mellitus.

Clinical Parameter	No of patients with records (203)	Percent of patients with records (100)
Age	203	100
Sex	203	100
Occupation	136	67.0
Weight	14	6.9
BMI	0	0
Duration of DM	185	91.1
HX of HBP	179	88.1
Duration of HBP(179)	44	24.6
Urinary symptoms	125	61.6
Oedema	14	6.9
BP	100	100
Sensation	12	5.9
Fundoscopy	0	0

BMI – body mass index; DM – diabetes mellitus; HX – history; HBP – hypertension; BP – blood pressure

Table 1: Audit of clinical parameters.

Laboratory Parameters	No of patients with records (203)	Percent of patients with records (100)
Fbs at 1st	188	92.6
Fbs at last	197	97.0
HbA1c	0	0
Electrolyte	67	33.0
Creatinine	85	41.9
eGFR	0	0
Lipid profile	43	21.2
Uric acid	1	0.5
PCV	88	43.4
Urinalysis	93	45.8
Microalbuminuria	0	0

Fbs – fasting blood sugar; HbA1c – glycosylated haemoglobin; eGFR – estimated glomerular filtration rate; PCV – packed cell volume

Table 2: Audit of laboratory parameters.

It is noteworthy that in this study none of the patients had record of assessment of fundoscopy or BMI, less than 10% had record that their sensory modalities or weight was assessed, and also records of assessment of features of diabetic nephropathy like oedema or urinary symptoms was found in less than 60%. Furthermore about 10% of the patients do not have records of duration of diabetes mellitus or history of hypertension. Thus the assessment of these parameters in these patients was very poor. Similar inadequate assessment of diabetes mellitus patients had been reported in studies in other part of the country [9,10] and this eclipse the benefit and expected impact on early and prompt detection and possible intervention of diabetic nephropathy and other complications of diabetes mellitus including associated cardiovascular risk factors. However it is of note that the age, sex and blood pressure was assessed in all the patients in this study. Even though this is commendable, it does not compensate for the gross inadequacies of the assessment of the clinical parameters of these patients. More so these aforementioned parameters are usually assessed by the support staff and not the attending physicians.

Diabetic nephropathy is usually defined as a clinical syndrome classically characterized by proteinuria and/or deteriorating glomerular filtration rate. It has 5 stages of hyperfiltration, normoalbuminuria, microalbuminuria, overt nephropathy and end stage kidney disease depending on the level of the proteinuria or glomerular filtration rate [7,11]. This shows that diagnosis and staging of diabetic nephropathy depends on the level of proteinuria and glomerular filtration rate of the patient. Proteinuria is usually assessed using timed or spot urine while GFR is assessed with measured or estimated technique. The use of spot urine (dip stick or urine protein:creatinine ratio) and estimated/calculated (Cockcroft and Gault or MDRD) method were preferred for proteinuria and GFR respectively because of its convenience and reliability [4,12]. Microalbuminuria (urine protein 30-299 mg/day), overt proteinuria (urine protein \geq 300 mg/day) and GFR less than 60 ml/dl is indicative of kidney disease. Only 41.9% and 45.8% of patients that participated in this study had record of urinalysis and serum creatinine respectively, and none of them had record of microalbuminuria or GFR [7,13]. Among the patients with record of urinalysis or serum creatinine 22.3% and 37.6% of them had overt proteinuria and eGFR less than 60 ml/min/1.73 m² respectively. This shows that though the screening of diabetic nephropathy using both parameters is poor in this study, the prevalence of diabetic nephropathy is higher than reported from a similar study in Nigeria [10]. This result of degree of assessment/screening for diabetic nephropathy is poor and falls short of the recommendation, which states that all adult patients with diabetes mellitus should undergo assessment of their kidney status using proteinuria and GFR at diagnoses and periodically depending on the state of the patient and the kidney [4,5].

Hyperlipidaemia, anemia, hyperglycaemia, hypertension, cigarette smoking and proteinuria has been reported as factors that propagate kidney diseases including diabetic nephropathy and other systemic complications of diabetes mellitus [4,5,14,15]. The paucity of adequate screening of these propagating factors of diabetic nephropathy was also noted in this study. Lipid profile and Packed Cell Volume (PCV) was assessed in 21.1% and 43.2% respectively. None of the patients was assessed for smoking, glycated haemoglobin and microalbuminuria in all their follow up at the medical outpatient clinic.

The dismal screening of diabetic nephropathy is not restricted to Nigeria as studies from Bosnia and Herzegovina [16], Malaysia [17], South African [18] and Sri Lanka [19] had recorded poor assessment of risk factors for diabetic nephropathy in patients with diabetes mellitus.

Reports from developed countries rather showed that over 70% of diabetes mellitus patients were screened for risk factors of diabetic nephropathy in the course of their management [20,21].

The suboptimal assessment of diabetes mellitus patient is not restricted to screening for diabetic nephropathy as there has been reported gross under assessment of patients with diabetes mellitus in Nigeria. Furthermore screening of kidney diseases in at risk groups have been reported to be poor [22]. There are multiplicities of factors contributing to this malady in the developing countries including Nigeria. These factors could be poverty and ignorance in the part of patient, ignorance or laxity in the part of the medical personnel, or non-availability of the laboratory facility, lack of protocols and guidelines for management (detection, prevention and treatment) of diabetic nephropathy and other complications of diabetes mellitus in the health institution. The study location has good and standard laboratory facilities to screen these patients appropriately, thus the challenges relies either with the patient or the medical personnel who in this study is a physician or resident in Medicine.

Medical practice especially renal medicine in the developing countries including Nigeria has been plagued with many daunting challenges. This has been perpetuated by many policies that do not encourage improvement in output of medical practice. The cost of medical practice is not cheap and thus in many developed countries there are policies that alleviate medical treatment for the populace. However in developing countries including Nigeria patients and medical personnel bear the entire cost of medical care in presence of other competing needs. Thus there are minimal or no awareness campaign to educate the patient and the populace at large on prevention, monitoring and screening of diseases like diabetic nephropathy. The patients are so poor that they find it very difficult to procure medication and other necessary investigations, thus adherence to medications or regular follow up is not feasible. The medical personnel finds it difficult to cope with very busy clinic, can't afford appropriate update and retraining programs, and occasionally their diligence and dedication is not forthcoming because of lack of motivation. There is no laid down guideline, audit, supervision and monitoring in most institution including the study location.

The result of this study surmises the poor state of our medical care and the sorrowful antecedent outcome of these diabetic patients considering that diabetic nephropathy is expensive to manage and has high mortality. Thus there is need for urgent attention to the health sector in terms of health education, regular clinical audit, update programs for health personnel, reliable medical insurance system, upgrade of the health facilities and population friendly policies that will encourage prevention of diseases and its complication.

References

1. US Renal Data System (2003) USRDS 2003 Annual Data Report: Atlas of End-Stage Renal Disease in the United States. National Institute of Diabetes and Digestive and Kidney Diseases, National Institute of Health, Bethesda, MD.
2. Brechner RJ, Cowie CC, Howie LJ, Herman WH, Will JC, et al. (1993) Ophthalmic examination among adults with diagnosed diabetes mellitus. *JAMA* 270: 1714-1718.
3. Vijan S, Stevens DL, Herman WH, Funnell MM, Standiford CJ (1997) Screening, prevention, counseling, and treatment for the complications of type II diabetes mellitus. Putting evidence into practice. *J Gen Intern Med* 12: 567-580.
4. Molitch ME, DeFronzo RA, Franz MJ, Keane WF, Mogensen CE, et al. (2004) Nephropathy in diabetes. *Diabetes Care* 27 Suppl 1: S79-83.
5. Adler AI, Stevens RJ, Manley SE, Bilous RW, Cull CA, et al. (2003)

- Development and progression of nephropathy in type 2 diabetes: the United Kingdom Prospective Diabetes Study (UKPDS 64). *Kidney Int* 63: 225-232.
6. Retnakaran R, Cull CA, Thorne KI, Adler AI, Holman RR; UKPDS Study Group (2006) Risk factors for renal dysfunction in type 2 diabetes: U.K. Prospective Diabetes Study 74. *Diabetes* 55: 1832-1839.
 7. National Kidney Foundation (2002) K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 39: S1-266.
 8. Koro CE, Lee BH, Bowlin SJ (2009) Antidiabetic medication use and prevalence of chronic kidney disease among patients with type 2 diabetes mellitus in the United States. *Clin Ther* 31: 2608-2617.
 9. Okoro EO, Adejumo AO, Oyejola BA (2002) Diabetic care in Nigeria: report of a self-audit. *J Diabetes Complications* 16: 159-164.
 10. Agaba EI, Puepet FH, Ugoya SO, Agaba PA, Adabe R, et al. (2009) Chronic kidney disease screening and renoprotection in type 2 diabetes. *Ann Afr Med* 8: 52-54.
 11. Bakris GL (2011) Recognition, pathogenesis, and treatment of different stages of nephropathy in patients with type 2 diabetes mellitus. *Mayo Clin Proc* 86: 444-456.
 12. Levey AS, Coresh J, Balk E, Kausz AT, Levin A, et al. (2003) National Kidney Foundation practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Ann Intern Med* 139: 137-147.
 13. Caramori ML, Fioretto P, Mauer M (2003) Low glomerular filtration rate in normoalbuminuric type 1 diabetic patients: an indicator of more advanced glomerular lesions. *Diabetes* 52: 1036-1040.
 14. Gall MA, Hougaard P, Borch-Johnsen K, Parving HH (1997) Risk factors for development of incipient and overt diabetic nephropathy in patients with non-insulin dependent diabetes mellitus: prospective, observational study. *BMJ* 314: 783-788.
 15. Ravid M, Brosh D, Ravid-Safran D, Levy Z, Rachmani R (1998) Main risk factors for nephropathy in type 2 diabetes mellitus are plasma cholesterol levels, mean blood pressure, and hyperglycemia. *Arch Intern Med* 158: 998-1004.
 16. Novo A, Jokić I (2008) Medical audit of diabetes mellitus in primary care setting in Bosnia and Herzegovina. *Croat Med J* 49: 757-762.
 17. Chew BH, Palikat JM, Nur Syamimi A, Nor Azillah A, See JK, et al. (2012) An Audit of Type 2 Diabetes Care in a Malaysian Public Community Polyclinic. *Pertanika J Sci & Technol* 20: 97-102.
 18. Levitt NS, Bradshaw D, Zwarenstein MF, Bawa AA, Maphumolo S (1997) Audit of public sector primary diabetes care in Cape Town, South Africa: high prevalence of complications, uncontrolled hyperglycaemia, and hypertension. *Diabet Med* 14: 1073-1077.
 19. Ranasinghe CC, Rodrigo PC, Premaeatne S, Adikari AMMC, Rajapakse S (2010) An audit on the care of type diabetes mellitus in a tertiary care medical clinic in Sri Lanka. *Int J Diab Dev Ctries* 30: 3.
 20. Färnkvist LM, Lundman BM (2003) Outcomes of diabetes care: a population-based study. *Int J Qual Health Care* 15: 301-307.
 21. Khunti K, Ganguli S, Baker R, Lowy A (2001) Features of primary care associated with variations in process and outcome of care of people with diabetes. *Br J Gen Pract* 51: 356-360.
 22. de Lusignan S (2006) Chronic kidney disease. *Br J Gen Pract* 56: 885.