

## Bacterial Outer Membrane Proteins – Dependent Complement Activation

Bożena Futoma-Koloch\*

Department of Microbiology, Institute of Genetics and Microbiology, University of Wrocław, Wrocław, Poland

\*Corresponding author: Bożena Futoma-Koloch, Department of Microbiology, Institute of Genetics and Microbiology, University of Wrocław, Przybyszewskiego 63-77, 51-148 Wrocław, Poland, Tel: 55298589295; E-mail: [bozena.futoma-koloch@uwr.edu.pl](mailto:bozena.futoma-koloch@uwr.edu.pl)

Received date: Oct 30, 2015; Accepted date: Nov 02, 2015; Published date: Nov 07, 2015

Copyright: ©2015 Futoma-Koloch B. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Editorial

Complement (C) plays a pivotal role in the innate defence against microorganisms that invade animal and human bodies. Its activation through the classical (CP), lectin (LP) and alternative (AP) pathways lead to opsonisation and lysis of pathogens by incorporation of the membrane attack complex (MAC) into the membrane surrounding the microorganism's body. Bacteria getting into contact with human blood or plasma have developed a variety of strategies to evade C attack. Of special interest are outer membrane proteins (OMP) located in the outer membrane (OM) of gram-negative bacteria, which were shown to be involved in interactions with complement's factors [1]. Some OMP bind host's C regulatory proteins, while some of the others directly inactivate C components. Bacterial OMP may provide resistance against bactericidal action of the C system; however some of them may increase sensitivity of bacteria to serum [2]. Finding factors that control C components deposition on bacterial cells is the way to explain the sophisticated mechanism of bacteraemia leading to sepsis. Additionally, the mechanisms of OMP-mediated activation are currently intensively investigated, because of its high diversity and principal antigen vaccine potential. OMP are surface factors that are often utilized by gram-negative bacteria to avoid C-mediated recognition and destruction. Firstly, many OMP can bind specific fluid-phase regulators, such as factor H or C4bp protein. Respiratory pathogen *Moraxella catarrhalis* directly interacts with C<sub>3</sub>, as it produces C<sub>3</sub>-binding protein UspA [3]. The analysis of C<sub>3</sub> activation on *Salmonella* isolates belonging to the O48 serogroup showed that OMP in the range of molecular weights of 35-48 kDa determined their sensitivity to normal human serum (NHS) by C<sub>3</sub> activation [4]. It on the bacteria does not guarantee is worth mention that activation of the C system by surface structures the elimination of the microorganisms' cells from the host fluids. For example, no association between C<sub>3b</sub> deposition on the *N. meningitidis* cells and their lysis was found [5]. What is more, there are numerous OMP, which cleave complement's components or interfere with a correct formation of MAC, the final step in C activation. Well documented is the research on human enteropathogen *Yersinia enterocolitica* strain lacking OmpR regulator protein. OmpR exhibited extremely high resistance to the bactericidal activity of NHS through Ail (factor involved in the serum-resistant (SR) phenotype) expression compared with the wild-type strain [6]. Within OMP isolated from *Salmonella* O<sub>48</sub> serogroup, surface proteins characteristic of the bacterial strains sensitive or resistant to the bactericidal action of normal cord serum were appointed [7]. Interesting data were published for urinary tract pathogen *Proteus mirabilis* treated by the sequential passage in 90% normal bovine serum. Serum sensitive *P. mirabilis* O<sub>18</sub> strain after prolonged treatment by animal serum produced certain classes of surface proteins in excess, which were related in their quantity to these isolated from the resistant strain *P. mirabilis* O<sub>3</sub> [8]. Component C<sub>4</sub>bp regulates both CP and LP of C by preventing the assembly of C<sub>4</sub>b<sub>2</sub>a (the CP/LP C<sub>3</sub>

convertase) and accelerating the decay of this complex. Many pathogens are equipped with protein surface factors that bind C<sub>4</sub>bp. They are found for instance in bacterial species of *Neisseria meningitidis*, *Neisseria meningitidis*, *Neisseria gonorrhoeae*, and *Y. enterocolitica* [9-11]. The major OMP of *N. gonorrhoeae* is Por, which comprises 60% of the protein content of OM. The binding of C<sub>4</sub>bp to Por<sub>1</sub>A represents the C evasion mechanism [10]. It was demonstrated that serogroup B *N. meningitidis* that produce Pores also bind C<sub>4</sub>bp. The results of serum bactericidal assay showed that the strains expressing PorA are more resistant to C lysis than PorA-negative strains [9]. *Y. enterocolitica* has two classes of OMP: YadA and Ail, which act as receptors for C<sub>4</sub>bp. These proteins are the main SR determinants of these bacteria [11]. The fact is that the AP spontaneously initiates in serum. What is more, this pathway enhances C activation initiated by the CP and the LP. To protect own tissues organisms are equipped with membrane-bound C inhibitors and fluid-phase inhibitors. A significant, soluble-phase, the AP regulator is fH that act as cofactors for factor I in C<sub>3</sub>b inactivation; it competes with factor B for C<sub>3</sub>b binding and, for this reason, prevent C<sub>3</sub>bBb (the C<sub>3</sub> convertase of AP) formation. In addition, it accelerates the decay of C<sub>3</sub>bBb [12]. Factor H binding is a frequent mechanism used by different pathogens such as *Borrelia burgdorferi*, *N. gonorrhoeae*, and *Haemophilus influenzae* [12-14]. In turn, the virulence factor PgtE (outer membrane protease) of *Salmonella enterica* not only cleaves fH, but also other C components: factor B, C<sub>3</sub>, and C<sub>3</sub>b [15]. In the context of mobilization of C components of the distinct pathways, porins of *Klebsiella pneumoniae* were more effective activators of the CP. The molecule isolated from this pathogen named OmpK36 interacted directly with C<sub>1</sub>q complement protein independently of antibodies [16,17]. Former report suggested that OMP of molecular weights of 34 kDa and 36 kDa of *Salmonella Typhimurium* were involved in C activation. It has been shown that these porins enabled AP and CP pathways initiation [18]. Less information is available about the initiation of the LP by the presence of OMP. Lectin pathway begins with the binding of mannose-binding lectin (MBL) to repeating sugar moieties on the bacteria. Mannose-binding lectin was shown to bind to *N. meningitidis* surface proteins Opa and PorB, independent of the calcium ions [19].

### Acknowledgment

This work was supported by a "Academy of Development as the key to strengthening human resources of the Polish economy" Grant No. BPZ.506.50.2012.MS co-financed by the European Union under the European Social Fund.

### References

1. Rautemaa R, Meri S (1999) Complement-resistance mechanisms of bacteria. *Microbes Infect* 1: 785-794.

2. Futoma-Koloch B, Bugla-Ploskonska G, Sarowska J (2012) Searching for outer membrane proteins typical of serum-sensitive and serum-resistant phenotypes of *Salmonella*. InTech, Croatia.
3. Hallström T, Nordström T, Tan TT, Manolov T, Lambris JD, et al. (2011) Immune evasion of *Moraxella catarrhalis* involves ubiquitous surface protein A-dependent C3d binding. J Immunol 186: 3120-3129.
4. Futoma-Koloch B, Godlewska U, Guz-Regner K, Dorotkiewicz-Jach A, Klaus E, et al. (2015) Presumable role of outer membrane proteins of *Salmonella* containing sialylated lipopolysaccharides serovar Ngozi, sv. Isaszeg and subspecies Arizona in determining susceptibility to human serum. Gut Pathog 7: 18.
5. Drogari-Apiranthitou M, Kuijper EJ, Dekker N, Dankert J (2002) Complement activation and formation of the membrane attack complex on serogroup B *Neisseria meningitidis* in the presence or absence of serum bactericidal activity. Infect Immun 70: 3752-3758.
6. Skorek K, Raczowska A, Dudek B, Mińska K, Guz-Regner K, et al. (2013) Regulatory protein OmpR influences the serum resistance of *Yersinia enterocolitica* O:9 by modifying the structure of the outer membrane. PLoS One 8: e79525.
7. Bugla-Ploskonska G, Korzeniowska-Kowal A, Guz-Regner K (2011) Reptiles as a source of *Salmonella* O48-clinically important bacteria for children: the relationship between resistance to normal cord serum and outer membrane protein patterns. Microb Ecol 61: 41-51.
8. Futoma-Koloch B, Bugla-Ploskonska G, Doroszkiewicz W, Kaca W (2006) Survival of *Proteus mirabilis* O3 (S1959), O9 and O18 strains in normal human serum (NHS) correlates with the diversity of their outer membrane proteins (OMPs). Po J Microbiol 55: 153-156.
9. Jarva H, Ram S, Vogel U, Blom AM, Meri S (2005) Binding of the complement inhibitor C4bp to serogroup B *Neisseria meningitidis*. J Immunol 174: 6299-6307.
10. Ram S, Cullinane M, Blom AM, Gulati S, McQuillen DP, et al. (2001) Binding of C4b-binding protein to porin: a molecular mechanism of serum resistance of *Neisseria gonorrhoeae*. J Exp Med 193: 281-295.
11. Kirjavainen V, Jarva H, Biedzka-Sarek M, Blom AM, Skurnik M, et al. (2008) *Yersinia enterocolitica* serum resistance proteins YadA and ail bind the complement regulator C4b-binding protein. PLoS Pathog 4: 1000140.
12. Hellwage J, Meri T, Heikkilä T, Alitalo A, Panelius J, et al. (2001) The complement regulator factor H binds to the surface protein OspE of *Borrelia burgdorferi*. J Biol Chem 276: 8427-8435.
13. Ngampasutadol J, Ram S, Gulati S, Agarwal S, Li C, et al. (2008) Human factor H interacts selectively with *Neisseria gonorrhoeae* and results in species-specific complement evasion. J Immunol 180: 3426-3435.
14. Rosadini CV, Ram S, Akerley BJ (2014) Outer membrane protein P5 is required for resistance of nontypeable *Haemophilus influenzae* to both the classical and alternative complement pathways. Infect Immun 82: 640-649.
15. Riva R, Korhonen TK, Meri S (2015) The outer membrane protease PgtE of *Salmonella enterica* interferes with the alternative complement pathway by cleaving factors B and H. Front Microbiol 6: 63.
16. Alberti S, Alvarez D, Merino S, Casado MT, Vivanco F, et al. (1996) Analysis of complement C3 deposition and degradation on *Klebsiella pneumoniae*. Infect Immun 64: 4726-4732.
17. Alberti S, Marqués G, Camprubí S, Merino S, Tomás JM, et al. (1993) C1q binding and activation of the complement classical pathway by *Klebsiella pneumoniae* outer membrane proteins. Infect Immun 61: 852-860.
18. Galdiero F, Tufano MA, Sommese L, Folgore A, Tedesco F (1984) Activation of complement system by porins extracted from *Salmonella typhimurium*. Infect Immun 46: 559-563.
19. Estabrook MM, Jack DL, Klein NJ, Jarvis GA (2004) Mannose-binding lectin binds to two major outer membrane proteins, opacity protein and porin, of *Neisseria meningitidis*. J Immunol 172: 3784-3792.