

Bilateral Radial Artery Aneurysm: Case Report and Review of Literature

Thai H*, Vasyluk A and Rits Y

Vascular Surgery Fellow, Wayne State University SOM, Detroit Medical Center, USA

Abstract

While arterial aneurysm in the lower extremities is common, upper extremity aneurysms are relatively infrequent. Radial artery aneurysms are sometimes associated with trauma and connective tissue disorders. Very few cases of radial artery aneurysm have been reported in the literature. We have encountered a case of bilateral radial artery aneurysms in a 61 year old male. A review of the English language literature was performed using Pub Med database, resulting in 31 reported cases of radial artery aneurysms, with most cases being either traumatic or idiopathic in etiology. Of these reports, bilateral aneurysms had clear etiologies, including Marfan's vasculopathy, granulomatous arteritis, arteriosclerotic disease and intra-arterial drug injection. To our knowledge, this is the first reported case of primary bilateral radial artery aneurysms.

Keywords: Radial artery aneurysms; Peripheral aneurysms; Vasculitis

Case Report

A 61 year old left-hand dominant man with a history of frequent falls presented with bilateral hand pain and cellulitis. He has a history of falls in the weeks prior to presentation with redness, pain and swelling of his right wrist. Approximately a week later his left wrist and forearm also became swollen and erythematous. Physical exam revealed a moderately edematous left forearm with bilateral tender pulsatile masses at the wrists. Color flow duplex showed a maximum diameter of 1.3 and 1.5 cm of the left and right radial artery aneurysms, respectively, with intact palmar arches and bilateral radial dominance.

His past medical history includes hypertension and ischemic stroke two years ago. There was no indication from the history of connective tissue disease, and workup for vasculitis (sedimentation rate and C-reactive protein) was negative. A screening duplex of the popliteal arteries and CT of the chest and abdomen did not reveal other aneurysms.

Due to the size of the aneurysms and persistent symptoms operative management was discussed and the patient was taken to the operating room for subsequent aneurysm resection. The larger and more symptomatic aneurysm (right) side was resected first. Intraoperative findings showed significant inflammatory reaction and scar tissue encasing the aneurysm wall. There was an excellent doppler signal in the palmar and digits with the radial artery clamped. Thus we chose not to perform a bypass. A month later the patient returned for the left radial aneurysm resection. This aneurysm had septation and significant intramural thrombus. There was a radial artery dependent palmar circulation thus an interposition graft was performed with a reversed great saphenous vein. Patient experienced a full and uncomplicated recovery. Histological examination of the right aneurysm revealed fibromyxoid degeneration of the media with elastic tissue degeneration, while the left specimen showed increased perivascular inflammatory infiltrate but was not conclusive for vasculitis.

Discussion

True aneurysms of the radial artery are rare, with most reported being false aneurysms, traumatic or iatrogenic in origin. Given normal inflammatory markers and recent history of trauma, these aneurysms are likely secondary to trauma provoked. A search of English-language literature reveals 29 cases of true radial artery aneurysms,

mostly reported as single cases or case series. Tables 1 and 2 of these 4 was bilateral [1-4]. The most commonly etiologies were idiopathic and traumatic aneurysms, although it has been suggested that some of the cases of unknown etiology may be due to unrecalled trauma [3]. Other less common etiologies included mycotic aneurysms [5,6] atherosclerosis [7], NF-1 vasculopathy [8,9], vasculitis [3], and Marfan's syndrome [1,10]. The reported cases show a male predominance, with a mean age of 57 years, and most typically occur in the distal part of the artery, around the wrist and "snuffbox", where the artery is most superficial.

Most cases present as an asymptomatic swelling [2,7,11,12] but can present with paraesthesias due to radial nerve compression [13], symptoms of embolization [14] or rupture [8]. Diagnosis is often made on physical exam which shows a pulsatile mass, as well as non-invasive imaging such as doppler ultrasound. While most such aneurysms are traumatic or idiopathic in nature, efforts should be made to evaluate for vasculitis or connective tissue disease where appropriate, as well as screening for other aneurysms. Radial artery aneurysms are rare and its management is not well described. Early intervention is recommended to prevent distal embolization as well as addressing pain and compressive symptoms [15].

The most commonly reported approach for radial artery aneurysms is ligation and excision [5,12,13,16-23] which appears to be the preferred in cases where the ulnar artery is the dominant arterial supply for the hand [13,15]. In cases where the ulnar collaterals are poor or there is radial dominant circulation, options include excision and anastomosis [3,7,24] or excision with interposition vein graft [11,17] if a tension free anastomosis could not be accomplished, as in our case. A few reports exist of long term follow up for radial artery

***Corresponding author:** Thai H, Vascular Surgery Fellow, Wayne State University SOM, Detroit Medical Center, USA, Tel: 206-484-3664; E-mail: hcthai@gmail.com

Received June 10, 2015; **Accepted** January 21, 2016; **Published** January 29, 2016

Citation: Thai H, Vasyluk A, Rits Y (2016) Bilateral Radial Artery Aneurysm: Case Report and Review of Literature. *J Vasc Med Surg* 4: 247. doi:[10.4172/2329-6925.1000247](https://doi.org/10.4172/2329-6925.1000247)

Copyright: © 2016 Thai H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Author	Sex	Gender	Size/Location	Etiology	Treatment
Yukios [1]	74	Female	9 mm, 5 mm, snuffbox (bilateral)	Marfans	Ligation + excision (right)
Malt [2]	56	Male	2.0 cm, 1.5 cm, wrist (bilateral)	Arteriosclerotic	Resection + anastomosis (right)
Leitner [3]	69	Female	1.5 cm, 2.0 cm wrist (bilateral)	Granulomatous arteritis	Ligation + excision (bilateral)
Coppola [4]	40	Male	Wrist (bilateral)	Intrararterial drug injection	Ligation + excision (bilateral)
Shaabi [13]	65	Female	2 × 1.5 cm snuffbox	Idiopathic/non-traumatic	Ligation + excision
Luzzani [19]	63	Female	1.0 × 1.1 cm snuffbox	Idiopathic/non-traumatic	Ligation + excision
Yaghoubian [26]	77	Male	1.0 × 1.5 cm distal to snuffbox	Idiopathic/non-traumatic	Observation
Claudio [18]	47	Female	1.1 × 1.0 cm wrist	Idiopathic/non-traumatic	Ligation + excision
Walton [25]	40	Male	1.5 cm snuffbox	Idiopathic/non-traumatic	Observation
Turner [14]	55	Male	2.0 cm distal RA (antecubital fossa)	Idiopathic/non-traumatic	Resection + anastomosis
Filis [20]	45	Male	2.0 × 3.0 cm snuffbox	Idiopathic/non-traumatic	Ligation + excision
Jedynak [15]	60	Male	Snuffbox	Idiopathic/non-traumatic	Ligation + excision
Lee [24]	42	Male	Left wrist	Idiopathic/non-traumatic	Resection + anastomosis
Behar [21]	62	Male	1.9 cm wrist	Repetitive trauma	Ligation + excision
Turowski [11]	70	Female	2 cm × 7 cm wrist	Traumatic	Reconstruction with interposition vein graft
Kadowaki [5]	61	Male	2.0 × 1.5 cm snuffbox	Mycotic	Ligation + excision
Singh [9]	45	Male	Proximal radial artery	NF-1 vasculopathy	Ligation (no excision)
De Santis [8]	48	Female	Multiple proximal + distal aneurysms	NF-1 vasculopathy	Ligation + excision
Thorrens [7]	60	Male	1.5 × 2.0 cm snuffbox	Atherosclerotic	Resection + anastomosis
Goertz [10]	52	Male	Snuffbox	Marfans	Ligation + excision

Table 1: Size location/Etiology with multiple patients.

Author	Number pts	Location	Etiology	Management
Gray [12]	2	Distal radial artery/wrist	Repetitive trauma Idiopathic	Ligation + excision Observation
Johnson [6]	4	U/K	Mycotic	Ligation + excision
Mayall [22]	1(41M)	Distal radial artery/wrist	Trauma	Ligation + excision
Ho [17]	1	Wrist	Blunt trauma	Reconstruction w/ vein graft
Igari [16]	1 (72F)	Snuffbox	Idiopathic	Ligation + excision
Poirier [23]	1(69M)	Wrist	Mycotic	Ligation + excision

U/K = Unknown, N/A = Not Available

Table 2: Case series/reports with multiple patients.

reconstruction, ranging from 3 months to 5 years follow up, with most reporting no complications, including growth or recurrence of the aneurysm, but there is one published report of re-occurrence of aneurysm at 18 months [17]. We recommend pre-operative evaluation of palmar circulation with a digital doppler ultrasound.

In summary we recommend radial aneurysms to be repaired due to risks of embolization in addition to pain and compressive symptoms [25,26]. Preoperative laboratory testing should include a SED rate and CRP to exclude inflammatory components. Evaluation of the palmar circulation is necessary for operative planning. Treatment can include simple ligation, excision with primary anastomosis or with an interposition graft.

References

- Yukios U, Matsuno Y, Imaizumi M, Mori Y, Iwata H, et al. (2009) Bilateral radial artery aneurysms in the anatomical snuff box seen in marfan syndrome patient: case report and literature review. *Ann Vasc Dis* 2: 185-189.
- Malt S (1978) An arteriosclerotic aneurysm of the hand. *Arch Surg* 113: 762-763.
- Leitner DW, Ross JS, Neary JR (1985) Granulomatous radial arteritis with bilateral, nontraumatic, true arterial aneurysms within the anatomic snuffbox. *J Hand Surg Am* 10: 131-135.
- Coppola G, Amann-Vesti BR, Koppensteiner R (2003) Aneurysms of radial arteries and severe anemia. *Vasa* 32: 178.
- Kadowaki M, Hashimoto M, Nakashima M (2013) Radial mycotic aneurysm complicated with infective endocarditis caused by *Streptococcus sanguinis*. *Intern Med* 52: 2361-2365.
- Johnson JR, Ledgerwood AM, Lucas CE (1983) Mycotic aneurysm. New concepts in therapy. *Arch Surg* 118: 577-582.
- Thorrens S, Trippel OH, Bergan JJ (1966) Arteriosclerotic aneurysms of the hand. Excision and restoration of continuity. *Arch Surg* 92: 937-939.
- De Santis F, Negri G, Martini G, Mazzoleni G (2013) Multiple aneurysms of the radial artery in a woman with neurofibromatosis type 1 presenting as aneurysm rupture. *J Vasc Surg* 58: 1394-1397.
- Singh S, Riaz M, Wilmshurst AD, Small JO (1998) Radial artery aneurysm in a case of neurofibromatosis. *Br J Plast Surg* 51: 564-565.
- Goertz O, Kapalschinski N, Stricker I, Hauser J (2010) [Aneurysm of the radial artery in a patient with marfan syndrome]. *Handchir Mikrochir Plast Chir* 42: 307-309.
- Turowski GA, Amjadi N, Sterling A, Thomson JG (1997) Aneurysm of the radial artery following blunt trauma to the wrist. *Ann Plast Surg* 38: 527-530.
- Gray RJ, Stone WM, Fowl RJ, Cherry KJ, Bower TC (1998) Management of true aneurysms distal to the axillary artery. *J Vasc Surg* 28: 606-610.
- Shaabi HI (2014) True idiopathic saccular aneurysm of the radial artery. *J Surg Case Rep* 2014.
- Turner WH, Turnbull AR (1988) True radial artery aneurysm. *Br J Surg* 75: 233.
- Jedynak J, Frydman G (2012) Idiopathic True Aneurysm of the Radial Artery: A Rare Entity. *EJVES Extra* 24: e21-e22.
- Igari K, Kudo T, Toyofuku T, Jibiki M, Inoue Y (2013) Surgical treatment of aneurysms in the upper limbs. *Ann Vasc Dis* 6: 637-641.

17. Ho PK, Weiland AJ, McClinton MA, Wilgis EF (1987) Aneurysms of the upper extremity. *J Hand Surg Am* 12: 39-46.
18. Claudio NF, Ferreira MADA (2010) True aneurysm of the radial artery: a case report. *J Vasc Bras* 9: 239-240.
19. Luzzani L, Bellosta R, Carugati C, Talarico M, Sarcina A (2006) Aneurysm of the radial artery in the anatomical snuff box. *Eur J Vasc Endovasc Surg* 11: 94-96.
20. Filis K, Arhontovassilis F, Theodorou D, Theodossiades G, Manouras A (2007) True radial artery aneurysm in a mild haemophilia A patient. *Haemophilia* 13: 440-442.
21. Behar JM, Winston JS, Knowles J, Myint F (2007) Radial artery aneurysm resulting from repetitive occupational injury: Tailor's thumb. *Eur J Vasc Endovasc Surg* 34: 299-301.
22. Mayall JC, Mayall RC, Mayall AC, Mayall LC (1991) Peripheral aneurysms. *Int Angiol* 10: 141-145.
23. Poirier RA, Stansel HC Jr (1972) Arterial aneurysms of the hand. *Am J Surg* 124: 72-74.
24. Lee BY, Kim KK, Madden JL (1977) True aneurysm of the radial artery: report of a case with long-term follow-up. *J Am Geriatr Soc* 25: 376-378.
25. Walton NP, Choudhary F (2002) Idiopathic radial artery aneurysm in the anatomical snuff box. *Acta Orthop Belg* 68: 292-294.
26. Yaghoubian A, de Virgilio C (2006) Noniatrogenic aneurysm of the distal radial artery: a case report. *Ann Vasc Surg* 20: 784-786.

Citation: Thai H, Vasyluk A, Rits Y (2016) Bilateral Radial Artery Aneurysm: Case Report and Review of Literature. *J Vasc Med Surg* 4: 247. doi:[10.4172/2329-6925.1000247](https://doi.org/10.4172/2329-6925.1000247)

OMICS International: Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700 Open Access Journals
- 50,000 Editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus, Google Scholar etc.
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.editorialmanager.com/lifesciences