

## Bilateral Transcranial Direct-current Stimulation (tDCS) of Dorsolateral Prefrontal Cortex during Specific Working Memory Tasks

Saggini R<sup>1</sup>, Barassi G<sup>2</sup>, Carmignano SM<sup>3</sup>, Ancona E<sup>3</sup>, Di felice P<sup>4</sup>, Giannuzzo G<sup>4</sup>, Banchetti A<sup>5</sup> and Bellomo RG<sup>4</sup>

<sup>1</sup>Department of Medical Sciences, Oral and Biotechnology, Full Professor of PRM, "G.d'Annunzio" University, Chieti, Italy

<sup>2</sup>Faculty of Physiotherapy, PhD Lecturer, "G.d'Annunzio" University, Chieti, Italy

<sup>3</sup>School of Specialties in PRM, "G.d'Annunzio" University, Chieti, Italy

<sup>4</sup>Faculty of Physiotherapy, Lecturer "G.d'Annunzio" University, Chieti, Italy

<sup>5</sup>Physiotherapist, "G.d'Annunzio" University, Chieti, Italy

<sup>6</sup>Department of Medicine and Science of Aging, Associated Professor of PRM, "G.d'Annunzio" University, Chieti, Italy

\*Corresponding author: Giovanni Barassi, V.le Abruzzo 322, 66100 Chieti, Italy., Tel: +39 0871 3553006 – 3553007; E-mail: coordftgb@unich.it

Received date: August 03, 2016; Accepted date: August 04, 2016; Published date: August 08, 2016

Copyright: © 2016 Barassi G, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

Many authors analyzed the effect of direct current stimulation (tDCS), but the interest in tDCS effect on the human central nervous system for clinical application developed only in the recent years. Aim of current study is to evaluate the impact of tDCS on WM performance and better understand neuromodulatory effects of tDCS, in order to hypothesize applications for the treatment of memory deficits in stroke, Parkinson's and Alzheimer's diseases.

This study was conducted at the Centre of Physical and Rehabilitation Medicine, "Gabriele d'Annunzio" University in Chieti, in accordance with the Helsinki Declaration of 2013. 20 healthy subjects between 20 and 30 years old, 7 females and 13 males were enrolled. The population was divided into two groups; Group A, that underwent transcranial stimulation and Group B, that underwent sham stimulation. During the stimulation patients carry out specific test to assess working memory (Dual n-Back game). We performed 3 sessions in one week, every other day. An improvement in performance was recorded in both groups, statistically significant in the experimental group in 1 Back test. tDCS prefrontal stimulation can modulate working memory performance: further studies are required that would expand the search field and application of tDCS mainly in pathological conditions such as stroke and Parkinson's disease.

**Keywords:** Working memory; Transcranial direct-current stimulation; Rehabilitation working memory impairment

### Introduction

The interest *in vivo* of effects of transcranial direct current stimulation (tDCS) is widely spreading nowadays. Many authors have been study the effect of tDCS [1-4] but the interest in tDCS effects on the human central nervous system for clinical application has developed only in the recent years. tDCS is a non-invasive form of brain stimulation that involves the application of mild electrical currents to the scalp via large electrodes. tDCS alters cortex excitability through modulating, rather than directly stimulating, neuronal activity [5]. A small direct electrical current, typically 1-2 mA is induced across the brain from an anode to a cathode electrode [6] However, the effects of tDCS are not limited to modulations in cortical excitability during stimulation, and may outlast the stimulation period by several minutes or even hours [7-10]. These after effects of tDCS are associated with a number of different physiological modifications, including local changes in ionic concentrations (hydrogen, calcium) and levels of cyclic adenosine monophosphate (cAMP), alterations in protein synthesis, and modulation of N-methyl-D-aspartate (NMDA) receptor efficacy [11-17]. Thus increasing or decreasing the firing rate of the neurons depending upon their alignment with the current flow.

Recent studies use non-invasive brain stimulation (NIBS) techniques, such as repetitive transcranial magnetic stimulation (rTMS) [18-20] and tDCS [21,22], to increase dorsolateral prefrontal cortex (DLPFC) activity and, consequently, working memory (WM) performance. Working memory is the capacity to keep in mind some information for a short time to allow monitoring and manipulating information in short-term memory, providing the interface between perception, long-term memory and action that enables goal-directed behavior [23,24].

Some studies, thanks to the development of neuroimaging techniques, indicate that WM is localized across multiple regions in the brain particularly in the dorsolateral region of the prefrontal cortex [25,26].

Barbey et al. [27] showed that anatomical site of general intelligence and working memory is circumscribed, concentrated within the core of white matter fiber tracts that connect DLPFC with the inferior parietal cortex and that terminate in the superior parietal lobe. The observed reliance upon white matter fiber tracts suggests that working memory and other high-level cognitive processes are supported by the interregional communication among many brain areas, emphasizing the central role of the DLPFC and parietal cortex [28]. It has been demonstrated that 10 minutes [29] of 1 mA anodal tDCS, applied to the left DLPFC, increase verbal performance of a WM task completed during the last 5 minutes of stimulation, compared with sham stimulation. Previous studies of normal healthy people who had no

neurological abnormality were also carried out [15,30]. Furthermore tDCS-induced changes in excitability or tDCS application preceding learning processes have been shown to cause inverse or preventative effects on proceeding manipulations of neuronal excitability and synaptic plasticity [31].

Others reports of tDCS effects on WM suggest that enhancement is possible through anodal stimulation of the left the dorsolateral prefrontal cortex (DLPFC), a finding that makes sense given the tendency for anodal stimulation to enhance neuronal excitability [32,33] and the role of left DLPFC in WM [34].

Aim of current study is to evaluate the impact of tDCS on WM performance and better understand neuromodulatory effects of tDCS, train the use of tDCS device, in order to hypothesize applications for the treatment of memory deficits in stroke, Parkinson's and Alzheimer's diseases.

## Materials and Methods

This single-blind control procedure was conducted at the Centre of Physical and Rehabilitation Medicine, "Gabriele d'Annunzio" University in Chieti in accordance with the Helsinki Declaration of 2013. 20 healthy subjects between 20 and 30 years old, 7 females and 13 males were enrolled. The population was divided into two groups; Group A (10 subjects) underwent trans-cranial stimulation and Group B (10 subjects) underwent sham stimulation. These subjects had never performed tDCS before, they did not know their group membership and all participants signed an informed consent.

Exclusion criteria were: a history of epilepsy, metal prosthesis, pace maker implant, cardiac or neurologic disease and any medication therapy or pregnant. Each subject performed 3 sessions (T1-T2-T3) in one week every other day. In Group A, tDCS was transferred using saline-soaked electrode in buckskin (35 cm<sup>2</sup>) and it was delivered using a battery-driven, constant-current stimulator HDCstim system (Newronika EN-ITA 00 2010).


During left DLPFC stimulation, the active electrode was over F3 and the reference above the right side- F3-F4 according to the 10–20 international system for electroencephalogram electrode placement [35-44]; the intensity was set at 1.5 mA (starter peak time of 5 sec) for a total time of 13 minutes. Intensity of electric current was kept under safety thresholds. Sham stimulation (Group B) was conducted with 120 s of tDCS applied at the onset, the same electrode placement was used (F3-F4 according to the 10–20 international system for electroencephalogram electrode placement), after 120 sec the device was programmed to stop.

During session subjects carry out after a Dual N-Back game test which is commonly used to study working memory.

The Dual N-Back game involves remembering a sequence of spoken letters and a sequence of positions of a square at the same time, and identifying when a letter or position matches the one that appeared earlier [35-39].

For the current study dual N-Back test was administered on a personal computer and structured in to two levels of difficulty: 1-Back first level and 2-Back second level. In 1-Back test 24 visual and auditory stimuli were provided, each one lasted 1, 5 sec with a gap between one stimulus and the next one of 2, 3 sec, for a total time of 91 sec. 1 Dual-Back: an auditory stimulus (a,b,c etc.) and simultaneously one visual-spatial stimulus consists of a blue square that changes

position within a 9-panes grid is presented to the subject. The subject is instructed to press a button when there is a match is in position, the square reappears in a position where it was previously, that sound, when repeating a letter already heard earlier. 2-Back test had three kinds of stimuli: a spatial visual, simultaneously visual and an auditory stimulation. Visual spatial stimulus consisted in an image that, with each apparition, changed its position in a space grid, the visual stimulation consisted in the same image that changed shape; the auditory stimulation consisted of an alphabetic letter. Dual 2-Back: it has the same characteristics of the Dual n 1-Back, but with additional difficulty. The subject does not see the square that changes position but a figure; it may introduce a car, a boat, a plane, a star etc., which changes position and shape. So subjects have to remember: the earlier listened letters, position, and figure previously appeared. Stimuli provided were 24; each one lasted for 1.5 sec with latency between one and the other of 2.3 sec, for a total time of 91 sec. (Figure 1)



**Figure 1:** Timeline and sequence of the session both for group A that group B.


During the first session all participants were instructed on the use of n-Back test with 48 total trial stimuli between level 1 Back and level 2 Back tests, these results were not considered for data analysis. After 8 minutes the subjects underwent dual n-Back test and this time data was considered. During the second and the third session subjects did not undergo training but they directly did 1 and 2 Back test. At each session were recorded the obtained scores (number of correct answers) is 1 to 2 for the Back which Back testing.

## Statistical analysis

Data are analysed by Wilcoxon-Mann-Whitney test with NCSS9<sup>®</sup> software and one way Anova for Windows. The minimum level of significance was set to  $p < 0.05$ .

## Results

In group A for the 1 Back: the average number of correct answers (indicated as a score) has significant increased from T1 ( $51.8 \pm 21.61$ ) to T2 ( $63.4 \pm 24.38$ )  $p=0.04883$  ( $p < 0.05$ ). The increase is confirmed to T3 compared to T1 ( $71.5 \pm 21.9$ )  $p=0.045$  ( $p \leq 0.05$ ). Instead, B group (sham group) shows decrease in the average score from T1 ( $55 \pm 21.9$ ) to T2 ( $33.2 \pm 22.2$ )  $p=0.068$  and at T3 ( $67 \pm 27.12$ ) the increase is not significant (Figure 2).




**Figure 2:** Average score trend in 1 Back at T1-T2-T3 in Group A and Group B.

There was a significant effect of tDCS evaluated with one way ANOVA as shown in Table 1.

ANOVA					
Anova one way	d.f.	SS	MS	F	P value
Between groups	5	3.187,3333	637.4667	2.5093	0.0409
Intra group	54	13.718	254.037		
Total	59	16.905,3333			

**Table 1:** One way Anova for 1 Back. df degrees of freedom; SS deviance; MS variance; F test F.

Average of 2 Back scores: the group A showed a non-significant increase in T2 to T3 (T1 50.2 ± 13.4, T2 55.7 ± 13.8, T3 p=0.89791). In group B: there was a stabilization of the averages of scores to T2 (T1 46.1 ± 21.39, T2 46.7 ± 15.4) and an increase at T3 (63.4 ± 15.6) that is not significant p=0.013 (Figure 3).



**Figure 3:** Average score trend in 2 Back at T1-T2-T3 in Group A and Group B.

These data are confirmed by the analysis of variance with one way ANOVA (Table 2).

ANOVA					
Anova one way	SS	df	MS	F	p value
Between groups	937.4	2	468.7	1.105	0.3457
Intra group	11.452,9	27	424.1815		
Totale	12.390,3	29			

**Table 2:** One way Anova for 2 Back. df degrees of freedom; SS deviance; MS variance; F test F.

The improvements in Group A are shown to be steady and homogeneous, meanwhile the improvement in group B seem to have a spike in T3 probably two to the training effect.

## Discussion

The selective influence of tDCS on different cognitive processes, after the bilateral application has already been amply demonstrated in the primary motor cortex [40], in the prefrontal cortex [41,42], in the anterior temporal lobe, in the fronto-temporal region [43] and posterior parietal cortex [30].

Ohn et al. [10] reported that a 20-or 30-min anodal stimulation boosted short-term memory performance significantly more than 10 min of stimulation did, demonstrating that the effects of tDCS intensify with a longer stimulation period. Lally et al. [44] confirmed these results in a larger cohort, but only when subjects were tested during the stimulation session (online), without a persisting effect 48 h later. Marshall et al. [45] reported increased reaction time in the same task during both anodal and cathodal bilateral intermittent stimulation over the DLPFC. Moreover, Andrews et al. [46] investigated the impact of 1 session of anodal tDCS delivered during a WM task (n-Back task) on performances on a subsequent WM task (digit span forward). Upon completion of the n-Back task, they observed a significant improvement in performance on the digit span forward task. In Javadi and Walsh's [47] and Javadi et al. [48] studies, anodal tDCS that was applied during initial encoding enhanced performance on a subsequent word memory task. This raises the possibility that anodal tDCS used during initial encoding can be combined with anodal tDCS administered during subsequent consolidation in order to bring larger memory improvements to people. Alonzo et al. [49] explored the effects of anodal stimulation (0.0571 mA/cm<sup>2</sup> current density; M1/contralateral orbit montage; 20 min duration) on MEP amplitude over the course of 5 days (Monday-Friday). Although these researchers reported variable baseline levels across the week, the ratio of pre- to post-stimulation group average MEP amplitudes did not significantly change from day-to-day. Using a similar protocol, Gálvez et al. [50] reported similar findings: namely, whereas baseline levels changed throughout the week, the group averaged after-effects of daily stimulation did not significantly vary across 5 days. Jean et al. [51] investigated the effect (tDCS) applied over the prefrontal cortex on the improvement of verbal, visuo-spatial working memory and naming in healthy adults. The subjects were divided into four groups randomly and they underwent sham or anodal tDCS over the left or right prefrontal cortex, for 20 minutes at a direct current of 1 mA. The word/interference significantly improved in the left prefrontal tDCS group compared with that of the sham group while the visuospatial attention

significantly increased in the right prefrontal tDCS group compared with that of the sham group. tDCS can induce verbal working memory improvement and naming facilitation by stimulating the left prefrontal cortex. Donckery et al. [52] in a study on healthy subjects showed that tDCS is able to increase planning performance and cognitive skill learning. This study concluded that excitability decreasing cathodal tDCS mediates its beneficial effect through early noise reduction of neuronal activity. Nelson et al. [53] have shown the efficacy of tDCS in a study in which participants performed a task of air traffic control simulation requiring detect possible collision paths.

Tremblay et al. [54] show that the position of electrodes determine different effects. Previous studies have shown that stimulation with bilateral assembly determines an increase of mental flexibility: [55], language comprehension, generation of untruthful answer [56], attention and language performance [57], automaticity for learned materials [58].

One possible way to reduce uncertainty about effect of tDCS is to monitor the brain impact of tDCS separately and independently of behavioural and cognitive effects. Techniques such as EEG, TMS-EEG, magnetic resonance spectroscopy, functional magnetic resonance imaging and modelling of induced currents have all been shown to be effective in characterizing the physiologic effects of tDCS.

We only performed stimulation and test for only 1 week, if we performed more follow up tests with different periods, we could have detected persistent durations of tDCS effects and the changing patterns of each test more accurately. If you replicated with a larger sample, more numerous sessions and with a follow up, the data may eventually be confirmed, with potential long-term effects that also extend in the hours or days following the experimental session. Moreover we must remind that the subjects in exam are not from a clinical population, thus making the individual differences more relevant, and the number of patients is not huge so we cannot make a statement that can be valid in general. However, more research needs to be conducted to examine if the beneficial effects of tDCS during encoding and reactivation are additive, or if there is a limit on the extent to which tDCS can enhance memory performance.

Concluding our data could be explained according to Teo et al. [59] and Hoy et al. [60] could be that the greater cortical excitability needed to perform the n-back task with its consistent WM load resulted in a cumulative effect above which the cortex is not over excited in the last part of tDCS stimulation session.

Furthermore the idea of the study is also born to learn more about the use of the machine and then start trials in those diseases characterized by an impairment of working memory. WM impairment is a major feature of a number of neurologic and psychiatric disorders, including schizophrenia, Alzheimer's disease, Parkinson's disease, and major depression [61-64]. Some studies have indicated that antipsychotics, particularly the newer atypical antipsychotics, may improve cognitive functioning in schizophrenia [65,66]. Despite this, no effective treatment has yet been established so new and thorough research will be needed in this field.

## References

1. Bindman LJ, Lippold OCJ, Redfearn JWT (1964) the action of brief polarizing currents on the cerebral cortex of the rat 1 during current flow and 2 in the production of long-lasting after-effects. *J Physiol* 172: 369-382.
2. Elsberg Ca (1917) Experiments on Motor Nerve Regeneration and the Direct Neurotization of Paralyzed Muscles by their Own and by Foreign Nerves. *Science* 45: 318-320.
3. Fuortes MG (1954) Direct current stimulation of motoneurons. *J Physiol* 126: 494-506.
4. Hern JE, Landgren S, Phillips CG, Porter R (1962) Selective excitation of corticofugal neurones by surface-anodal stimulation of the baboon's motor cortex. *J Physiol* 161: 73-90.
5. Nitsche MA, Cohen LG, Wassermann EM, Priori A, Lang N, et al. (2008) Transcranial direct current stimulation: State of the art 2008. *Brain Stimul* 1: 206-223.
6. Floel A, Cohen LG (2007) Contribution of noninvasive cortical stimulation to the study of memory functions. *Brain Res Rev* 53: 250-259.
7. Nitsche MA, Nitsche MS, Klein CC, Tergau F, Rothwell JC, et al. (2003) Level of action of cathodal DC polarisation induced inhibition of the human motor cortex. *Clinical Neurophysiology* 114: 600-604.
8. Nitsche MA, Paulus W (2000) Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation. *The Journal of Physiology* 527: 633-639.
9. Bindman LJ, Lippold OC, Redfearn JW (1962) Long-lasting changes in the level of the electrical activity of the cerebral cortex produced by polarizing currents. *Nature* 196: 584-585.
10. Ohn SH, Park CI, Yoo WK, Ko MH, Choi KP, et al. (2008) Time-dependent effect of transcranial direct current stimulation on the enhancement of working memory. *Neuroreport* 19: 43-47.
11. Ardolino G, Bossi B, Barbieri S, Priori A (2005) Non-synaptic mechanisms underlie the after-effects of cathodal transcutaneous direct current stimulation of the human brain. *J Physiol* 568: 653-663.
12. Gartside IB (1968) Mechanisms of sustained increases of firing rate of neurons in the rat cerebral cortex after polarization: reverberating circuits or modification of synaptic conductance? *Nature* 220: 382-383.
13. Hattori Y, Moriwaki A, Hori Y (1990) Biphasic effects of polarizing current on adenosine-sensitive generation of cyclic AMP in rat cerebral cortex. *Neuroscience Letters* 116: 320-324.
14. Islam N, Aftabuddin M, Moriwaki A, Hattori Y, Hori Y (1995) Increase in the calcium level following anodal polarization in the rat brain. *Brain Res* 684: 206-208.
15. Liebetanz D, Nitsche MA, Tergau F, Paulus W (2002) Pharmacological approach to the mechanisms of transcranial DC-stimulation-induced after-effects of human motor cortex excitability. *Brain* 125: 2238-2247.
16. Nitsche MA, Jaussi W, Liebetanz D, Lang N, Tergau F, et al. (2004) Consolidation of human motor cortical neuroplasticity by D-cycloserine. *Neuropsychopharmacology* 29: 1573-1578.
17. Wagner T, Valero-Cabre A, Pascual-Leone A (2007) Noninvasive human brain stimulation. *Annual Review of Biomedical Engineering* 9: 527-565.
18. Dayan E, Censor N, Buch ER, Sandrini M, Cohen LG (2013) Noninvasive brain stimulation: from physiology to network dynamics and back. *Nat Neurosci* 16: 838-844.
19. Demirtas-Tatlidede A, Vahabzadeh-Hagh AM, Pascual-Leone A (2013) Can noninvasive brain stimulation enhance cognition in neuropsychiatric disorders? *Neuropharmacology* 64: 566-578.
20. Fregni F, Pascual-Leone A (2007) Technology insight: Noninvasive brain stimulation in neurology-perspectives on the therapeutic potential of rTMS and tDCS. *Nature Clinical Practice Neurology* 3: 383-393.
21. Heimrath K, Sandmann P, Becke A, Muller NG, Zaehle T (2012) Behavioral and electrophysiological effects of transcranial direct current stimulation of the parietal cortex in a visuo-spatial working memory task. *Front Psychiatry* 3: 56.
22. D'Esposito M, Postle BR, Rypma B (2000) Prefrontal cortical contributions to working memory: Evidence from event-related fMRI studies. *Exp Brain Res* 133: 3-11.
23. Baddeley AD, Thomson N, Buchanan M (1975) Word length and the structure of short-term memory. *J Verbal Learn. Verbal Behav* 14: 575-589.



24. Baddeley A, Petrides M (1996) Specialized systems for the processing of mnemonic information within the primate frontal cortex. *Philos Trans R Soc Lond B Biol Sci* 351: 1461-1462.
25. Cabeza R, Nyberg L (2000) Neural bases of learning and memory: functional neuroimaging evidence. *Curr Opin Neurol* 13: 415-421.
26. Daniel TA, Katz JS, Robinson JL (2016) Delayed match-to-sample in working memory: A BrainMap meta-analysis. *Biol Psychol* 120: 10-20.
27. Barbey AK, Colom R, Solomon J, Krueger F, Forbes C, et al. (2012) An integrative architecture for general intelligence and executive function revealed by lesion mapping. *Brain* 135: 1154-1164.
28. Jung RE, Haier RJ (2007) The Parieto-Frontal Integration Theory (P-FIT) of intelligence: converging neuroimaging evidence. *Behav Brain Sci* 30: 135-135.
29. Fregni F, Boggio PS, Nitsche M, Bermanpohl F, Antal A, et al. (2005) Anodal transcranial direct current stimulation of prefrontal cortex enhances working memory. *Exp Brain Res* 166: 23-30.
30. Cohen Kadosh R, Soskic S, Iuculano T, Kanai R, Walsh V (2010) Modulating neuronal activity produces specific and long-lasting changes in numerical competence. *Curr Biol* 20: 2016-2020.
31. Lang N, Siebner HR, Ernst D (2004) Preconditioning with transcranial direct current stimulation sensitizes the motor cortex to rapid-ratetranscranial magnetic stimulation and controls the direction of after-effects. *Biol Psychiatry* 56: 634-639.
32. Rosenkranz K, Nitsche MA, Tergau F, Paulus W (2000) Diminution of training-induced transient motor cortex plasticity by weak transcranial direct current stimulation in the human. *Neurosci Lett* 296: 61-63.
33. Priori A (2003) Brain polarization in humans: a reappraisal of an old tool for prolonged non-invasive modulation of brain excitability. *Clin Neurophysiol* 114: 589-595.
34. Carpenter PA, Just MA, Reichle ED (2000) Working memory and executive function: evidence from neuroimaging. *Curr Opin Neurobiol* 10: 195-199.
35. Sandrini M, Fertonani A, Cohen LG, Miniussi C (2012) Double dissociation of working memory load effects induced by bilateral parietal modulation. *Neuropsychologia* 50: 396-402.
36. Poreisz C, Boros K, Antal A, Paulus W (2007) Safety aspects of transcranial direct current stimulation concerning healthy subjects and patients. *Brain Research Bulletin* 72: 4-6.
37. Gevins A, Cuttillo B (1993) Spatiotemporal dynamics of component processes in human working memory. *Electroencephalogr Clin Neurophysiol* 87: 128-143.
38. Carter CS, Braver TS, Barch DM, Botvinick MM, Noll D, et al. (1998) Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* 280: 747-749.
39. Smith EE, Jonides J (1998) Neuroimaging analyses of human working memory. *Proc Natl Acad Sci U S A* 95: 12061-12068.
40. Cerruti C, Schlaug G (2009) Anodal transcranial direct current stimulation of the prefrontal cortex enhances complex verbal associative thought. *J Cogn Neurosci* 21: 1980-1987.
41. Hecht D, Walsh V, Lavidor M (2010) Transcranial direct current stimulation facilitates decision making in a probabilistic guessing task. *J Neurosci* 30: 4241-4245.
42. Fecteau S, Knoch D, Fregni F, Sultani N, Boggio P, et al. (2007) Diminishing risk-taking behavior by modulating activity in the prefrontal cortex: a direct current stimulation study. *J Neurosci* 27: 12500-12505.
43. Penolazzi B, Di Domenico A, Marzoli D, Mammarella N, Fairfield B, et al. (2010) Effects of Transcranial Direct Current Stimulation on episodic memory related to emotional visual stimuli. *PLoS One* 5: e10623.
44. Lally N, Nord CL, Walsh V, Roiser JP (2010) Does excitatory fronto-extracerebral tDCS lead to improved working memory performance? *Version 2 F1000Res* 2: 219.
45. Marshall L, Mölle M, Siebner HR, Born J (2005) Bifrontal transcranial direct current stimulation slows reaction time in a working memory task. *BMC Neurosci* 6: 23.
46. Andrews SC, Hoy KE, Enticott PG, Daskalakis ZJ, Fitzgerald PB (2011) Improving working memory: the effect of combining cognitive activity and anodal transcranial direct current stimulation to the left dorsolateral prefrontal cortex. *Brain Stimul* 4: 84-89.
47. Javadi AH, Walsh V (2012) Transcranial direct current stimulation (tDCS) of the left dorsolateral prefrontal cortex modulates declarative memory. *Brain Stimul* 5: 231-241.
48. Javadi AH, Cheng P, Walsh V (2012) Short duration transcranial direct current stimulation (tDCS) modulates verbal memory. *Brain Stimul* 5: 468-474.
49. Alonzo A, Brassil J, Taylor JL, Martin D, Loo CK (2012) Daily transcranial direct current stimulation (tDCS) leads to greater increases in cortical excitability than second daily transcranial direct current stimulation. *Brain Stimulation* 5: 208-213.
50. Gálvez V, Alonzo A, Martin D, Loo CK (2013) Transcranial direct current stimulation treatment protocols: should stimulus intensity be constant or incremental over multiple sessions? *Int J Neuropsychopharmacol* 16: 13-21.
51. Jeon SY, Han SJ (2012) Improvement of the working memory and naming by transcranial direct current stimulation. *Ann Rehabil Med* 36: 585-595.
52. Dockery CA, Hueckel-Weng R, Birbaumer N, Plewnia C (2009) Enhancement of planning ability by transcranial direct current stimulation. *J Neurosci* 29: 7271-7277.
53. Nelson JT, McKinley RA, Goloba EJ, Warmc JS, Parasuramand R (2014) Enhancing vigilance in operators with prefrontal cortex transcranial direct current stimulation (tDCS). *Neuroimage* 85: 909-917.
54. Tremblay S, Lepage JF, Latulipe-Loiselle A, Fregni F, Pascual-Leone A, et al. (2014) The uncertain outcome of prefrontal tDCS. *Brain Stimul* 7: 773-783.
55. Leite J, Carvalho S, Fregni F, Boggio PS, Gonçalves OF (2012) The effects of crosshemispheric dorsolateral prefrontal cortex transcranial direct current stimulation (tDCS) on task switching. *Brain Stimul* 5: 231e4.
56. Nozari N, Thompson-Schill SL (2013) More attention when speaking: does it help or does it hurt? *Neuropsychologia* 51: 2770-2780.
57. Mamelì F, Mrakic-Spota S, Vergari M (2010) Dorsolateral prefrontal cortex specifically processes general e but not personal e knowledge deception: multiple brain networks for lying. *Behav Brain Res* 211: 164e8.
58. Fecteau S, Boggio P, Fregni F, Pascual-Leone A (2013) Modulation of untruthful responses with non-invasive brain stimulation. *Front Psychiatry* 3: 1e9.
59. Teo F, Hoy KE, Daskalakis ZJ, Fitzgerald PB (2011) Investigating the Role of Current Strength in tDCS Modulation of Working Memory Performance in Healthy Controls. *Front Psychiatry* 2: 45.
60. Hoy KE, Arnold SL, Emonson MR, Daskalakis ZJ, Fitzgerald PB (2014) An investigation into the effects of tDCS dose on cognitive performance over time in patients with schizophrenia. *Schizophr Res* 155: 96-100.
61. Goldman-Rakic PS (1994) Working memory dysfunction in schizophrenia. *J Neuropsychiatry Clin Neurosci* 6: 348-357.
62. Morris RG (1994) Working memory in Alzheimer-type dementia. *Neuropsychol* 8: 544-554.
63. Winograd-Gurvich C, Fitzgerald PB, Georgiou-Karistianis N, Bradshaw JL, White OB (2006) Negative symptoms: a review of schizophrenia, melancholic depression and Parkinson's disease. *Brain Res Bull* 70: 312-321.
64. Marini P, Ramat S, Ginestroni A, Paganini M (2003) Deficit of short-term memory in newly diagnosed untreated parkinsonian patients: reversal after L-dopa therapy. *Neurol Sci* 24: 184-185.
65. Lindenmayer JP, Khan A, Iskander A, Abad MT, Parker B (2007) A randomized controlled trial of olanzapine versus haloperidol in the treatment of primary negative symptoms and neurocognitive deficits in schizophrenia. *J Clin Psychiatry* 68: 368-379.
66. Riedel M, Spellmann I, Strassnig M, Douhet A, Dehning S, et al. (2007) Effects of risperidone and quetiapine on cognition in patients with schizophrenia and predominantly negative symptoms. *Eur Arch Psychiatry Clin Neurosci* 257: 360-370.